

Molecular Genetics of Aichivirus C (Porcine Kobuvirus) in China

Lei Chen¹, Ling Zhu¹, Zhi-wen Xu^{1*}, Wan-zhu Guo¹

¹Animal Biotechnology Center, College of Veterinary Medicine of Sichuan Agricultural University, 46[#] Xinkang Road, Yucheng District, Ya'an, 625014, Sichuan province of P.R. China

*Corresponding author: Zhi-wen Xu

E-mail address:

LC: morn_chen@126.com (Contacting author)

LZ: abtczl72@126.com

ZWX: abtcxzw@126.com

WZG: wzguo@126.com

Abstract

Aichivirus C is a member of the *Kobuvirus* genus within the *Picornaviridae* family and is widely distributed in both healthy and diarrheic pigs from China. Due to the high detection rate in the severe diarrhea in China, *Aichivirus C* is thought to be a potential pathogen of pig diarrhea. The review represented the discovery of *Aichivirus C* in China, and made a brief summary about molecular and epidemiology characterizations of Chinese *Aichivirus C* strains. Multiple *Aichivirus C* strains and *Aichivirus C* variants are circulating in China. Recombination events were also observed in Chinese *Aichivirus C* strains. More further studies are needed to clarify the evolutionary features and pathogenicity of *Aichivirus C*.

Introduction

Members of the *Picornaviridae* family are small, non-enveloped viruses with a linear, single-stranded, and positive-sense RNA genome. This family includes important pathogens of both human and animals and is divided into 17 genera: *Enterovirus*, *Aphthovirus*, *Cardiovirus*, *Hepatovirus*, *Parechovirus*, *Erbovirus*,

Teschovirus, *Avihepatovirus*, *Aquamavirus*, *Cosavirus*, *Dicipivirus*, *Megrivirus*, *Salivirus*, *Sapelovirus*, *Senecavirus*, *Tremovirus*, and *Kobuvirus*. The *Kobuvirus* genus contains three officially recognized species: *Aichivirus A* (human *Aichi virus*), *Aichivirus B* (*Bovine kobuvirus*), and *Aichivirus C* (*Porcine kobuvirus*) [1]. The RNA genomes of kobuviruses range from 8.2–8.4 kb, which include a 5'UTR, a leader (L) protein, three structural proteins (VP0, VP3, and VP1), seven non-structural proteins (2A, 2B, 2C, 3A, 3B and 3D), a 3'UTR and a poly (A) tail [2]. The *Kobuvirus* has a wide range of host specificity and kobuviruses have been detected in humans, cattle, pigs, sheep, wild boars, bats, dogs, cats, goats and rodents [3-12]. The *Aichivirus A* was first isolated in Japan from faecal samples of a patient suffering from acute gastroenteritis in 1991 and the *Aichivirus B* was detected in faecal samples from clinically healthy cattle in 2003 in Japan [3, 4]. The *Aichivirus C* was first reported in faecal samples of pig collected from a Hungarian farm in 2007, and later in 2009. Yu et al. have reported the first detection of *Aichivirus C* in China [5, 13]. Since then, *Aichivirus C* has been detected in Thailand, Spain, Japan, Korea, the United States, Brazil, the Netherlands, and recently in the Czech Republic [13-21].

Researchers from worldwide have done much work on *Aichivirus C* to clarify its epidemic characteristics, distribution, evolutionary features and genome sequences. Among all those currently published reports about *Aichivirus C*, most of which described the infection situations in China. Chinese scholars from different districts have carried out many approaches to determine the prevalence and molecular characterizations of *Aichivirus C* in China.

Discovery of *Aichivirus C* in China

When detecting calicivirus in porcine fecal specimens from healthy piglets by reverse transcription-PCR (RT-PCR), an unexpected nonspecific band was observed on agarose gel electrophoresis. Nucleotide sequence of this 1,185bp fragment was determined, and further molecular and phylogenetic analysis revealed that the porcine faecal specimen contained a novel species of *Kobuvirus*. This was the first detection of *Aichivirus C* in China [13]. The complete nucleotide sequence of *Aichivirus C*

strain Y-1-CHINA/2008 isolated from China was released by Yu et al. 2009. The isolate Y-1-CHI was 88.62%, 58.66%, and 48.86% identical to those of *Aichivirus C* (strain S-1-HUN), *Aichivirus B* (strain U-1), and *Aichivirus A* (strain A846/88), respectively. Phylogenetic tree revealed that Y-1-CHI and S-1-HUN belong to the same species, *Aichivirus C* [22].

Epidemiology of *Aichivirus C* in China

The *Aichivirus C* is widely distributed in China, in both apparently healthy and diarrheic pigs. Up to now, 28/34 (82.4%) first-level administrative subdivision districts have been reported the existence of *Aichivirus C* in porcine intestine and faecal samples (**Figure 1**) [13, 23-31].

In different regions, the prevalence rate of *Aichivirus C* ranged from 11.2 to 100% (Table 1) [13, 23-31]. Total infection rate of *Aichivirus C* was extremely high in a few provinces: 100% (8/8) in Anhui, 91.7% (11/12) in Henan, 88.1% (37/42) in Hunan, 96.2% (25/26) in Neimenggu, 84.2% (16/19) in Xinjiang. In addition, statistical analysis has concluded that *Aichivirus C* infection was more susceptible to young animals and pigs with diarrhoea [31, 32].

The *Aichivirus C* has been reported in pig serum in Hungary and Brazil [14, 33]. In China, pig serum from Gansu (33.3%, 2 out of 6) and also from Hunan, Hubei and Henan (total infection rate of those three provinces: 6.5%, 6 out of 92) have been tested for infection with *Aichivirus C*. Furthermore, recently, the *Aichivirus C* was detected in sow colostrum samples by Xiang et al. 2013 from IVDC (China Institute of Drug Control): 66.7% (2/3) in Beijing, 80% (4/5) in Xinjiang, and 84.2% (16/19) in Neimenggu [30].

Molecular genetics of *Aichivirus C* in China

There are total 11 *Aichivirus C* complete genome sequences available in GenBank from China. The complete RNA genomes of *Aichivirus C*, WUH1, Y-1-CHI, SH-W-CHN, K-11/2012/CH and K-4/2012/CH strains, were similar to the sequence of *Aichivirus C* prototype strain S-1-HUN, which consist of 8210 nucleotides (nt) in length excluding the poly(A) tail. The genome sequence of S-1-HUN shared 89.2%,

88.6%, 89%, 86% and 89% identity at the nucleotide level to WUH1, Y-1-CHI, SH-W-CHN, K-11/2012/CH and K-4/2012/CH, respectively [22, 23, 27, 34]. Interestingly, compared to the standard *Aichivirus C* strain S-1-HUN, there were six *Aichivirus C* variants in China. The CH/HNXX-4/2012 had a 30-amino-acid deletion in the 2B protein and a threonine amino acid insertion in the VP1 gene [35]. The genome sequence of CH/HZ is 8101 nt (excluding the poly-A tail); an adenine insertion at position 124, an 18-nt deletion in the 5'UTR, and a 90-nt deletion in the 2B coding region were also observed [25]. The RNA genomes of GS-1/2012/CH and GS-2/2012/CH consist 8121 nt, excluding the poly (A) tail, and they both have a 90-nucleotide deletion in the 2B protein and a single nucleotide insertion in the 3'UTR. Besides, possible 3C/3D cleavage site of GS-1/2012/CH and GS-2/2012/CH is Q/C, while it is Q/S in S-1-HUN genome sequence [27]. The *Aichivirus C* isolates XX and swKoV CH441 have genomes of 8147 nt and 8149 nt, respectively, which were shorter than the S-1-HUN (8210 nt).

Partial sequences of *Aichivirus C* 3D gene and VP1 gene from China were also largely characterized. The 3D RNA-dependent RNA polymerase gene is a relatively conserved region and has been used to determine phylogenetic relationships among kobuviruses [2]. The VP1 region is the most exposed and immunodominant portion of the capsid protein, and is the most variable structural protein among all kobuviruses [2, 3]. Characterizations of the 3D protein gene and VP1 gene play an important role in the molecular epidemiology and genetic evolution of *Aichivirus C*.

Phylogenetic trees based on the partial 3D or VP1 sequences revealed that there were multiple lineages of *Aichivirus C* circulating in China. In addition, Chen et al. 2013 have recently reported co-infection of multiple *Aichivirus C* strains in a single pig, and significant recombination breakpoints observed in the co-infection strains demonstrated that multiple *Aichivirus C* strains in the same pig could have arisen from recombination events [31]. Wang et al. 2012 indicated that early recombination events might have facilitated the genome of SH-W-CHN [23]. Significant recombination signals were also detected in GS-1/2012/CH [27]. Amino acid mutations were observed in both structural proteins and non-structural proteins.

Mutations and recombination events may have contributed to the genetic diversity of *Aichivirus C* and act as a driving force for the evolution of virus genomes. However, such finding requires further investigations.

Discussion and Summary

Since the end of 2010, massive outbreaks of diarrhoea have occurred in suckling piglets in China; however, the etiological agent has yet to be determined. Affected pigs exhibited signs of watery diarrhoea, dehydration, and vomiting with morbidity ranging 80–100% and mortality between 50–90%. High detection rate of *Aichivirus C* in those porcine diarrhoea samples made researchers suspect that *Aichivirus C* may be a causative agent of the diarrhoea or has atleast contributed to aggravate the disease.

Failure of culturing *Aichivirus C* in vitro has hindered the research about its pathogenicity. Considering *Aichivirus C* is also widely existent in healthy swinery, most people probably would ignore its importance. Less intentions have been paid to these asymptomatic infections, providing suitable conditions for evolution and persisting of *Aichivirus C* in pig hosts. Recombination events and variants are common in Chinese *Aichivirus C* group. Recombination and mutation possibly could convert *Aichivirus C* to a potential causative agent. We could not exclude the possibility that *Aichivirus C* variants generated through recombination or other evolutionary forces are associated with the large-scale outbreak of severe diarrhoea in suckling piglets in China. Further characterization and epidemiological studies are required to determine the exact role of *Aichivirus C* variants in swine disease. We hope data about *Aichivirus C* from China would facilitate its evolutionary researches and pathogenicity study progress in near future.

References:

1. International Committee on Taxonomy of Viruses.

<http://ictvonline.org/proposals/2011.020aV.A.v2.Kobuvirus-Sp.pdf> Accessed in: 17 May 2013

2. Reuter G, Boros Á, Pankovics P: **Kobuviruses—a comprehensive review.** *Reviews in medical virology* 2011, **21**:32-41.

3. Yamashita T, Ito M, Kabashima Y, Tsuzuki H, Fujiura A, Sakae K: **Isolation and characterization of a new species of kobuvirus associated with cattle.** *Journal of general virology* 2003, **84**:3069-3077.
4. Yamashita T, Kobayashi S, Sakae K, Nakata S, Chiba S, Ishihara Y, Isomura S: **Isolation of Cytopathic Small Round Viruses with BS-C1 Cells from Patients with Gastroenteritis.** *Journal of Infectious Diseases* 1991, **164**:954-957.
5. Reuter G, Boldizsár Á, Kiss I, Pankovics P: **Candidate new species of Kobuvirus in porcine hosts.** *Emerging infectious diseases* 2008, **14**:1968.
6. Reuter G, Boros Á, Pankovics P, Egyed L: **Kobuvirus in domestic sheep, Hungary.** *Emerging infectious diseases* 2010, **16**:869.
7. Reuter G, Nemes C, Boros Á, Kapusinszky B, Delwart E, Pankovics P: **Porcine kobuvirus in wild boars (*Sus scrofa*).** *Archives of virology* 2013, **158**:281-282.
8. Kapoor A, Simmonds P, Dubovi EJ, Qaisar N, Henriquez JA, Medina J, Shields S, Lipkin WI: **Characterization of a canine homolog of human Aichivirus.** *Journal of virology* 2011, **85**:11520-11525.
9. Lee M, Jeoung H, Lim J, Song J, Song D, An D: **Kobuvirus in South Korean black goats.** *Virus genes* 2012, **45**:186-189.
10. Carmona-Vicente N, Buesa J, Brown PA, Merga JY, Darby AC, Stavisky J, Sadler L, Gaskell RM, Dawson S, Radford AD: **Phylogeny and prevalence of kobuviruses in dogs and cats in the UK.** *Veterinary microbiology* 2013.
11. Li L, Victoria JG, Wang C, Jones M, Fellers GM, Kunz TH, Delwart E: **Bat guano virome: predominance of dietary viruses from insects and plants plus novel mammalian viruses.** *Journal of virology* 2010, **84**:6955-6965.
12. Phan TG, Kapusinszky B, Wang C, Rose RK, Lipton HL, Delwart EL: **The fecal viral flora of wild rodents.** *PLoS pathogens* 2011, **7**:e1002218.
13. Yu J, Jin M, Zhang Q, Li H, Li D, Xu Z, Li J, Cui S, Yang S, Liu N: **Candidate porcine kobuvirus, China.** *Emerging infectious diseases* 2009, **15**:823.
14. Barry AF, Ribeiro J, Alfieri AF, van der Poel WH, Alfieri AA: **First detection of kobuvirus in farm animals in Brazil and the Netherlands.** *Infection, Genetics and Evolution* 2011, **11**:1811-1814.

15. Ribeiro J, de Arruda Leme R, Alfieri AF, Alfieri AA: **High frequency of Aichivirus C (porcine kobuvirus) infection in piglets from different geographic regions of Brazil.** *Tropical animal health and production* 2013;1-6.
16. Khamrin P, Maneekarn N, Kongkaew A, Kongkaew S, Okitsu S, Ushijima H: **Porcine kobuvirus in piglets, Thailand.** *Emerg Infect Dis* 2009, **15**:2075-2076.
17. Halaihel N, Fuertes H, de Blas I, Gironés O: **Kobuvirus and Hepatitis-E virus infections in piglets: Epidemiology and possible zoonosis.** *Virology* 2011, **3**:134.
18. Khamrin P, Maneekarn N, Hidaka S, Kishikawa S, Ushijima K, Okitsu S, Ushijima H: **Molecular detection of kobuvirus sequences in stool samples collected from healthy pigs in Japan.** *Infection, Genetics and Evolution* 2010, **10**:950-954.
19. An D, Jeoung H, Jeong W, Lee HS, Park J, Kim B: **Porcine kobuvirus from pig stool in Korea.** *Virus genes* 2011, **42**:208-211.
20. Verma H, Mor SK, Abdel-Glil MY, Goyal SM: **Identification and molecular characterization of porcine kobuvirus in US swine.** *Virus genes* 2013:1-3.
21. Dufkova L, Scigalkova I, Moutelikova R, Malenovska H, Prodelalova J: **Genetic diversity of porcine sapoviruses, kobuviruses, and astroviruses in asymptomatic pigs: an emerging new sapovirus GIII genotype.** *Archives of virology* 2012:1-10.
22. Yu J, Xu Z, Li B, Zhang Q, Cui S, Jin M, Duan Z: **Analysis and characterization of the complete genome of a member of a new species of kobuvirus associated with swine.** *Archives of virology* 2011, **156**:747-751.
23. Wang C, Lan D, Cui L, Yang Z, Yuan C, Hua X: **Molecular characterization of a porcine kobuvirus strain in China.** *Archives of virology* 2012, **157**:573-578.
24. Chen RJ, Wu XM, Che YL, Wang LB, Wei H, Zhuang XS, Liu YT, Yan S, Zhou LJ: **Cloning and Evolution Analysis of the 3D Gene of Porcine Kobuvirus in Fujian.** *Fujian Journal of Agricultural Sciences* 2012, **27**:941-944.
25. Shi D, Zhang S, Chen J, Shi H, Zhang X, Feng L: **Molecular characterization of a porcine kobuvirus variant strain in China.** *Archives of virology* 2013:1-5.
26. Zhang S, Shi D, Chen JF, Shi HY, Zhang X, Liu XZ, Liu SX, Wang L, Chen HY, Feng L: **Epidemiological survey of porcine kobuvirus and genetic variation analysis of 3D gene.** *Chinese Journal of Preventive Veterinary Medicine* 2013, **35**:95-98.

27. Fan S, Sun H, Ying Y, Gao X, Wang Z, Yu Y, Li Y, Wang T, Yu Z, Yang S: **Identification and Characterization of Porcine Kobuvirus Variant Isolated from Suckling Piglet in Gansu Province, China.** *Viruses* 2013, **5**:2548-2560.
28. Zhang WB, Leng C, Duan J, Deng SZ: **Establishment and Application of Porcine Kobuvirus RT-PCR detection method.** *China Animal Husbandry & Veterinary Medicine* 2013, **1**:10.
29. Tang XB, Hu RM, Liu WH, Ni DB: **Establishment of Porcine Kobuvirus RT-PCR detection method and epidemiological survey of porcine kobuvirus in Hubei Province.** *Journal of HuaZhong Agriculture University* 2012, **31**: 485-489.
30. Xiang Y, Xue QH, Yin CS, Zhi HB: **Development and Application of a Nested RT-PCR Assay for Detection of Porcine Kobuvirus.** *Chinese Journal of Veterinary Drug* 2013, **47**:5-8.
31. Chen L, Zhu L, Zhou Y, Xu Z, Guo W, Yang W: **Molecular and phylogenetic analysis of the porcine kobuvirus VP1 region using infected pigs from Sichuan Province, China.** *Virology journal* 2013, **10**:281.
32. Wang C, Lan D, Hua X: **Porcine kobuvirus from pig stool specimens in Shanghai, China.** *Virus genes* 2011, **43**:350-352.
33. Reuter G, Kecskeméti S, Pankovics P: **Evolution of porcine kobuvirus infection, Hungary.** *Emerging infectious diseases* 2010, **16**:696.
34. Lin Y, Xiao S, Zeng Y, Song T, Zeng S, Chen H, Fang L: **Complete genome sequence of porcine kobuvirus strain WUH1.** *Journal of virology* 2012, **86**:7010.
35. Cao W, Zheng H, Zhang K, Jin Y, Lv L, Yang F, Liu X: **Complete Genome Sequence of the Porcine Kobuvirus Variant CH/HNXX-4/2012.** *Journal of virology* 2012, **86**:11947.

Figure 1. Distribution of *Aichivirus C* in China

Different shades of red colour indicate various-total infection rates of *Aichivirus C* in porcine faeces and intestine samples. Light red to dark red colours are corresponding to 0% to 100% *Aichivirus C* infection rates. For districts where no literature has been reported, the presence of *Aichivirus C* is indicated by no colour. Districts in orange indicate *Aichivirus C* exists in this region, but currently the prevalence rate is unclear.

Table 1. Prevalence rate of *Aichivirus C* in pigs from different regions of China

District	Infection Rate	Faeces Consistency (Diarrheal or Healthy)
Beijing	11.2% (133/1190)	Healthy
Shanghai	61.2% (30/49)	Diarrheal
Total: 38.8% (45/116)	22.4% (15/67)	Healthy
Hebei	30.12% (97/322)	Healthy
Zhejiang	60% (3/5)	Diarrheal
Anhui	100% (8/8)	Diarrheal
Jiangxi	59.7% (37/62)	Diarrheal
Henan	91.7% (11/12)	Diarrheal
Hubei	76.8% (262/341)	Diarrheal
Total: 66.3% (271/409)	13.2% (9/68)	Healthy
Hunan	88.1% (37/42)	Diarrheal
Sichuan	64.3% (72/112)	Diarrheal
Total: 53.4% (87/163)	29.4% (15/51)	Healthy
Gansu	40.1% (57/142)	Diarrheal
Guangxi	11.8% (2/11)	Diarrheal
Neimenggu	96.2% (2/26)	Unknown
Xinjiang	84.2% (16/19)	Unknown

