



Identification of Myomaker in Yellowfin Seabream (*Acanthopagrus latus*) (Hottuyn, 1782) and its Transcriptional Regulation by Two MyoDs

Kecheng Zhu^{1,2,3}, Peiying He¹, Baosuo Liu^{1,2,3}, Huayang Guo^{1,2,3}, Nan Zhang^{1,2,3}, Liang Guo^{1,2,3}, Shigui Jiang^{1,2,3} and Dianchang Zhang^{1,2,3,*}

¹Key Laboratory of South China Sea Fishery Resources Exploitation and Utilization, Ministry of Agriculture and Rural Affairs; South China Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences, 510300, Guangzhou, Guangdong Province, PR China

²Guangdong Provincial Engineer Technology Research Center of Marine Biological Seed Industry, Guangzhou, Guangdong Province, PR China

³Guangdong Provincial Key Laboratory of Fishery Ecology and Environment, Guangzhou, Guangdong Province, PR China

ABSTRACT

Myomaker is a muscle-specific membrane protein that is essential for myoblast fusion. Myomaker is regulated by myoblast determination protein (MyoD), a muscle-specific basic helix-loop-helix (bHLH) transcription factor in higher vertebrates. However, the transcriptional regulatory mechanism of the *myomaker* gene has not been explored in marine fishes. In the present study, molecular cloning, bioinformatic analysis and transcriptional analysis of *Acanthopagrus latus* myomaker (*Almyomaker*) were performed. The open reading frame (ORF) sequence of *Almyomaker* is 858 bp, which encodes a polypeptide of 285 amino acids. Moreover, phylogenetic and gene structure analysis indicates that *Almyomaker* is highly conserved among vertebrates. The tissue distribution pattern shows that *Almyomaker* is more highly expressed in white muscle than in other tissues. Furthermore, to explore whether two MyoDs are modulators of *Almyomaker*, a promoter analysis was performed using progressive deletion mutations of *Almyomaker*. The results of promoter activity assays show that *Almyomaker* expression is notably activated by two MyoDs. Transcriptional activity of the *Almyomaker* promoter was observed to dramatically decrease after targeted mutation of the MyoD1 M1 and MyoD2 M2 binding sites. In summary, MyoD1 and MyoD2 play an important role in the regulation of *Almyomaker* expression and may promote myoblast fusion during muscle development and growth by modulating *Almyomaker* expression.

Article Information

Received 16 January 2020

Revised 14 February 2020

Accepted 04 March 2020

Available online 14 October 2020

Authors' Contribution

KCZ, SGJ and DCZ designed the research and wrote the paper. PYH and KCZ performed the research. HYG and NZ analyzed the data. BSL and LG contributed reagents/materials/analysis tools.

Key words

Acanthopagrus latus, Myomaker, MyoD, Promoter activity, Mutation analyses.

INTRODUCTION

The fusion of myoblasts is an important process to generate multinucleated myofibers during skeletal muscle regeneration and development (Kim *et al.*, 2015). To date, several proteins, such as myoferlin (Doherty *et al.*, 2005), myogenin and myoD (Tapscott, 2005), Myocyte enhancer factor 2s (mef2s) (Hinitz and Hughes, 2007), Ras-related C3 botulinum toxin substrate 1 (Rac1) (Vasyutina *et al.*, 2009), nephrin (Sohn *et al.*, 2009), Junctional adhesion molecule b and c (Jamb and Jamc) (Powell and Wright, 2011; Shi *et al.*, 2019), CD9 (Charrin *et al.*, 2013), and CD81 (Charrin *et al.*, 2013), involved in the process

of muscle development, have been already identified. However, myomaker, also called transmembrane protein 8c (TMEM8c), is a muscle-specific protein that is absolutely indispensable for myoblast fusion and sufficient to promote fibroblast fusion with muscle cells in zebrafish (*Danio rerio*), mice (*Mus musculus*), and chickens (*Gallus gallus*) (Millay *et al.*, 2013, 2014, 2016; Landemaine *et al.*, 2014; Luo *et al.*, 2015; Zhang *et al.*, 2017). Knockdown of *myomaker* in zebrafish, mice, and chickens indicates that *myomaker* is necessary for myoblast fusion and that loss of myomaker function causes abnormal muscle development (Millay *et al.*, 2013, 2014; Landemaine *et al.*, 2014; Luo *et al.*, 2015; Shi *et al.*, 2018). The amino acid (aa) sequence of myomaker is highly conserved throughout vertebrate species (Millay *et al.*, 2013), and its function in myogenesis is also conserved between mammals and fish (Landemaine *et al.*, 2014). Myomaker is a hydrophobic protein

* Corresponding author: zhangdch@scsfri.ac.cn
0030-9923/2021/0001-0001 \$ 9.00/0

Copyright 2021 Zoological Society of Pakistan

consisting of 221 aa that localizes to intracellular vesicles and the plasma membrane in skeletal myocytes (Millay *et al.*, 2013). The structural features of myomaker span the bilayer seven times with an extracellular N-terminal region and cytosolic C-terminal tail with unknown conserved functional domains (Millay *et al.*, 2016).

Myogenin (MyoG) and MyoD, members of myogenic regulatory factors (MRFs), are pivotal transcription factors (TFs) in myogenesis and can control the transcription of most muscle-related genes (Braun *et al.*, 1989; Edmondson and Olson, 1990; Berkes and Tapscott, 2005; Braun and Gautel, 2011). The two MRFs play a critical role in the regulation of myoblast differentiation. The transcription of myomaker is mediated through the activity of muscle-specific TFs MyoG and MyoD (Luo *et al.*, 2015; Quinn *et al.*, 2017; Zhang *et al.*, 2017; Ganassi *et al.*, 2018). Additionally, during the acute and chronic muscle regeneration process, MyoD can induce *myomaker* expression in muscle satellite cells and accelerate myoblast fusion. Then, the expression of *myomaker* decreased rapidly after fusion (Millay *et al.*, 2014; Landemaine *et al.*, 2014; Demonbreun *et al.*, 2015; Zhang and Roy, 2017). In the MyoD signalling pathway, a downstream gene, *myomixer*, is activated by MyoD. Nonmyoblasts can fuse into multicellular cells when myomaker and myomixer are overexpressed (Bi *et al.*, 2017; Quinn *et al.*, 2017; Zhang *et al.*, 2017). In avians, the essential role of *myomaker* is in myoblast fusion and shows that MyoD can regulate *myomaker* expression (Luo *et al.*, 2015). Two highly conserved e-box mutations in the 5'-regulatory region sequence of the *myomaker* gene in mice and chickens have been verified, suggesting that the importance of the e-box is in regulating the transcription and expression of *myomaker* (Millay *et al.*, 2014; Luo *et al.*, 2015). Moreover, when two MyoD binding sites of the *myomaker* promoter are deleted in chickens, its promoter activity is significantly reduced, indicating that MyoDs have a positive regulatory effect on *myomaker* (Luo *et al.*, 2015). The C-terminal region of myomaker is essential for the function of three cysteine residues, which are speculated to be palmitoylated (Millay *et al.*, 2016). Myomaker displays N-terminal glycine predicted to sustain myristoylation (Bologna *et al.*, 2004). However, the cellular mechanism of *myomaker* and the regulatory mechanism of its expression during myogenesis have not been determined in marine fish.

Yellowfin seabream (*Acanthopagrus latus*) (Hottuyn, 1782), Sparidae, and Perciformes are considered significant aquaculture fish in southern China due to their economic value. Nevertheless, the muscle growth rate is too low in *A. latus*. Consequently, this fish is known as a specific model for exploring regulatory mechanisms in muscle development in marine fish. In the present study,

to investigate the underlying function of *Almyomaker* and transcriptional regulation of two AlMyoDs, this study focused on illuminating the consequence of MyoD in the activation of *Almyomaker* expression. A muscle-specific membrane protein gene from *A. latus*, *myomaker* (*Almyomaker*), was confirmed. Promoter activity assays via the mutation of potential MyoD binding sites are executed to determine key elements in the *Almyomaker* candidate sequence. The present study may contribute to further exploration of *myomaker* function in marine fish and help to elucidate the regulatory mechanism for myoblast fusion.

MATERIALS AND METHODS

Animals and tissues collection

Fish (body weight: 289 ± 18.5 g) were collected from Yangjiang Marine Fish Farm in Guangdong Province, China. For the present study, three healthy fish were used, and 13 tissues (heart, male gonad, eye, skin, brain, fin, spleen, small intestine, gill, white muscle, kidney, liver, and stomach) were isolated to analyze the *myomaker* expression level in different tissues. Then, the flash was frozen in liquid nitrogen and stored at -80°C until use.

Gene cloning and bioinformatics of myomaker

Total RNA (1 μg) was extracted from *A. latus* white muscle by TRIzol Reagent (Takara, Japan). To synthesize cDNA, the Prime ScriptTM RT reagent Kit (Takara) was used according to the manufacturer's instructions. Putative *myomaker*, MyoD1 and MyoD2 sequences were acquired according to genomic data of *A. latus* (Sequence Read Archive under BioProject PRJNA566024). To verify the accuracy of the derived sequence, gene-specific primers were designed (Table 1). The PCR system (volume) and procedure employed were described previously (Zhu *et al.*, 2014). The amplified products, linked into the pEASY-T1 vector (TransGen Biotech, China), were purified by a DNA Purification Kit (Tiangen, China) and then sequenced (Invitrogen, China). Confirmed recombinants were transformed into competent Trans1-T1 cells (TransGen Biotech, China). A BLAST search on the supposed *myomaker* open reading frame (ORF) sequence further confirmed the accuracy and validity of the analysis.

The derived aa sequence from the cloned *Almyomaker* ORF was aligned with other myomaker orthologue proteins from the NCBI and Ensembl databases (Fig. 2). Multiple sequence alignment was performed by ClustalX version 2.0 (Larkin *et al.*, 2007) with default parameters. Phylogenetic analyses for all myomaker aa sequences were achieved using MEGA 6.0 (Tamura *et al.*, 2013) with maximum likelihood (ML) methods (LG + G

model, bootstrap 1000). All available *myomaker* genome sequences were obtained from Ensembl (<http://asia.ensembl.org/>) and Genome Browser (<http://genome.ucsc.edu/cgi-bin/hgBlat>). To predict the signal peptides and transmembrane domain, the Signal P3.0 server (<http://www.cbs.dtu.dk/services/SignalP/>), TMHMM (<http://www.cbs.dtu.dk/services/TMHMM/>), and SMART (<http://smart.embl-heidelberg.de/>) were used. Moreover, the genome structure and phylogenetic tree were embellished using Adobe PhotoShop CS6 (Adobe, San Jose, CA) and FigTree v1.4.2 (<http://tree.bio.ed.ac.uk/software/figtree/>), respectively.

Table I.- Primers used for sequence cloning, deletion mutant construction, and qRT-PCR.

Subject and Primers	Nucleotide sequence
Primers for sequence cloning	
Myomaker-ORF F:	ATGGGTGCGTTTATTGCCAAGAT
Myomaker-ORF R:	TCACTCCAGCCATTCATCT
MyoD1-ORF-F	CGCGGATCCATGGAGCTGTCCGATATCT
MyoD1-ORF-R	CCGCTCGAGCTATAGGACTTGATAGATCA
MyoD2-ORF-F	CGCGGATCCATGGATCTGTCCGACCTTCC
MyoD2-ORF-R	CCGCTCGAGTCAGAGCGGCTCCTGGATGCT
Deletion mutant construction	
Myomaker-pF1	CGGGGTACCCACACAAAATCCAGATCAGC
Myomaker-pF2	CGGGGTACCCGCCAAAAGGGTTAAATC
Myomaker-pF3	CGGGGTACCTGAAGACATTCATACAGA
Myomaker-pF4	CGGGGTACCATGGCTACAATATGGCGTG
Myomaker-pF5	CGGGGTACCAAACCTTAAAGGGCCAAGCTG
Myomaker-pR	CCGCTCGAGCATGGTGGAGGGCCAGTAGACA
Primers for qRT-PCR	
Myomaker-F	CTGGGCGACTTTGATGAGC
Myomaker-R	AGTGATGATGAAGACAGCGGA
EF1 α -F	AAGCCAGGTATGGTTGTCAACTTT
EF1 α -R	CGTGGTGCATCTCCACAGACT

Real-time quantitative PCR analysis

Specific primers for real-time quantitative PCR (qRT-PCR) were designed by Primer Premier 5.0 (Premier Biosoft, USA) based on cloned nucleotide sequences (Table I) (Zhu *et al.*, 2020). The *myomaker*, *MyoDs* and elongation factor 1- α (*EF-1 α*) were tested and used as a target and reference gene, respectively. qRT-PCR was implemented in a quantitative thermal cycle (Mastercycler® eprealplex; Eppendorf, Germany). The PCR system (volume) was described above. The qRT-PCR program was conducted as follows: 95°C for 2 min followed by 40 cycles of 95°C for 10 s, 56°C for 10 s, and 72°C for 20 s. Gene expression was determined using the

2^{- $\Delta\Delta C_t$} method (Livak *et al.*, 2001).

Plasmid construction, cell culture and dual-luciferase reporter assays

Briefly, to clone the target fragment of *Almyomaker*, genomic DNA was isolated from the white muscle of *A. latus*, as previously described in other marine fish (Sun *et al.*, 2013). The ORF sequence upstream of the *myomaker* gene was obtained from genomic data of *A. latus*. Moreover, to obtain the recombinants of *MyoD1* and *MyoD2*, the integrated sequences of *MyoD1* and *MyoD2* ORFs were inserted into the pCDNA3.1-Flag vector (Invitrogen, USA) using specific primers (Table I) (Zhu *et al.*, 2020). To confirm the effect of AlMyoD1 and AlMyoD2 on *Almyomaker* expression, five different truncated regions from *Almyomaker* were amplified by peculiar primers with *KpnI* and *XhoI* restriction sites (Table I). The forward primers (Myomaker-pF1, Myomaker-pF2, Myomaker-pF3, Myomaker-pF4, and Myomaker-pF5) were designed with a 5' *KpnI* site, and the common reverse primer (Myomaker-pR) was designed with a 3' *XhoI* site (Table I). These primers were used to acquire the target region (Myomaker-pF1, 2,090 bp) and four truncated fragments ((i) Myomaker-pF2, 1,483 bp; (ii) Myomaker-pF3, 1,211 bp; (iii) Myomaker-pF4, 436 bp and (iv) Myomaker-pF5, 302 bp) (Fig. 5). PrimeSTAR Master Mix (Takara, Japan) was used to amplify the five truncated mutants. The PCR program consisted of 95°C for 4 min, followed by 30 cycles of 95°C for 40 s, 56°C for 40 s, and 72°C for 1 min. Subsequently, the PCR products were purified by the general DNA Purification Kit (Tiangen, China). A pGL3-basic (Promega, USA) vector and purified PCR products were digested with *KpnI* and *XhoI* and linked by T4 DNA ligase (Takara, Japan) overnight at 16°C. Recombined plasmids were extracted using the EndoFree Plasmid Giga Kit (Tiangen, China), and recombinants were validated by sequencing, as described above.

Furthermore, to investigate the possible function of *MyoD1* or *MyoD2* binding sites on the core *myomaker* promoter and 5'UTR sequences, three truncated mutations of recombinant plasmids were established. The TF binding site prediction (TFBS)-JASPAR database (<http://jaspar.genereg.net/>), TRANSFAC®, and MatInspector® were used to search for potential binding sites in the *myomaker* sequence with *MyoD1* and *MyoD2*, respectively. Then, according to the manufacturer's protocol, site-directed mutagenesis was conducted with a QuickChange II Site-Directed Mutagenesis Kit (Vazyme, USA) from the deletion mutant pGL3-basic-myomaker-p5, which was regarded as the wild-type plasmid. The prediction of three binding sites (M1, M2, and M3) was directly deleted, and the schematic diagram and sequences of homologous

TF binding sites are shown in [Figure 6A](#) and [Table II](#), respectively. The role of TF binding site mutations on the promoter activity of *Almyomaker* was explained by a dual luciferase assay, as described below (Genecreate, China).

Table II.- Sequences of putative binding sites on *AlMyomaker-P5* sequence.

Putative binding sites	Nucleotide sequence	Mutated pattern
M1	GGACAGCTGGGT	Deletion
M2	GGACATCTACTT	Deletion
M3	GACTGAGGAAC	Deletion

Human embryonic kidney (HEK293T) cells were cultured in DMEM (Gibco, USA) with 10% fetal bovine serum (FBS) (Invitrogen, USA) accompanied by 100 μ g/mL streptomycin and 100 U/mL penicillin (Thermo Fisher Scientific, USA) at 37°C in a humidified incubator under 5% CO₂. The procedure of transfection and dual luciferase reporter assays were described by [Li *et al.* \(2017\)](#). Relative luciferase activities (the ratio of firefly and renilla luciferase activities) were measured and calculated using the VICTOR™ X2 Multi-label Plate Reader (PerkinElmer, Inc., Waltham, MA, USA).

Statistical analysis

All trials were executed in triplicate in the present study. All values are shown with three replicates $\pm$ SE. Significant differences were calculated by one way ANOVA tests. *P* values <0.05 were considered to be significant.

RESULTS

Sequence characterization of Almyomaker

The genomic sequence of *Almyomaker* is 3,383 bp, including 6 exons and 5 introns (Supplementary Fig. S1). The full-length ORF of *Almyomaker* is 855 bp (Fig. 1A), encoding peptides of 285 amino acids with a predicted molecular weight of 32.14 kDa and a theoretical isoelectric point of 9.44 (Accession No. MN266854). A conserved domain (DUF3522 superfamily starting from the 3rd amino acid and ending at the 185th amino acid) was detected using BLAST (Huang *et al.*, 2019). Hydrophobicity profiles show that the *Almyomaker* protein includes more hydrophobic amino acids than hydrophilic amino acids (Fig. 1B). The results of TMHMM indicate that there are 6 obvious transmembrane domains in the *Almyomaker* protein (Fig. 1C), which is consistent with the analysis of goose *Anser cygnoides* (He *et al.*, 2017).




Fig. 1. Sequence characterization of the *Myomaker* gene in *Acanthopagrus latus*. A, the nucleotide sequence of the *Myomaker* gene and the deduced amino acid sequence of *T. ovatus*. Initiation and termination codons are marked by green. Six transmembrane domains are shown in red; B, hydrophobicity profile of *Myomaker*; C, TMHMM posterior probabilities of *Myomaker*.




Fig. 2. Amino acid sequences of Myomaker homologues in vertebrates. A, amino acid alignment of Myomaker proteins from *A. latus* (Al), *Oreochromis niloticus* (On), *Xiphophorus maculatus* (Xm), *Oryzias latipes* (Ol), *Poecilia formosa* (Pf), *Danio rerio* (Dr), *Tetraodon nigroviridis* (Tn), *Gallus gallus* (Gg), *Latimeria chalumnae* (Lc), *Pelodiscus sinensis* (Ps), *Homo sapiens* (Hs), and *Mus musculus* (Mm). Conserved sequences are marked with asterisks. B, Percent identities of Myomaker amino acids compared to the above 12 Myomaker amino acids, and the accession numbers of the Myomaker sequences used are also listed.

The myomaker amino acids in *A. latus* reveal sequence homology with those in the other metazoan species (Fig. 2A) and are most highly homologous with *Xiphophorus maculatus*, *Oryzias latipes*, *Poecilia formosa*, and *Tetraodon nigroviridis*. Moreover, a BLAST analysis indicated that the Almyomaker protein sequence shares high sequence identity with myomaker sequences from other teleosts containing tilapia (*O. niloticus*, 92 %), platyfish (*X. maculatus*, 87.1%), medaka (*O. latipes*, 86.7%), and molly (*P. formosa*, 86.3%) and low sequence

identity with humans (*Homo sapiens*, 56.5%) and mice (*Mus musculus*, 54.7%) (Fig. 2B).

Almyomaker structural and phylogenetic analyses

The genomic structural features and phylogenetic relationship of myomaker were determined and constructed in metazoans (Fig. 3). The lengths and distributions of the exons and introns of metazoan myomaker genes are displayed in Supplementary Table I. Six exons and five introns were identified in myomaker gene sequences,

except for zebrafish (*Danio rerio*), Tetraodon (*Tetraodon nigroviridis*), Chinese softshell turtle (*Pelodiscus sinensis*), Coelacanth (*Latimeria chalumnae*), chicken (*Gallus gallus*), *H. sapiens* and *M. musculus* myomaker, which only possess five exons and four introns. Furthermore, the sizes of the exon sequences reveal that there is nearly no diversity among species. Additionally, the phylogenetic tree was constructed with the full-length myomaker aa of various fish, mammalia, aves, and amphibia species. *Almyomaker* is grouped together with other Perciformes, such as *O. niloticus*. The phylogenetic tree clearly shows that fish evolve into one branch followed by Amphibians,

Aves and then Mammals (Huang *et al.*, 2019).

Tissue expression of *Almyomaker*

The mRNA levels of *Almyomaker* are determined by qRT-PCR in various kinds of tissues. qRT-PCR shows that *Almyomaker* is broadly distributed in various tissues, and the abundance of mRNA varies among tissues. The highest *Almyomaker* transcriptions were detected in the white muscle followed by the brain and spleen, whereas the lowest *Almyomaker* expression levels were found in the liver, skin and stomach (Fig. 4).




Fig. 3. Phylogenetic relationship and structure of the *Almyomaker* gene with other vertebrates. Genome structure analysis of *Almyomaker* genes according to their phylogenetic relationship. Lengths of exons and introns of each *Myomaker* gene are displayed proportionally. Different colour boxes and lines represent exons and introns, respectively. The identical colour boxes represent homologous sequences.




Fig. 4. Relative expression levels of *Almyomaker* in different tissues. The twelve tissues are heart, gonad, eye, skin, brain, fin, spleen, small intestine, gill, white muscle, kidney, liver, and stomach. Different letters indicate significant differences ($p < 0.05$).




Fig. 5. Promoter activity analysis of the *Almyomaker* gene. A, the structure of the *Almyomaker* promoter and 5'UTR region. Five recombinant plasmids were denoted myomaker-p1 (-1822 to +267), myomaker-p2 (-1215 to +267), myomaker-p3 (-943 to +267), myomaker-p4 (-168 to +267) and myomaker-p5 (-34 to +267). B and C, Transcriptional activity of the *Almyomaker* promoter. These plasmids were transfected along with the transcription factors MyoD1 (B) and MyoD2 (C) into HEK 293T cells. Dual-luciferase activity was driven by the *Almyomaker* promoter upon the transfection of pcDNA3.1-MyoD1, pcDNA3.1-MyoD2 or pcDNA3.1 into HEK 293T cells. Data are presented as the means of three replicates $\pm$ SE. Different letters indicate significant differences ($p < 0.05$).




Fig. 6. Construction of truncated mutants for the identification of predicted transcription factor (TF) binding sites in the *Almyomaker* candidate sequence. A, the nucleotide sequence and predicted binding sites in the core region of the *Almyomaker*-p5. TSS indicates the transcription start site. Effects of three mutants on *Almyomaker*-p5 promoter activity transfected with pcDNA3.1-MyoD1 (B) or pcDNA3.1-MyoD2 (C) or pcDNA3.1. Binding sites are shown with boxes. Data are presented as the means of three replicates $\pm$ SE. Different letters indicate significant differences ($p < 0.05$).

Two *AlMyoDs* activate *Almyomaker* expression

The amplified candidate *Almyomaker* promoter and

5'-UTR region (2,090 bp) are upstream nontranslational sequences. To comprehend the binding region of two *MyoDs*

in the *Almyomaker* sequence, a full-length target fragment and 4 truncated mutants were inserted with a promoter-less luciferase reporter vector, pGL3-basic (Promega, USA). The result of promoter activity analysis shows that the construct of myomaker-p5 (-34 bp to +267 bp) is higher than that of other constructs with cotransfection of MyoD1 or MyoD2, suggesting that the core binding region is located at -34 bp to +267 bp, which may include several MyoD1 or MyoD2 binding sites (Fig. 5). Therefore, the sequence of construct myomaker-p5 (-34 bp to +267 bp) was used for further functional analysis.

To understand the MyoD1 and MyoD2 binding sites in *Almyomaker*, the presumptive binding sites are mutated (Fig. 6A; Table II). The effects on promoter activity were investigated in 293T cells, together transfected with each mutant and MyoD1 or MyoD2. The results reveal that mutation of the M1 binding site (GGACAGCTGGGT, -14 bp to -3 bp) generates a marked reduction in promoter activity (Fig. 6B), showing that M1 is the MyoD1 binding site on the *Almyomaker* promoter. Moreover, mutation of the M2 binding site (GGACATCTACTT, +92 bp to +103 bp) resulted in a significant reduction in promoter activity (Fig. 6C), suggesting that M2 is the potential MyoD2 binding site in the *Almyomaker* gene 5'-UTR region. Notably, one predicted binding site (M3) does not activate luciferase activity with MyoD1 or MyoD2, suggesting that this site is not required for triggering *Almyomaker* expression with two MyoDs.

DISCUSSION

The present study investigated the potential mechanisms of the transcriptional regulation of *myomaker* by MyoD1 and MyoD2 in *A. latipes*. The sequence and functional characterization, tissue expression and regulation relationship between two AlMyoDs and *Almyomaker* were determined. The *Almyomaker* aa sequence was 54.7%–92.0% identical to myomaker proteins from other teleosts.

In general, the expression of several genes depends on RNA polymerases and TFs binding to specific sequences on the promoters of target genes in eukaryotic organisms (O'Malley, 1977; Xie *et al.*, 2018; Zhu *et al.*, 2019). Furthermore, the integrity and activity of a promoter could regulate gene transcription. The TF MyoDs monitored skeletal muscle growth in mammals and altered the transcription of muscle-related genes in teleosts and avians (Millay *et al.*, 2014; Luo *et al.*, 2015). Notably, evidence has demonstrated that overexpression of MyoD could increase downstream gene *myomaker* expression in zebrafish (Bi *et al.*, 2017; Quinn *et al.*, 2017; Zhang *et al.*, 2017). It was unclear whether MyoD could upregulate

Almyomaker in marine fishes. In the present study, dual-luciferase reporter assays were conducted to realize regulatory mechanisms whereby MyoD1 and MyoD2 were believed to accommodate *Almyomaker* expression. The results of truncated mutants indicated that *Almyomaker* reporter activity was activated by the overexpression of two AlMyoDs. The core binding region in the *Almyomaker* promoter was -34 to +267 bp (Fig. 6). This evidence was the first to show that the expression of *Almyomaker* could be upregulated by both AlMyoD1 and AlMyoD2 in marine fishes.

MyoDs possess a highly conserved bHLH domain, which is necessary for heterodimerization with characteristic DNA binding to the e-box motif (consensus CANNTG) detected in the regulatory regions of their target genes (Murre *et al.*, 1989; Lassar *et al.*, 1989, 1991; Davis and Weintraub, 1992). Consequently, to investigate the two MyoD binding sites on the *Almyomaker* promoter, three mutant vectors were constructed. Mutation of the MyoD1 M1 binding site (GGACAGCTGGGT) and MyoD2 M2 binding site (GGACATCTACTT) resulted in observably decreased promoter activity (Fig. 6B, C), suggesting that the MyoD1 binding M1 site and MyoD2 binding M2 site were essential for *Almyomaker* promoter activity, respectively. The sequences of the above two binding sites were representative e-box motifs (Davis and Weintraub, 1992). In brief, MyoD1 and MyoD2 could mediate *myomaker* promoter expression by combining e-box motifs (M1 and M2 binding sites) in fish, respectively, which was consistent with the findings in *M. musculus* and *G. gallus* (Millay *et al.*, 2014; Luo *et al.*, 2015). Moreover, the MyoD2 M2 binding site was located in the 5'UTR, which was analogous to previous studies (Wu *et al.*, 2018; Khan *et al.*, 2019).

Phylogenetic analysis of myomakers in tetrapods and teleosts showed that a protein of uniform length (221 aa) was observed in ancestral nonteleost fish, and teleost myomaker sequences were classified into three groups according to myomaker protein length (Landemaine *et al.*, 2019). In the present study, phylogenetic analysis showed that a typical phylogeny revealing the amino acid sequence of *Almyomaker* was closely matched to myomakers of *O. niloticus*, *X. maculatus*, *O. latipes*, *P. formosa*, and *T. nigroviridis*, with analogous lengths. This result was similar to that reported in Landemaine *et al.* (2019). A genome structure analysis showed that all myomakers contained 6 exons and 5 introns in metazoans, except for *D. rerio*, *T. nigroviridis*, *P. sinensis*, *L. chalumnae*, *G. gallus*, *H. sapiens* and *M. musculus* myomaker, which only possessed 5 exons and 4 introns, suggesting that the last exon might be lost during evolution.

Previous studies have focused on the role of myomaker

during embryonic development and have determined that myomaker was indispensable for the facilitation of myoblast fusion in such species as *D. rerio*, *G. gallus*, and *M. musculus* (Millay *et al.*, 2013; Landemaine *et al.*, 2014; Luo *et al.*, 2015; He *et al.*, 2017; Zhang and Roy, 2017). The mRNA expression patterns of the *myomaker* gene during 8 different postnatal developmental stages in the Japanese flounder (*Paralichthys olivaceus*) showed that the expression of *myomaker* at 180 dph was higher than that at other periods (Huang *et al.*, 2019). Moreover, myomaker is expressed only in skeletal muscle in *M. musculus*, *G. gallus* and *Oncorhynchus mykiss* (Millay *et al.*, 2013; Landemaine *et al.*, 2019). In the present study, the tissue-specific expression pattern revealed that the highest *Almyomaker* mRNA expression was detected in white muscle, showing that *Almyomaker* played an important role in muscle development. *Almyomaker* is also broadly expressed in other tissues; however, in mammals, chicks and fish, *Myomaker* is nearly undetectable in other organs (Millay *et al.*, 2013; Landemaine *et al.*, 2019). This discrepancy may be attributable to species diversity.

CONCLUSION

In conclusion, the full-length *Almyomaker* genome sequence was cloned. The tissue expression profile indicated that the mRNA level of *Almyomaker* was highest in the heart and gonad among the detected tissues. The luciferase activity analysis showed that the region from -34 bp to +267 bp includes the core binding region. Mutation analyses indicated that the activity of the *Almyomaker* promoter significantly decreased after the targeted mutation of the M1 and M2 binding sites with MyoD1 and MyoD2, respectively. This study mainly focused on the transcription of *Almyomaker* by MyoDs in heterologous cells. In the future, more attention should be paid to the mRNA or protein levels of *Almyomaker* after overexpression of MyoDs in endogenous cells.

ACKNOWLEDGMENTS

The study was supported by the Natural Science Foundation of Guangdong Province (2017A030310594), the National Infrastructure of Fishery Germplasm Resources Project (2019DKA30470) and the Science and Technology Infrastructure Construction Project of Guangdong Province (2019B030316030).

Supplementary material

There is supplementary material associated with this article. Access the material online at: <https://dx.doi.org/10.17582/journal.pjz/20200116020138>

Statement of conflict of interests

The authors declare no competing interests.

REFERENCES

- Berkes, C.A. and Tapscott, S.J., 2005. MyoD and the transcriptional control of myogenesis. *Semin. Cell Dev. Biol.*, **16**: 585-595. <https://doi.org/10.1016/j.semcdb.2005.07.006>
- Bi, P.P., Ramirez-Martinez, A., Li, H., Cannavino, J., McAnally, J.R., Shelton, J.M., Sanchez-Ortiz, E., Bassel-Duby, R. and Olson, E.N., 2017. Control of muscle formation by the fusogenic micropeptide myomixer. *Science*, **356**: 323-327. <https://doi.org/10.1126/science.aam9361>
- Bologna, G., Yvon, C., Duvaud, S. and Veuthey, A.L., 2004. N-Terminal myristoylation predictions by ensembles of neural networks. *Proteomics*, **4**: 1626-1632. <https://doi.org/10.1002/pmic.200300783>
- Braun, T., Buschhausen-Denker, G., Bober, E., Tannich, E. and Arnold, H.H., 1989. A novel human muscle factor related to but distinct from MyoD1 induces myogenic conversion in 10T1/2 fibroblasts. *EMBO J.*, **8**: 701-709. <https://doi.org/10.1002/j.1460-2075.1989.tb03429.x>
- Braun, T. and Gautel, M., 2011. Transcriptional mechanisms regulating skeletal muscle differentiation, growth and homeostasis. *Nat. Rev. Mol. Cell Biol.*, **12**: 349-361. <https://doi.org/10.1038/nrm3118>
- Charrin S., Latil M., Soave S., Polesskaya A., Christien F., Boucheix C. and Rubinstein E., 2013. Normal muscle regeneration requires tight control of muscle cell fusion by tetraspanins CD9 and CD81. *Nat. Commun.*, **4**: 1674. <https://doi.org/10.1038/ncomms2675>
- Davis, R.L. and Weintraub, H., 1992. Acquisition of myogenic specificity by replacement of three amino acid residues from MyoD into E12. *Science*, **256**: 1027-1030. <https://doi.org/10.1126/science.1317057>
- Demonbreun, A.R., Biersmith, B.H. and McNally, E.M., 2015. Membrane fusion in muscle development and repair. *Semin. Cell Dev. Biol.*, **45**: 48-56. <https://doi.org/10.1016/j.semcdb.2015.10.026>
- Doherty, K.R., Cave, A., Davis, D.B., Delmonte, A.J., Posey, A. and Earley, J.U., 2005. Normal myoblast fusion requires myoferlin. *Development*, **132**: 5565-5575. <https://doi.org/10.1242/dev.02155>
- Edmondson, D.G. and Olson, E.N., 1990. A gene with homology to the myc similarity region of MyoD1 is expressed during myogenesis and is sufficient to

- activate the muscle differentiation program. *Genes Dev.*, **4**: 1450. <https://doi.org/10.1101/gad.4.8.1450>
- Ganassi, M., Badodi, S., Quiroga, H.P.O., Zammit, P.S., Hinitis, Y. and Hughes, S.M., 2018. Myogenin promotes myocyte fusion to balance fibre number and size. *Nat. Commun.*, **9**: 4232. <https://doi.org/10.1038/s41467-018-06583-6>
- He, K., Ren, T., Zhu, S., Liang, S. and Zhao, A., 2017. Transiently expressed pattern during myogenesis and candidate miRNAs of Tmem8C in goose. *J. Genet.*, **96**: 39-46. <https://doi.org/10.1007/s12041-016-0737-8>
- Hinitis, Y. and Hughes, S.M., 2007. Mef2s are required for thick filament formation in nascent muscle fibres. *Development*, **134**: 2511-2519. <https://doi.org/10.1242/dev.007088>
- Huang, Y., Wu, S., Zhang, J., Wen, H., Zhang, M. and He, F., 2019. Methylation status and expression patterns of myomaker gene play important roles in postnatal development in the Japanese flounder (*Paralichthys olivaceus*). *Gen. Comp. Endocrinol.*, **280**: 104-114. <https://doi.org/10.1016/j.ygcen.2019.04.017>
- Khan, R., Raza, S.H.A., Junjvlieke, Z., Wang, X.Y., Garcia, M., Elnour, I.E., Wang H.B. and Zan, L.S., 2019. Function and transcriptional regulation of bovine TORC2 gene in adipocytes: Roles of C/EBP γ , XBP1, INSM1 and ZNF263. *Int. J. Mol. Sci.*, **20**: 4338. <https://doi.org/10.3390/ijms20184338>
- Kim, J.H., Jin, P., Duan, R. and Chen, E.H., 2015. Mechanisms of myoblast fusion during muscle development. *Curr. Opin. Genet. Dev.*, **32**: 162-170. <https://doi.org/10.1016/j.gde.2015.03.006>
- Landemaine, A., Rescan, P.Y. and Gabillard, J.C., 2014. Myomaker mediates fusion of fast myocytes in zebrafish embryos. *Biochem. biophys. Res. Commun.*, **451**: 480-484. <https://doi.org/10.1016/j.bbrc.2014.07.093>
- Landemaine, A., Ramirez-Martinez, A., Monestier, O., Sabin, N., Rescan, P.Y., Olson, E.N. and Gabillard, J.C., 2019. Trout myomaker contains 14 minisatellites and two sequence extensions but retains fusogenic function. *J. biol. Chem.*, **294**: 6364-6374. <https://doi.org/10.1074/jbc.RA118.006047>
- Larkin, M.A., Blackshields, G., Brown, N.P. and Higgins, D.G., 2007. ClustalW and ClustalX version 2. *Bioinformatics*, **23**: 2947-2948. <https://doi.org/10.1093/bioinformatics/btm404>
- Lassar, A.B., Buskin, J.N., Lockshon, D., Davis, R.L., Apone, S., Hauschka, S.D. and Weintraub, H., 1989. MyoD is a sequence specific DNA binding protein requiring a region of myc homology to bind to the muscle creatine kinase enhancer. *Cell*, **58**: 823-831. [https://doi.org/10.1016/0092-8674\(89\)90935-5](https://doi.org/10.1016/0092-8674(89)90935-5)
- Lassar, A.B., Davis, R.L., Wright, W.E., Kadesch, T., Murre, C., Vorona, A., Baltimore, D. and Weintraub, H., 1991. Functional activity of myogenic HLH proteins requires hetero-oligomerization with E12/E47-like proteins *in vivo*. *Cell*, **66**: 305-315. [https://doi.org/10.1016/0092-8674\(91\)90620-E](https://doi.org/10.1016/0092-8674(91)90620-E)
- Li, S.L., Monroig, O., Wang, T.J., Yuan, Y.H., Navarro, J.C., Hontoria, F., Liao, K., Tocher, D.R., Mai, K.S. and Xu, W., 2017. Functional characterization and differential nutritional regulation of putative Elovl5 and Elovl4 elongases in large yellow croaker (*Larimichthys crocea*). *Scient. Rep.*, **7**: 2303. <https://doi.org/10.1038/s41598-017-02646-8>
- Livak, K.J. and Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{- $\Delta\Delta C_t$} method. *Methods*, **25**: 402-408. <https://doi.org/10.1006/meth.2001.1262>
- Luo, W., Li, E., Nie, Q.H. and Zhang, X.Q., 2015. Myomaker, Regulated by MyoD, MyoG and miR-140-3p, Promotes chicken myoblast fusion. *Int. J. mol. Sci.*, **16**: 26186-26201. <https://doi.org/10.3390/ijms161125946>
- Millay, D.P., Gamage, D.G., Quinn, M.E., Min, Y.L., Mitani, Y., Bassel-Duby, R. and Olson, E.N., 2016. Structure-function analysis of myomaker domains required for myoblast fusion. *Proc. natl. Acad. Sci. U.S.A.*, **113**: 2116-2121. <https://doi.org/10.1073/pnas.1600101113>
- Millay, D.P., O'Rourke, J.R., Sutherland, L.B., Bezprozvannaya, S., Shelton, J.M., Bassel-Duby, R. and Olson, E.N., 2013. Myomaker is a membrane activator of myoblast fusion and muscle formation. *Nature*, **499**: 301-305. <https://doi.org/10.1038/nature12343>
- Millay, D.P., Sutherland, L.B., Bassel-Duby, R. and Olson, E.N., 2014. Myomaker is essential for muscle regeneration. *Genes Dev.*, **28**: 1641-1646. <https://doi.org/10.1101/gad.247205.114>
- Murre, C., McCaw, P.S., Vaessin, H., Caudy, H., Jan, L.Y., Jan, Y.N., Cabrera, C.V., Buskin, J.N., Hauschka, S.D., Lassar, A.B., Weintraub, H. and Baltimore, D., 1989. Interactions between heterologous helix-loop-helix proteins generate complexes that bind specifically to a common DNA sequence. *Cell*, **58**: 537-544. [https://doi.org/10.1016/0092-8674\(89\)90434-0](https://doi.org/10.1016/0092-8674(89)90434-0)
- O'Malley, B.W., Towle, H.C. and Schwartz, R.J., 1977. Regulation of gene expression in eucaryotes. *Annu. Rev. Genet.*, **11**: 239-275. <https://doi.org/10.1146/annurev.ge.11.120177.001323>

- Powell, G.T. and Wright, G.J., 2011. Jamb and Jamc are essential for vertebrate myocyte fusion. *PLoS Biol.*, **9**: e1001216. <https://doi.org/10.1371/journal.pbio.1001216>
- Quinn, M.E., Goh, Q.N., Kurosaka, M., Gamage, D.G., Petrany, M.J., Prasad, V. and Millay, D.P., 2017. Myomerger induce fusion of non-fusogenic cells and is required for skeletal muscle development. *Nat. Commun.*, **8**: 15665. <https://doi.org/10.1038/ncomms15665>
- Shi, J., Cai, M.X., Si, Y.F., Zhang, J.S. and Du, S.J., 2018. Knockout of myomaker results in defective myoblast fusion, reduced muscle growth and increased adipocyte infiltration in zebrafish skeletal muscle. *Hum. Mol. Genet.*, **27**: 3542-3554. <https://doi.org/10.1093/hmg/ddy268>
- Si, Y.F., Wen, H.S. and Du, S.J., 2019. Genetic mutations in jamb, jamc, and myomaker revealed different roles on myoblast fusion and muscle growth. *Mar. Biotechnol.*, **21**: 111-123. <https://doi.org/10.1007/s10126-018-9865-x>
- Sohn, R.L., Huang, P., Kawahara, G., Mitchell, M., Guyon, J., Kalluri, R., Kunkel, L.M. and Gussoni, E., 2009. A role for nephrin, a renal protein, in vertebrate skeletal muscle cell fusion. *Proc. natl. Acad. Sci.*, **106**: 9274-9279. <https://doi.org/10.1073/pnas.0904398106>
- Sun, L.Y., Zhang, D.C., Jiang, S.G., Guo, H.Y. and Zhu, C.Y., 2013. Isolation and characterization of 21 polymorphic microsatellites in golden pompano *Trachinotus ovatus*. *Conserv. Genet. Resour.*, **5**: 1107-1109. <https://doi.org/10.1007/s12686-013-9942-4>
- Tamura, K., Stecher, G., Peterson, D., Filipinski, A. and Kumar, S., 2013. MEGA6: Molecular evolutionary genetics analysis, Version 6.0. *Mol. Biol.*, **3**: 2725-2729. <https://doi.org/10.1093/molbev/mst197>
- Vasyutina, E., Martarelli, B., Brakebusch, C., Wende, H. and Birchmeier, C., 2009. The small G-proteins Rac1 and Cdc42 are essential for myoblast fusion in the mouse. *Proc. natl. Acad. Sci.*, **106**: 8935-8940. <https://doi.org/10.1073/pnas.0902501106>
- Wu, X., Xu, F.L., Ding M., Zhang, J.J., Yao, J. and Wang, B.J., 2018. Characterization and functional analyses of the human HTR1A gene: 5' regulatory region modulates gene expression in vitro. *BMC Genet.*, **19**: 115. <https://doi.org/10.1186/s12863-018-0708-6>
- Xie, D., Fu, Z., Wang, S., You, C., Monroig, Ó., Tocher, D.R. and Li, Y., 2018. Characteristics of the fads2 gene promoter in marine teleost *Epinephelus coioides* and role of Sp1-binding site in determining promoter activity. *Scient. Rep.*, **8**: 5305. <https://doi.org/10.1038/s41598-018-23668-w>
- Tapscott, S.J., 2005. The circuitry of a master switch: MyoD and the regulation of skeletal muscle gene transcription. *Development*, **132**: 2685-2695. <https://doi.org/10.1242/dev.01874>
- Zhang, Q., Vashisht, A.A., O'Rourke, J., Corbel, S.Y., Moran, R., Romero, A., Miraglia, L., Zhang, J., Durrant, E. and Schmedt, C., 2017. The microprotein Minion controls cell fusion and muscle formation. *Nat. Commun.*, **8**: 15664. <https://doi.org/10.1038/ncomms15664>
- Zhang, W.B. and Roy, S., 2017. Myomaker is required for the fusion of fast-twitch myocytes in the zebrafish embryo. *Dev. Biol.*, **423**: 24-33. <https://doi.org/10.1016/j.ydbio.2017.01.019>
- Zhu, K.C., Chen, L.P., Zhao, J.K., Wang, H.J., Wang, W.M., Li, Z. and Wang, H.L., 2014. Molecular characterization and expression patterns of myogenin in compensatory growth of *Megalobrama amblycephala*. *Comp. Biochem. Physiol. B*, **170**: 10-17. <https://doi.org/10.1016/j.cbpb.2014.01.001>
- Zhu, K.C., Song, L., Guo, H.Y., Guo, L., Zhang, N., Liu, B.S., Jiang, S.G. and Zhang, D.C., 2019. Elovl4a participates in LC-PUFA biosynthesis and is regulated by PPARα in golden pompano *Trachinotus ovatus* (Linnaeus 1758). *Scient. Rep.*, **9**: 4684. <https://doi.org/10.1038/s41598-019-41288-w>
- Zhu, K.C., Liu, B.S., Guo, H.Y., Zhang, N., Guo, L., Jiang, S.G. and Zhang, D.C., 2020. Functional analysis of two MyoDs revealed their role in the activation of myomixer expression in yellowfin seabream (*Acanthopagrus latus*) (Hottuyn, 1782). *Int. J. Biol. Macromol.*, <https://doi.org/10.1016/j.ijbiomac.2019.11.139>



Supplementary Material

Identification of Myomaker in Yellowfin Seabream (*Acanthopagrus latus*) (Hottuyn, 1782) and its Transcriptional Regulation by Two MyoDs

Kecheng Zhu^{1,2,3}, Peiying He¹, Baosuo Liu^{1,2,3}, Huayang Guo^{1,2,3}, Nan Zhang^{1,2,3}, Liang Guo^{1,2,3}, Shigui Jiang^{1,2,3} and Dianchang Zhang^{1,2,3,*}

¹Key Laboratory of South China Sea Fishery Resources Exploitation and Utilization, Ministry of Agriculture and Rural Affairs; South China Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences, 510300, Guangzhou, Guangdong Province, PR China

²Guangdong Provincial Engineer Technology Research Center of Marine Biological Seed Industry, Guangzhou, Guangdong Province, PR China

³Guangdong Provincial Key Laboratory of Fishery Ecology and Environment, Guangzhou, Guangdong Province, PR China

Supplementary Table I.- Lengths of exons and introns of each *Myomaker* gene.

Species	E1	I1	E2	I2	E3	I3	E4	I4	E5	I5	E6
<i>Acanthopagrus latus</i>	135	704	115	967	149	338	117	249	141	260	201
<i>Oreochromis niloticus</i>	135	119	115	1187	149	311	117	160	141	1467	210
<i>Oryzias latipes</i>	135	248	115	731	149	142	117	104	141	1715	201
<i>Poecilia formosa</i>	135	936	115	1144	149	502	117	114	141	1209	204
<i>Xiphophorus maculatus</i>	135	919	115	848	149	1008	117	114	141	4917	201
<i>Danio rerio</i>	135	1764	115	694	149	1933	117	99	147	-	-
<i>Tetraodon nigroviridis</i>	135	75	115	658	149	79	117	159	144	-	-
<i>Pelodiscus sinensis</i>	135	4396	115	6029	149	2988	117	4519	144	-	-
<i>Latimeria chalumnae</i>	135	2095	115	3509	149	2558	117	2984	144	-	-
<i>Gallus gallus</i>	135	2203	115	1893	149	589	117	904	144	-	-
<i>Homo sapiens</i>	135	4421	115	1151	149	3266	117	705	147	-	-
<i>Mus musculus</i>	135	4413	115	913	149	3472	117	364	147	-	-

* Corresponding author: zhangdch@scsfri.ac.cn

0030-9923/2021/0001-0001 \$ 9.00/0

Copyright 2021 Zoological Society of Pakistan

ATGGGTGCGTTTATTGCCAAGATGCTGCTTCCTACAGTCAGCAGTCTGGTGTTCCTGCCTGCAGCCAGC
 GTGGCCACCAAGAGGGGCTTCCACATGGAGGCCATGGTCTACTTCTTCACGATGTTCTTCACCGCGT
 GAGAATGCGGCGATGGCTCTGTTTCGATATCACGGATTGAAAATGTGTTGCTTGGTTGTGATGTTTGC
 TGTCTAAGTCACTGAACATGCGGGAAGCTGAGAACATTGATGTTTCGAGTTTGGATCAGAAGATAAC
 CCTAACGCCATATCATTTTAAAACTGCTTTTCATTTTATGTTTTTGCATCTGCTGTTCAACTTACCGGCAT
 TAAATTTGATCAGCGTAAGTAAGAAAAGTTGTGAAGAATTTACACAATCAGAGGTAAGGATGGGACTCG
 ATAGCATTTTATGTAATCACCATAAGATCTGTTTATCAACTTCAAATATCCTTTACAGGCTTTGCTTACTT
 ATTTTACATACAAATATAAATATATAAAAGATTATATAAATATATCATAAACCTTACTAACTCAAACCTTTT
 ATCTTCCCTCTCTTTAATTGACAGCATCTATGGTTTGTGCATCCCTGTTTGTCTATAAGAAGTGTGTCTCA
 AAGCTTTTATAATTTGTGCGTTTAAATCCAAACAGACAACCCAATTCACACATTGTTTTCATTATATC
 AAAAATGACCTACTGGTGATTTCACACCTTCTATAAGCGCAACCTGTTGTGACATTGAAACTGGGACT
 GTATGTGAAGGTCATTAATAACCTTATAGATACTCCTGCTGTCTGTTTGACTTGAACCTTGTATGTCTTTCT
 GTAGATTACCACGCGATGTGATGGACCAGGTCTCTCCATCCTGTGTTTCATGAAATATGACATTCTCGAG
 TACTTCAGTGTTTACAGCACAGCTCTCTCCATGTGGGTACACTCATAGGTAAGTACAGTCTGTGCGAC
 TCTATACACAGCGAAAGCTTGTGATTTCGTATCATGAACACACCGGAAAAACAGGGAACAGTAGAGG
 CCACAAACAACCTTTTCTTTTGAATATCTTTAGGTAGTGCTGCTTGACAGTGACGGACAGAAAGATAT
 TTACATGACCAACAGCTTAAATGTAAACACACTTTAAAGCATCACTCCTGAAAAAGGGCGTAAATTAG
 CATGTTAATCCAGGCCAAGCTGAGTTTTTTTTTTTTTTTTTGGCATTATTACACAAAATATCTTTTATCCA
 AAGGTTTTGTAAACCTTCTGAACAACCTGGAAGCTAGTGCTAGGGTATTTTTGCGTAAACTCAAGCTA
 GCGAGCACTGGCTAAAAAAGAATATGATTGATATGATTTTAAAGCTCATATCCGACAATTACACTTCCCA
 AGCACAAGTTCAAGAAATAAAAAATCCAGCTCTGGCTGCAACAGTTGTTGACAGCATATGTTGTGTCTT
 TGAGTTCATCTACATGTGCTCCGAAATGCCTGCTCTAATATGATCACACACAGACACACACACACAAA
 TACATTTGTTCTCTCCATCGACCAATTAAGCCCTGAGGTTTTGCTACAAGGCTTTTAAATCACATG
 CTGTGAGGGGCCAAATGCTTCGACAGCTGACTTTACTGATTGGTCTGCTCTCTTTGTCGGGTGGTGTA
 GCCTTCAGGATATTTGGCAATGGCGATGTGTGTCTTTCTCATGGAAGACAGCAGAATGCAAACTCTGA
 AAAGATTCCTGGTGTTATTTTTAGCCGAACCTGAAAACCTAGAGATTCTCTGGCATTACGGTATTTTTATA
 GGATTTGTGAGTGCATCTGGATTCTGGATAAGAAATGTATCTAATGTAATTTTGGTCTACATTAAGTAA
 AAATGTGCCTGCCTGTTAGAGTAACATACATGTATGTGTGTCCCTGTTCCAGCTCTGGGCGACTTTGAT
 GAGCCCCAGCGCTCCAGTATAACCATGTTTGGAGTCTGACGACTGCGGTGAGGATCTACCAGGATCG
 CTGGGGCTATGGGATCTACTCCGGACCCATTGGATCCGCTGTCTTCATCATCACTGTCAAATGGGTGAG
 TGGGCTTAAGAATTGGCAGGTTCCAGGAGTTAAGAATATGGTTATATATGTCGATATTTCTGTGATGTCT
 GATCCAAATCCGGCTCAGTCTTATATCTACTCCAACCTCCCCCTCATAAACTTCCATCTCCTTCATATAATC
 CCTCTTACCATCCTCTCCTTCCTGATCTCCACCGTTGGACATTTTACTTACGTCGTCACCTCTTTTATC
 TTATTATTGTCTCTGATTTTTTCTGCACTCTCTGTACTTGAGTCTAATTTTCCCATTCCTTTTAA
 CTCTCGTCATTGAATTCAGTGTGCCTTATCTGTATCACTTTAATTTCAAGCTGCAGAAGATGAAGGAGCT
 GAGGGCGGTGTATCCAGAGAAGAAAGTGTACACGCAGCAGGTCCGGTCCAGGCTGCTGCTTCGGTGCT
 CTCGCTCTGATGCTGCGTTTCTACTTTGAGGTGACAATCATCAACATAAACTAACCCAAACAAAGAA
 CTACCAGAAGATTAATCAGGGGTGGGCAGAGAGATTGTTTCTTGCACTAACAACACATTACATGTATTT
 ACTATTTTACCTTTATGAGCTCTGTAAATTTGGTTATAGAGTGACAAAAATGAGGAGTTTGATTAGGCTT
 CCCCCAACTATGTATTACACCTTTAATAAATGATATCAGCCTGAATAGAATGTGCTATATTCTCTTCACT
 GCAGGAGTGGGACTACGCCTATGTTACAGCTTCTACCATCTGTCTCTGGCTGTGTCTCTCGTCTGCT
 ACTGCCAAAGAAGAACCCTACGCAGGGACGGGACGAAACGCAGCTAAGCTTAGCTGCTTCACTCTC
 TGCTGCTGCTGAAGTTACTTTACTATATCTGTATAAGAGAGGGGAGCTGAAGCTGAAGGCAACTGGGC
 AGGTCACATGAGGGCAGGGTAGAGTTGGAGTCGTAGATTGTATGGACTAGTGAAAGCAAGAGACTCG
 GGACTGGAAGTAACATTCAATAATTTAGCGCTTAAACTGACTTTGCAATAAATCAGCATCAITCACT
 GTCATGTCAAGTCAATTCCTTAACAATGTCTAATCCTGCTTCCTCACTGTTCCATCTCCAGTCAAT
 GTCCCCCTGCTACTTCTAAAGAGAAGAAAGACAAGCCAAAGAAGAAATCGTCTCGGACCGTGTGGACA
 ATCCCCACTGAAAAGCTGTGGACACGAGGCTGTAGCACCCCAACTCTGCCTCTTTACAACCCCCCGCC
 CTCCACGCCAATCAAAGGACCCAGCATCAGCAAACTCAAAGAGATGAATGGCTGGAAGTGA

Supplementary Fig. S1. The full length of *Myomaker* genomic DNA in *Acanthopagrus latus*. Initiation and termination codons are marked by green. Exons are shown with yellow boxes.