



Transcriptome Analysis Reveals the Immune Response of Chicken Erythrocytes to Marek's Disease Virus Infection

Guan-bao Ning¹, Sheng Niu¹, Yue-jian Li^{1,2}, Xiao-xiao Lu^{1,3}, Shi-xiong Yang¹, Ali-Raza Jahejo¹, Ding Zhang¹, Wei-fang Hao⁴, Wen-wei Gao¹, Yu-jun Zhao¹, Jian-hui Li¹, Fang Yan¹, Rong-kun Gao¹, Yu-hai Bi^{1,5}, Wen-xia Tian^{1*} and Ling-xia Han^{6*}

¹College of Veterinary Medicine, Shanxi Agricultural University, Taigu 030801, China

²Weinan Animal Health Supervision Institute, Weinan 714000, China

³Wenxian Forestry Bureau, Wenxian 454850, China

⁴Taiyuan Center for Disease Control and Prevention, Taiyuan 030024, China

⁵CAS Key Laboratory of Pathogenic Microbiology and Immunology, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Disease, Institute of Microbiology, Center for Influenza Research and Early-warning (CASCIRE), Chinese Academy of Sciences, Beijing 100101, China

⁶Unit of Laboratory Animal and Comparative Medicine, State Key Laboratory of Veterinary Biotechnology, Harbin Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Harbin 150001, China

Guan-bao Ning and Sheng Niu contributed equally to this study.

ABSTRACT

Marek's disease virus (MDV) is a highly cell-associated, oncogenic alpha herpesvirus that causes Marek's disease (MD) in chickens. Chicken erythrocytes are nucleated cells of myeloid origin which contribute to gases exchange, transportation and immunity. The objective of this study was to reveal the immune responses of MDV-infected erythrocytes of chicken by transcriptome analysis. Twenty specific pathogen-free (SPF) chickens were divided into two groups as control and treatment. The chickens of treatment group were inoculated with MDV, and blood samples of two groups were obtained at 14 and 22 days post-infection (dpi) for RNA extraction. Transcriptome analysis was performed to explore the immune responses of chicken erythrocytes to MDV infection at 14 dpi and 22 dpi. In total, 129.6 million reads of two groups at two time points were mapped against the reference genome (*Gallus gallus* 5.0). A significant differential pattern of the immune response of chicken erythrocytes was identified by the GO and KEGG databases between the two phases of MDV infection. The results indicated that cytokine-cytokine receptor interaction, toll-like receptor signaling pathway, mitogen-activated protein kinases (MAPK) signaling pathway, Janus kinase/signal transducers and activators of transcription (JAK-STAT) signaling pathway were involved in the response to MDV infection. The expression levels of all most differentially expressed genes (DEGs) in these four pathways were upregulated significantly at 14 and 22 dpi, indicating that the immune responses of erythrocytes were induced by MDV. This study may help to elucidate the molecular mechanism of chicken erythrocytes relate to immune responses in MDV infection.

Article Information

Received 12 October 2019

Revised 02 January 2020

Accepted 08 January 2020

Available online 06 November 2020

Authors' Contribution

WXT, LXH and YHB contributed to conception and design of the research. GBN, SN, YJL, XXL, SXY, ARJ, DZ, WFH participated in sample collection and laboratory testing. WWG, YJZ, JHL, FY and RKG participated in the data analysis. GBN and SN drafted the basic manuscript and analyzed the data.

Key words

Marek's disease virus, Chicken, Erythrocyte, Immune response, Transcriptome

INTRODUCTION

Marek's disease (MD) is caused by Marek's disease virus (MDV) which is a lymphotropic cell-associated herpesvirus and induces the transformation of cluster of

differentiation (CD)4⁺ T cells in chicken. MD is also characterized by the infiltration of lymphoid cells into many organs such as liver, spleen and peripheral nerves (Couteaudier and Denesvre, 2014). The serotypes of MDVs were divided into three types: oncogenic MDV (such as MD5), non-oncogenic MDV (such as SB-1), herpesvirus of turkeys (HVT). These viruses have been evolving worldwide despite forty years of vaccination, so it has resulted in substantial economic losses. The pandemic strains have changed from mild MDV (mMDV) to very

* Corresponding author: wenxiantian@126.com; 11703120@qq.com
0030-9923/2021/0001-0001 \$ 9.00/0
Copyright 2021 Zoological Society of Pakistan

virulent MDV (vvMDV) and very virulent plus MDV (vv + MDV) (Boodhoo *et al.*, 2016; Hao *et al.*, 2014). The major histocompatibility complex (MHC) haplotypes of chickens is the locus with largest effect on MD genetic resistance and susceptibility, but other non-MHC genes are also thought to contribute to MD resistance through cellular interactions, variations in MDV-targeted cells, innate immunity, or cytokine regulation (McPherson and Delany, 2016). For example, the virulent strains of MDV are relatively sensitive to chickens with haplotypes B2, B5, B12, B13 and B15. In the initial phases of MDV pathogenesis, resistant and susceptible birds are similar in degree of viral replication. B-cell lysis and T-cell activation occur within 3-5 dpi and the viral latency begins around 7 to 10 dpi. (Boodhoo *et al.*, 2016). However, subsequent phases of pathogenesis and host responses differ greatly. It has been described that the susceptibility of MD is involved in virus re-activation at about 14 dpi (Boodhoo *et al.*, 2016). MDV re-activation is followed by the transformation and infiltration of CD4⁺CD8⁺ T cells, which results in immunosuppression about 21 dpi (Osterrieder *et al.*, 2006).

Erythrocytes are the most abundant cells in circulation and the most significant cells in transportation and exchange of gases. However, the immunomodulatory roles of erythrocytes during viral infections have not yet been thoroughly investigated. A recent research found that the expression of cytokines in salmon erythrocytes was induced by salmon anemia virus (Workenhe *et al.*, 2008). It has been reported that chicken erythrocyte expressed several toll-like receptors (TLRs) and upregulated the expression of some cytokines after TLR ligand treatments (Paolucci *et al.*, 2013).

Data from other groups have explored the pathogenesis of Marek's disease and immune response to MDV by the transcriptional analysis in various organs, and many differentially expressed genes between MDV infected and uninfected chickens have been identified (Sarson *et al.*, 2008; Kano *et al.*, 2009; Heidari *et al.*, 2010). For example, Heidari *et al.* (2010) have conducted the Gene Chip Chicken Genome Arrays to analyze the pattern of host gene expression in the spleen of MDV-infected chickens. Transcriptome studies have been used for exploring the host response to vaccination in MDV-infected chickens (Kano *et al.*, 2009). Further, the host responses to MDV infection in genetically resistant and susceptible chickens have also been obtained by using transcriptional analysis (Sarson *et al.*, 2008). However, there is little available information about the transcriptome expression of chicken erythrocyte in response to MDV infection. The present study aimed to elucidate the immune responses of chickens to MD by examining the transcriptional profiling in different phases of MDV infection.

MATERIALS AND METHODS

Animal infection experiments

Twenty 1-day-old specific-pathogen-free (SPF) chickens (B15 haplotype) and the MD5 strain (very virulent MDV) were kindly provided by Dr. Lingxia Han of Harbin Veterinary Research Institute. The MD5 strain was diluted by sterile phosphate buffered saline (PBS) (Solarbio, Beijing, China). The chickens were randomly divided into two groups (10 chickens/group): Each chicken of the infected group was immunized by an intra-abdominal injection of MD5 with the dose of 500 plaque-forming units (PFU)/500 μ L, and each chicken of uninfected control group was injected with the same volume of PBS. All chickens were reared in an animal biosafety level 2 laboratory, and the blood samples were collected from two groups (10 chickens each group) and pooled at 14 and 22 dpi respectively for total RNA extraction. The samples were designated as the control group (14C and 22C), and the treatment group (14T and 22T), where the sample names correspond to 14 dpi and 22 dpi respectively. The management of the experimental animals was in agreement with the welfare guidelines approved by the College of Animal Science and Veterinary Medicine of Shanxi Agricultural University, China (Number 88, 2010).

Erythrocyte isolation and RNA extraction

The erythrocytes were isolated from each blood sample collected at 14 and 22 dpi by a Histopaque-1119 solution (Sigma-Aldrich, Oakville, ON) and RNA extraction from these erythrocyte samples was performed as described previously (Niu *et al.*, 2018).

Library construction and RNA sequencing

Library construction and RNA sequencing were performed by Beijing BioMarker Technologies (Beijing, China) according to the institute's protocols. The RNA purity was checked by using the NanoPhotometer spectrophotometer (IMPLEN, CA, USA), and then RNA degradation and contamination were monitored on 1% agarose gel electrophoresis. The RNA integrity was checked using an RNA Nano 6000 Assay Kit of the Agilent Bioanalyzer 2100 system (Agilent Technologies, CA, USA). A total of 1 μ g purified RNA per sample was used as input material for the RNA library preparation. Sequencing libraries were generated using NEBNext®Ultra™ RNA Library Prep Kit for Illumina® (NEB, USA) according to the manufacturer's instructions, and index codes were added to attribute sequences to each sample. Subsequently, the libraries were sequenced on the Illumina HiSeq 2500. The detailed protocol has been described previously (Wei *et al.*, 2017). All transcriptome sequences were submitted

in the National Center for Biotechnology Information (NCBI) Sequence Read Archive under the SRA accession number SRP136403.

Quality control

Raw reads were firstly examined by using the in-house perl scripts. Clean reads were obtained by deleting the low-quality reads and the reads containing adapter or ploy-N from raw data. Then Q30, GC-content and sequence duplication level of the clean data were calculated. All the downstream analyses were based on clean data of high quality.

RNA-seq data analysis

Clean reads were mapped to the reference genome sequence (*Gallus_gallus* 5.0) by using the Tophat2 tools soft (Langmead *et al.*, 2009). Each unigene was transformed into FPKM (Fragments Per Kilo base per Million mapped reads) by using RSEM software (Li and Dewey, 2011). Gene function was annotated based on the following databases: Nr (NCBI non-redundant protein sequences) (Deng *et al.*, 2006); Pfam (Protein family) (Finn *et al.*, 2013); COG (Clusters of Orthologous Groups of proteins) (Tatusov *et al.*, 2000); Swiss-Prot (A manually annotated and reviewed protein sequence database) (Apweiler *et al.*, 2004); KEGG (Kyoto Encyclopedia of Genes and Genomes database) (Kanehisa *et al.*, 2004); GO (Gene Ontology) (Ashburner *et al.*, 2000).

Differential expression analysis

Prior to differential gene expression analysis, for each sequenced library, the read counts were adjusted by the edgeR program package through one scaling normalized factor. Differential expression analysis of two groups was performed using the DEGseq R package. P value was adjusted using q value for controlling the false discovery rate (Storey, 2003). Gene expressions with q-value < 0.005 and $|\log_2(\text{foldchange})| \geq 1$ found by DESeq were assigned as significantly differential expression.

RESULTS

Transcriptome assembly, read mapping, and annotation of new genes

The sequencing data of the 4 groups was described in Table I. The data quality assessment has been shown that quality Q30 was higher than 87.31% for each sample and the GC content was 52.48%-54.73%. 77.88%-82.64% clean reads were matched to chicken genome after filtering (Table I). The quality assessment showed that the transcriptome sequencing data was quite good and reliable. 1583 new genes were discovered that were not found or not annotated before in the reference genome. have not been

annotated in chicken reference genome. Among these, 568 genes were not present in the databases so were not annotated, and 1015 new genes were annotated according to the databases including NR, Swiss-Prot, GO, COG, and KEGG, and 568 new genes have not been annotated (Table II).

Functional annotation and classification of the DEGs between uninfected and infected groups

Total, up- and down-regulated DEGs were detected among these four groups (14T vs 14C, 22T vs 22C). The number of total DEGs (1880) was the highest between the treatment and the control group on 22 dpi, of which 1578 DEGs were up-regulated and 302 were down-regulated (Fig. 1). In the GO terms, three main functional classifications were determined from the obtained DEGs among the four groups: cellular component, biological process and molecular function (Fig. 2). KEGG pathway analysis indicated that the DEGs at each comparison could be grouped into 162 (14T vs 14C), 176 (22T vs 22C), pathways (Supplemental file 1).

Differentially expressed immune-related genes caused by MDV infection

Aiming to identify the DEGs involved in immune pathways and explore the specific changes in the expression of genes caused by MDV infection, all of the DEGs were classified into different functional categories by using GO enrichment analysis ($P < 0.05$). In the 14T/14C group (Supplemental file 2), genes related to defense response (GO: 0006952), blood vessel development (GO: 0001568), cell morphogenesis involved in neuron differentiation (GO: 0048667), response to oxygen-containing compound (GO: 1901700), blood vessel morphogenesis (GO: 0048514), regulation of mitogen-activated protein kinases (MAPK) cascade (GO: 0043408), activation of immune response (GO: 0002253), and cytokine production (GO: 0001816) were significantly enriched. In the data obtained at 22 dpi (Supplemental file 3), many genes were significantly enriched in the GO terms: the blood vessel development (GO: 0001568), response to oxidative stress (GO: 0006979), regulation of cytokine production (GO: 0001817), cytokine production (GO: 0001816), immune response (GO: 0006955), regulation of T cell mediated immunity (GO: 0002709), positive regulation of immune response (GO: 0050778), inflammatory response (GO: 0006954).

We used common expression patterns to further analyze the DEGs in the groups of 14T/14C, 22T/22C. Based on this method, all of the DEGs were placed into 56 groups, and then three immune-related groups were selected to analyze the immune response of erythrocyte to MDV infection (Fig. 3, Table III). The DEGs expression

Table I. Statistical analysis of transcriptome sequencing data.

Sample-ID	Total reads	Clean bases	GC content	%Q30	Mapped reads	Uniq mapped reads
14C	38,907,472	3,767,366,820	53.60%	89.44%	32,153,961 (82.64%)	31,438,669 (80.80%)
14T	42,900,662	4,517,615,886	53.53%	89.83%	33,955,630 (79.15%)	32,997,450 (76.92%)
22C	38,076,328	3,843,491,992	52.48%	89.98%	30,142,482 (79.16%)	29,256,084 (76.84%)
22T	41,237,630	4,163,714,940	54.73%	87.31%	33,403,166 (81.00%)	32,162,416 (77.99%)

of the three groups classified as involved in these immune-related GO terms: regulation of B cell cytokine production, NK T cell differentiation, interleukin-4 production, positive regulation of I-kappaB kinase/NF-kappaB signaling, positive regulation of interleukin-13 and interleukin-10 biosynthetic process and some other process. The genes involved in these immune-related GO terms showed increased expression in 14T and 22T as compared to 14C and 22C.

Table II. Statistical annotation of new genes.

Annotated databases	COG	GO	KEGG	Swiss-prot	Nr	All
New gene number	107	237	379	309	1004	1015

Table III. The GO terms description of three immune-related clusters in Fig. 3.

Groups	GO Term	P-value
Group 1	Regulation of B cell cytokine production	0.0001862635
	NK T cell differentiation	0.0002786086
	Regulation of platelet activation	0.0002786086
	Collagen type IV trimer	0.0003789154
	Interleukin-4 production	0.0005171461
Group 2	Cytosol	1.904526e-09
	Integrin-mediated signaling pathway	4.497712e-09
	Leukocyte migration	2.133534e-08
	Positive regulation of I-kappaB kinase/NF-kappaB signaling	1.897998e-07
	Magnesium ion binding	5.587603e-07
Group 3	Chromatin silencing at rDNA	0.0003357945
	Regulation of chromatin assembly	0.0004184276
	TAP complex	0.0008186206
	Positive regulation of interleukin-13 biosynthetic process	0.0008303057
	Positive regulation of interleukin-10 biosynthetic process	0.0008303057

The KEGG immune-related pathway mapping for DEGs

In the 14T/14C comparison, there are nine significantly enriched pathways, including endocytosis, ECM-receptor

interaction, focal adhesion, herpes simplex infection, cytokine-cytokine receptor interaction, phagosome, cell adhesion molecules, influenza A, ribosome, most of the enriched pathways were related to immune function. In the 22T/22C comparison, pathways related to the cytokine-cytokine receptor interaction, toll-like receptor signaling pathway, phagosome, calcium signaling pathway, MAPK signaling pathway, focal adhesion, JAK-STAT signaling pathway were significantly enriched.




Fig. 1. (A) Numbers of up-regulated and down-regulated DEGs between different cDNA libraries. Blue color represents up-regulated expression of DEGs; Brown color represents down-regulated expression of DEGs. **(B)** Volcano plot between different cDNA libraries. The log 2 fold represents the mean expression level. Red dots represent up-regulated genes, black dots represent no significant DEGs, the green dots represent down-regulated genes. 14T/14C, the DEGs were generated from the treatment group relative to the control group at 14 dpi; 22T/22C, the DEGs of treatment group relative to the control group at 22 dpi.




Fig. 2. Gene ontology (GO) categorization of the differentially expressed unigenes (DEG unigenes) in the comparisons (14T/14C and 22T/22C).





Fig. 3. The heat map of DEGs in three immune-related clusters. X-axis indicated different samples, and the y-axis indicated the gene expression level. Color key ranging from -2 to 2 represents $\log_2(\text{FPKM}+1)$, indicating low (green) to high (red) expression.






Fig. 5. Schematic diagrams of the toll-like receptor signaling pathway.

More importantly, the multiple occurrences of Cytokine-cytokine receptor interaction, Toll-like receptor signaling pathway, MAPK signaling pathway and many other immune-related pathways (enriched in the 14T/14C and 22T/22C) were particularly striking. These pathways may be a good starting point to characterize the difference in immune reactions during MDV infection. We focused on

the toll-like receptor signaling pathway, cytokine-cytokine receptor interaction, MAPK signaling pathway and Janus kinase/signal transducers and activators of transcription (JAK-STAT) signaling pathway (Figs. 4, 5, 6 and 7), and these pathways were enhanced for all groups. Of these, three main gene families (toll-like receptors, gallinacins, interleukins receptors) were upregulated in the 22T group

Table IV. Pathway analysis of DEGs based on the KEGG database.

Pathway term	Pathway ID	Unigene number
14T vs 14C		
Endocytosis	ko04144	23
ECM-receptor interaction	ko04512	21
Calcium signaling pathway	ko04020	17
Focal adhesion	ko04510	26
Vascular smooth muscle contraction	ko04270	12
Herpes simplex infection	ko05168	25
Cytokine-cytokine receptor interaction	ko04060	27
JAK-STAT signaling pathway	ko04630	15
Phagosome	ko04145	22
Insulin signaling pathway	ko04910	11
Toll-like receptor signaling pathway	ko04620	8
Tight junction	ko04530	14
Cell adhesion molecules (CAMs)	ko04514	20
Influenza A	ko05164	20
Ribosome	ko03010	20
Neuroactive ligand-receptor interaction	ko04080	14
RNA transport	ko03013	14
Pyrimidine metabolism	ko00240	14
Oxidative phosphorylation	ko00190	11
Regulation of actin cytoskeleton	ko04810	21
Purine metabolism	ko00230	18
Cell cycle	ko04110	15
MAPK signaling pathway	ko04010	18
22T vs 22C		
Cytokine-cytokine receptor interaction	ko04060	31
Toll-like receptor signaling pathway	ko04620	21
Carbon metabolism	ko01200	22
Phagosome	ko04145	33
Calcium signaling pathway	ko04020	29
Herpes simplex infection	ko05168	25
Endocytosis	ko04144	28
Lysosome	ko04142	21
Focal adhesion	ko04510	40
JAK-STAT signaling pathway	ko04630	28
Cell cycle	ko04110	23
FoxO signaling pathway	ko04068	27
Purine metabolism	ko00230	30
Influenza A	ko05164	28
MAPK signaling pathway	ko04010	32
Tight junction	ko04530	22
Regulation of actin cytoskeleton	ko04810	49
Cysteine and methionine metabolism	ko00270	11

Note: 14T/14C and 22T/22C means the DEGs between the control group and the treatment group at 14 and 22 dpi after MDV infection.

compared with the 22C group ([Supplemental file 4](#)). Further, the host defense peptides (GALs and CATHs and NK-lysin) were highly expressed in the MDV infection group at all comparisons.

DISCUSSION

MDV, the etiological agent of MD, is a highly cell-associated oncogenic α -herpesvirus that causes the infiltration of lymphoid cells into various organs. To provide insights into the immune responses of chicken erythrocytes to MDV infection, we investigated the dynamic changes of immune molecules in chicken erythrocytes by transcriptome analysis.

Cytokines play an important role in the regulation of immune responses and inflammation, and cytokine receptors act as transmembrane proteins to transmit a signal into the cell upon ligand binding. Many genome-wide association researches have also demonstrated that polymorphisms and mutations of cytokine receptors lead to some autoimmune disorders ([O'shea *et al.*, 2008](#)). It has been shown that high level of type I IFNs transcripts and moderate levels of *interleukin (IL)-8* transcripts were expressed in activated chicken erythrocytes, and these cytokine transcripts were up-regulated by *TLR3* and *TLR21* ligands poly I:C and CpG ODN ([Paolucci *et al.*, 2013](#)). In this study, the response of cytokines and cytokine receptors family members to MDV infection has been explored ([Fig. 4](#)), and we found that *C-X-C motif chemokine (CXCL)-12, 13* and *C-C motif chemokine (CCL)-19, 21* and *interleukin 2 receptor beta (IL-2RB)* were identified and found to be highly expressed in 14 and 22 d after MDV infection, but their expression levels in 22 dpi were lower compared with that of 14 dpi. Similarly, there were many components of cytokine-cytokine receptor interaction that were upregulated in 14 and 22 dpi. These data suggested that the expressions of cytokines and their receptors in chicken erythrocytes were induced by MDV infection and suppressed in the phase of immunosuppression. Furthermore, *vascular endothelial growth factor (VEGF)-C* and *VEGF-D* act an important role in lymphangiogenesis through the binding of *FMS-like tyrosine kinase 4 (FLT4)*, which promotes lymphatic vessel development and lymphatic endothelial cell (LEC) proliferation ([Karkkainen *et al.*, 2004](#); [Mäkinen *et al.*, 2001](#)). In this present study, the transcripts of *VEGF-C*, *VEGF-D* and their receptor *FLT4* were induced after the infection of MDV, indicating that chicken erythrocytes may participate in lymphangiogenesis by promoting these gene expressions during the development of the MDV infection. Overall, the cytokine-cytokine receptor interaction in chicken erythrocytes may play a significant role in the host immune response to MDV infection.





Fig. 6. Schematic diagrams of the JAK-STAT signaling pathway.

TLRs are a family of innate immune-recognition receptors that play important roles in recognizing molecular patterns associated with pathogen invasion, and they are expressed in various animals from protostomes to vertebrates (Jiménez-Dalmaroni *et al.*, 2016). Pathogen associated molecular patterns (PAMPs) can be recognized by TLRs that signal to the host in the presence of infection via Toll/IL-1 receptor (TIR) domain-containing adaptors, such as *MyD88* (Arleevskaya *et al.*, 2019). *TLR3*, one important member of TLRs family, plays an important role in the antiviral innate immune response (Shah *et al.*, 2016).

It has been reported that activation of *TLR3* inhibited MDV infection in chicken embryo fibroblast cells (Hu *et al.*, 2016). Similarly, our data showed that the expression of *TLR3* was identified in chicken erythrocytes and upregulated in 14 and 22 dpi, suggesting *TLR3* may recognize the double-stranded RNA (dsRNA) produced in the development of MDV infection, and then activate relate signaling pathways to limit the further spread. In addition, some other important TLR gene family members including *TLR2*, 4, 7, 8 were found to be highly expressed only in 22 dpi (Fig. 5), which indicated that these TLRs may focus on participating in

14T vs 14C



22T vs 22C




Fig. 7. Schematic diagrams of the MAPK signaling pathway.

the reactivation stage of MDV infection (Hu *et al.*, 2016). The common MyD88-dependent pathway and MyD88-independent signaling pathway can be used by TLRs to activate IRF-3 and NF- κ B, and then induce the production of type I interferons and pro-inflammatory cytokines (Ouyang *et al.*, 2018). In this study, *MyD88* was highly expressed after MDV infection in both 14 and 22 dpi, which was consistent with the results of *TLR3* expression. All the above results supported the possible defensive mechanism of chicken erythrocytes that the dsRNA produced in the development of MDV infection could be recognized by TLRs, and then the common MyD88-dependent pathway and MyD88-independent signaling pathway were used by TLRs and modulate host immune response when MDV invade into the blood.

As one of a handful of pleiotropic cascades, the JAK/STAT pathway was used to transduce a multitude of signals for a wide array of cytokines and growth factors in various animals (Morris *et al.*, 2018). Some important cellular events including cell proliferation, differentiation and apoptosis can be stimulated by JAK activation, which is critical to haematopoiesis and immune development. STATs are latent transcription factors that reside in the cytoplasm until activated. JAK/STAT signaling pathway was significantly enriched in all of three DEGs groups, suggesting an important role of the JAK/STAT signaling pathways in the immune response of erythrocytes to MDV infection. In this present study, the results showed that the JAKs including *JAK3* and *TYK2* were highly expressed after the infection of MDV in both of 14 and 22 dpi, which is consistent with the expression pattern of STATs including *STAT1*, *STAT4* and *STAT6* (Fig. 6). These results suggested that JAK/STAT cascade was activated by inducing the JAKs and STATs after MDV infection and it provides a direct mechanism to transform an extracellular signal into a transcriptional response (Niu *et al.*, 2018). In addition to JAK/STAT pathway effectors, suppressors of cytokine signaling (SOCS) family of proteins have been recognized as crucial negative regulators of JAK/STAT pathway and they also suppress cytokine signal transduction (Ghafouri-Fard *et al.*, 2018), and a negative feedback loop of the JAK/STAT circuitry can be induced by SOCS. The transcription and protein of the SOCS can be stimulated by activated STATs, and then the JAK/STAT pathway is turned off by SOCS protein binding phosphorylated JAKs and the receptors. Our data showed that SOCS had a higher expression after MDV infection in 14 and 22 dpi, and the expression level of SOCS in 22 dpi is higher than that of 14 dpi. These results support the hypothesis that MDV infection could activated JAKs by various cytokines such as IL2/3/6 and erythropoietin. Activated JAKs subsequently phosphorylated and targets many sites, including STATs, *tyrosine-protein phosphatase*

non-receptor type II (PTPN11) and *phosphoinositide-3-kinase (PI3K)*. The SOCS family of proteins can be activated by the high expression of STATs, and it may stimulate the negative feedback loop of the JAK/STAT circuitry. Mitogen-activated protein kinases (MAPKs) are evolutionarily conserved kinase modules that mediate intracellular signaling associated with fundamental cellular processes including growth, differentiation and apoptosis (Dhillon *et al.*, 2007). MAPK signaling pathways are comprised of a three-tier kinase module: MAPK kinase kinase (MAP3K), MAPK kinase (MAP2K) and MAPK (Kim and Choi, 2010). In our experiment, many DEGs including *tumor necrosis factor receptor (TNFR1)*, *interleukin 1 receptor type II (IL1R2)*, *MAP3K1*, *MAP3K3*, *MAP3K4*, *p38 MAP kinase (p38)* and *Ras* are in the MAPK signaling pathway. Of these DEGs, members of the Ras family of proteins including *K-Ras*, *H-Ras* and *M-Ras*, were upregulated by the MDV infection in 22 dpi, which indicated that Ras plays a key role in transmission of extracellular signals into erythrocytes in the development of MDV infection (Malumbres and Barbacid, 2003). The components of p38 signaling pathway, *p38*, *TNFR1* and *IL1R2* were highly expressed after the MDV infection, which suggested that *p38* could be activated through the proinflammatory cytokine receptors *TNFR1* and *IL1R2* (Kim and Choi, 2010). In addition to the DEG *MAP3K1* of 14 dpi, the expression of *MAP3K3* and *MAP3K4* were increased in 22 dpi, indicating that the JNK and p38 signaling pathways were activated after the MDV infection.

CONCLUSION

The immune responses of chicken erythrocyte to MDV infection was well explored by transcriptome analysis. Many DEGs showed significant differential expressions in the Cytokine-cytokine receptor interaction, Toll-like receptor signaling pathway, JAK/STAT signaling pathway and MAPK signaling pathway, which are responsible for immune responses to MDV infection. These results improve our understanding of MDV pathogenesis and complement our knowledge about the role of erythrocytes immune response on MDV infection mechanism in chicken.

ACKNOWLEDGEMENT

This work was supported by the National Key R and D Program (2016YFD0500800), Shanxi Scholarship Council of China (2017-073), Key scientific and technological innovation platform of Jinzhong (P171002-3), the National Natural Science Foundation of China (31072179), the Natural Science Foundation of Shanxi Province (2014011028-1), Key Technologies R and D Program of

Shanxi Province (20130311027-3), and “131” Leading Talent Project for Colleges and Universities of Shanxi Province.

Supplementary material

There is supplementary material associated with this article. Access the material online at: <https://dx.doi.org/10.17582/journal.pjz/20191012061013>

Statement of conflicts of interest

The authors have declared no conflict of interests.

REFERENCES

- Apweiler, R., Bairoch, A., Wu, C.H., Barker, W.C., Boeckmann, B., Ferro, S., Gasteiger, E., Huang, H., Lopez, R., Magrane, M. and Martin, M.J., 2004. UniProt: the universal protein knowledgebase. *Nucl. Acids Res.*, **32**: 115-119. <https://doi.org/10.1093/nar/gkh131>
- Arleevskaya, M.I., Larionova, R.V., Brooks, W.H., Bettacchioli, E. and Renaudineau, Y., 2019. Toll-like receptors, infections, and rheumatoid arthritis. *Clin. Rev. Allergy Immunol.*, **29**: 1-10. <https://doi.org/10.1007/s12016-019-08742-z>
- Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T. and Harris, M.A., 2000. Gene ontology: Tool for the unification of biology. *Nat. Genet.*, **25**: 25-29. <https://doi.org/10.1038/75556>
- Boodhoo, N., Gurung, A., Sharif, S. and Behboudi, S., 2016. Marek's disease in chickens: A review with focus on immunology. *Vet. Res.*, **47**: 119. <https://doi.org/10.1186/s13567-016-0404-3>
- Couteaudier, M. and Denesvre, C., 2014. Marek's disease virus and skin interactions. *Vet. Res.*, **45**: 36. <https://doi.org/10.1186/1297-9716-45-36>
- Deng, Y.Y., Li, J.Q., Wu, S.F., Zhu, Y.P., Chen, Y.W., He, F.C., 2006. Integrated nr database in protein annotation system and its localization. *Comput. Eng.*, **32**: 71-74.
- Dhillon, A.S., Hagan, S., Rath, O. and Kolch, W., 2007. MAP kinase signaling pathways in cancer. *Oncogene*, **26**: 3279-3290. <https://doi.org/10.1038/sj.onc.1210421>
- Finn, R.D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R.Y., Eddy, S.R., Heger, A., Hetherington, K., Holm, L., Mistry, J., Sonnhammer, E.L., 2013. Pfam: the protein families database. *Nucl. Acids Res.*, **42**: 222-2230. <https://doi.org/10.1093/nar/gkt1223>
- Ghafouri-Fard, S., Oskooei, V.K., Azari, I. and Taheri, M., 2018. Suppressor of cytokine signaling (SOCS) genes are downregulated in breast cancer. *World J. Surg. Oncol.*, **16**: 226. <https://doi.org/10.1186/s12957-018-1529-9>
- Hao, H., Li, C., Qiu, Y., Wang, F., Ai, W., Gao, J., Wei, L., Li, X., Sun, L., Wu, J. and Qin, G., 2014. Generation, transmission and infectivity of chicken MDV aerosols under experimental conditions. *Vet. Microbiol.*, **172**: 400-406. <https://doi.org/10.1016/j.vetmic.2014.06.003>
- Heidari, M., Sarson, A.J., Huebner, M., Sharif, S., Kireev, D. and Zhou, H., 2010. Marek's disease virus-induced immunosuppression: Array analysis of chicken immune response gene expression profiling. *Viral. Immunol.*, **23**: 309-319. <https://doi.org/10.1089/vim.2009.0079>
- Hu, X., Zou, H., Qin, A., Qian, K., Shao, H. and Ye, J., 2016. Activation of Toll-like receptor 3 inhibits Marek's disease virus infection in chicken embryo fibroblast cells. *Arch. Virol.*, **161**: 521-528. <https://doi.org/10.1007/s00705-015-2674-x>
- Jiménez-Dalmaroni, M.J., Gerswhin, M.E. and Adamopoulos, I.E., 2016. The critical role of toll-like receptors—from microbial recognition to autoimmunity: A comprehensive review. *Autoimmun Rev.*, **15**: 1-8. <https://doi.org/10.1016/j.autrev.2015.08.009>
- Kanehisa, M., Goto, S., Kawashima, S., Okuno, Y. and Hattori, M., 2004. The KEGG resource for deciphering the genome. *Nucl. Acids Res.*, **32**: 277-280. <https://doi.org/10.1093/nar/gkh063>
- Kano, R., Konnai, S., Onuma, M. and Ohashi, K., 2009. Microarray analysis of host immune responses to Marek's disease virus infection in vaccinated chickens. *J. Vet. med. Sci.*, **71**: 603-610. <https://doi.org/10.1292/jvms.71.603>
- Karkkainen, M.J., Haiko, P., Sainio, K., Partanen, J., Taipale, J., Petrova, T.V., Jeltsch, M., Jackson, D.G., Talikka, M., Rauvala, H. and Betsholtz, C., 2004. Vascular endothelial growth factor C is required for sprouting of the first lymphatic vessels from embryonic veins. *Nat. Immunol.*, **5**: 74. <https://doi.org/10.1038/ni1013>
- Kim, E.K. and Choi, E.J., 2010. Pathological roles of MAPK signaling pathways in human diseases. *Biochim. biophys. Acta Mol. Basis Dis.*, **1802**: 396-405. <https://doi.org/10.1016/j.bbadis.2009.12.009>
- Langmead, B., Trapnell, C., Pop, M. and Salzberg, S.L., 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome*

- Biol.*, **10**: R25. <https://doi.org/10.1186/gb-2009-10-3-r25>
- Li, B., Dewey, C.N., 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinform.*, **12**: 323. <https://doi.org/10.1186/1471-2105-12-323>
- Malumbres, M. and Barbacid, M., 2003. RAS oncogenes: the first 30 years. *Nat. Rev. Cancer*, **3**: 459. <https://doi.org/10.1038/nrc1097>
- Mäkinen, T., Veikkola, T., Mustjoki, S., Karpanen, T., Catimel, B., Nice, E.C., Wise, L., Mercer, A., Kowalski, H., Kerjaschki, D. and Stacker, S.A., 2001. Isolated lymphatic endothelial cells transduce growth, survival and migratory signals via the VEGF-C/D receptor VEGFR-3. *EMBO J.*, **20**: 4762-4773. <https://doi.org/10.1093/emboj/20.17.4762>
- McPherson, M.C. and Delany, M.E., 2016. Virus and host genomic, molecular, and cellular interactions during Marek's disease pathogenesis and oncogenesis. *Poult. Sci.*, **95**: 412-429. <https://doi.org/10.3382/ps/pev369>
- Morris, R., Kershaw, N.J. and Babon, J.J., 2018. The molecular details of cytokine signaling via the JAK/STAT pathway. *Protein Sci.*, **27**: 1984-2009. <https://doi.org/10.1002/pro.3519>
- Niu, G.J., Xu, J.D., Yuan, W.J., Sun, J.J., Yang, M.C., He, Z.H., Zhao, X.F. and Wang, J.X., 2018. Protein inhibitor of activated STAT (PIAS) negatively regulates the JAK/STAT pathway by inhibiting STAT phosphorylation and translocation. *Front. Immunol.*, **9**: 2392. <https://doi.org/10.3389/fimmu.2018.02392>
- Niu, S., Jahejo, A.R., Jia, F.J., Li, X., Ning, G.B., Zhang, D., Ma, H.L., Hao, W.F., Gao, W.W., Zhao, Y.J. and Gao, S.M., 2018. Transcripts of antibacterial peptides in chicken erythrocytes infected with Marek's disease virus. *BMC Vet. Res.*, **14**: 363. <https://doi.org/10.1186/s12917-018-1678-7>
- O'shea, J.J. and Murray, P.J., 2008. Cytokine signaling modules in inflammatory responses. *Immunity*, **28**: 477-487. <https://doi.org/10.1016/j.immuni.2008.03.002>
- Osterrieder, N., Kamil, J.P., Schumacher, D., Tischer, B.K., Trapp, S., 2006. Marek's disease virus: From miasma to model. *Nat. Rev. Microbiol.*, **4**: 283. <https://doi.org/10.1038/nrmicro1382>
- Ouyang, M.Z., Zhou, D., Zhu, Y., Zhang, M. and Li, L., 2018. The inhibition of MyD88 and TRIF signaling serve equivalent roles in attenuating myocardial deterioration due to acute severe inflammation. *Int. J. mol. Med.*, **41**: 399-408. <https://doi.org/10.3892/ijmm.2017.3239>
- Paolucci, S., Barjesteh, N., Wood, R.D. and Sharif, S., 2013. Chicken erythrocytes respond to Toll-like receptor ligands by up-regulating cytokine transcripts. *Res. Vet. Sci.*, **95**: 87-91. <https://doi.org/10.1016/j.rvsc.2013.01.024>
- Rawlings, J.S., Rosler, K.M. and Harrison, D.A., 2004. The JAK/STAT signaling pathway. *J. Cell Sci.*, **117**: 1281-1283. <https://doi.org/10.1242/jcs.00963>
- Sarson, A.J., Parvizi, P., Lepp, D., Quinton, M. and Sharif, S., 2008. Transcriptional analysis of host responses to Marek's disease virus infection in genetically resistant and susceptible chickens. *Anim. Genet.*, **39**: 232-240. <https://doi.org/10.1111/j.1365-2052.2008.01710.x>
- Schat, K.A. and Xing, Z., 2000. Specific and nonspecific immune responses to Marek's disease virus. *Dev. Comp. Immunol.*, **24**: 201-221. [https://doi.org/10.1016/S0145-305X\(99\)00073-7](https://doi.org/10.1016/S0145-305X(99)00073-7)
- Shah, M., Anwar, M.A., Kim, J.H. and Choi, S., 2016. Advances in antiviral therapies targeting Toll-like receptors. *Expert Opin. Investig. Drugs.*, **25**: 437-453. <https://doi.org/10.1517/13543784.2016.1154040>
- Storey, J.D., 2003. The positive false discovery rate: a Bayesian interpretation and the q-value. *Ann. Stat.*, **31**: 2013-2035. <https://doi.org/10.1214/aos/1074290335>
- Tatusov, R.L., Galperin, M.Y., Natale, D.A. and Koonin, E.V., 2000. The COG database: A tool for genome-scale analysis of protein functions and evolution. *Nucl. Acids Res.*, **28**: 33-36. <https://doi.org/10.1093/nar/28.1.33>
- Wei, J., Fan, S., Liu, B., Zhang, B., Su, J. and Yu, D., 2017. Transcriptome analysis of the immune reaction of the pearl oyster *Pinctada fucata* to xenograft from *Pinctada maxima*. *Fish Shellf. Immunol.*, **67**: 331-345. <https://doi.org/10.1016/j.fsi.2017.06.030>
- Workenhe, S.T., Kibenge, M.J., Wright, G.M., Wadowska, D.W., Groman, D.B. and Kibenge, F.S., 2008. Infectious salmon anaemia virus replication and induction of alpha interferon in Atlantic salmon erythrocytes. *J. Virol.*, **5**: 36. <https://doi.org/10.1186/1743-422X-5-36>
- Yamamoto, M., Sato, S., Hemmi, H., Hoshino, K., Kaisho, T., Sanjo, H., Takeuchi, O., Sugiyama, M., Okabe, M., Takeda, K. and Akira, S., 2003. Role of adaptor TRIF in the MyD88-independent toll-like receptor signaling pathway. *Science*, **301**: 640-643. <https://doi.org/10.1126/science.1087262>