

Short Communication

Characterization and Phylogenetic Analysis of the Mitochondrial Genome of *Clangula hyemalis* (Linnaeus, 1758)

Hengjiu Tian¹, Yan Liu², Rigude Bu³, Yaqian Fan⁴, Lin Li⁴, Shumiao Zhang⁵, Xiangnan Sun¹, Muhammad Younis Laghari⁶ and Meng Meng^{7,8*}

¹Beijing Wildlife Rescue and Rehabilitation Center, Beijing, 100029, China

²Beijing Key Laboratory of Captive Wildlife Technologies, Beijing Zoo, Beijing, 100044, China

³Academy of Inventory and Planning, National Forestry and Grassland Administration, Beijing, 100714, China

⁴Beijing Songshan National Nature Reserve, Beijing, 102115, China

⁵Beijing Milu Ecological Research Center, Beijing, 100076, China

⁶Department of Freshwater Biology and Fisheries, University of Sindh Jamshoro 76080-Pakistan

⁷China Wildlife Conservation Association, Beijing, 100029, China

⁸College of Wildlife and Protected Area, Northeast Forestry University, Harbin, 150040, China

ABSTRACT

The long-tailed duck (*Clangula hyemalis*) is a migratory bird and enduring rapid population declines, as extreme climatic alteration, and other reasons. It is also rare passage migrant into China. The mitochondrial genome (mitogenome) can provide information for phylogenetic analyses and evolutionary biology. In the present investigation, we generated complete mitogenome of the long-tailed duck using sanger sequencing. The mtDNA comprised 16,639 base pairs (bp), and contained 13 protein coding genes, 22 transfer RNAs and 2 ribosomal RNAs. Remarkably, this is the second and most complete published mitogenome for the *Clangula hyemalis* species. Our findings revealed that control region and ND4L would be expected to be as diversity fragments for identifying population-specific marker in *Clangula hyemalis*.

Article Information

Received 11 August 2023

Revised 05 June 2024

Accepted 19 June 2024

Available online 23 June 2025
(early access)

Authors' Contribution

HT and XS collected the sample. HT and MM designed the study. HT, LB, YF, LL and SZ analyzed the data. HT and MM wrote the paper. All authors read, edited, and approved the final manuscript.

Key words

Clangula hyemalis, Phylogenetic, Beijing, Mitochondrial genome, Population, Climate

The long-tailed duck (*Clangula hyemalis*) is one of the unique migratory birds which is the only living member of its genus *Clangula*. The wintering population of this sea duck is undergoing rapid population declines across Europe, as oil or other pollution and extreme climatic change, etc., (Dickson and Smith, 2013; Day *et al.*, 2015). Thus, it is classified as vulnerable species on the IUCN global red list of threatened species long-tailed ducks breed in the circumpolar Arctic and migrate to cold,

temperate waters in the non-breeding period (Robertson and Savard, 2002). The distribution of scarce long-tailed duck has been found in Central and Southwest of China for years, driven by climatic factors over the past years (Wu and Shi, 2016). Despite the low diversity for breeding Long-tailed Duck, the wild individuals from China were not published. Mitochondrial complete genome still be needed to provide more evidence in order to study genetic diversity and migratory connectivity in this species (Mackenzie *et al.*, 2022). Here, we described the mitogenome of wild-collected Long-tailed duck from China to provide basic genetic information about this species.

Material and methods

In 2020, we collected a wild migrating long-tailed duck from Beijing, China, (116°18'12"N and 40°10'11"E). A specimen was deposited at Beijing Wildlife Rescue and Rehabilitation Central Lab (contact person: Hengjiu Tian, hengjiutian@126.com, Beijing, China) under the voucher

* Corresponding author: mengmaoxian@163.com
0030-9923/2025/0001-0001 \$ 9.00/0



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

number BWRR-Ch-2020-001. The genomic DNA of each sample was extracted from the feature with TIANamp Genomic DNA Kit.

To get the complete mitochondrial genome, we selected custom-developed primers and minor gaps were resolving by overlapping primer-based PCR for DNA sequence walking (Table 1). All the purified PCR products were sequenced with sanger sequencing mode on an ABI 3730 XL machine. The mitochondrial genome was assembled following Xue *et al.* (2013), and then annotated using MitoAnnotator (Bernt *et al.*, 2013). The map of mitochondrial genome was drawing in <https://irscope.shinyapps.io/Chloroplast/>.

Results and discussion

The complete mitogenome of *Clangula hyemalis* resulted 16,639 bp in length, including 13 protein-coding genes (PCGs), 22 transfer RNAs, 2 ribosomal RNAs, and a putative control region (D-loop region), shown in Figure 1. Rearrangement and of duplication mitochondrial genome were not observed in this species. The nucleotide composition of mitogenome of *Clangula hyemalis* 29.1% for A, 33.2% for C, 15.8% for G, and 21.9% for T, with a slightly higher A+T content (51%). Only ND6 gene is encoded on the light strand, while other genes are distributed in the heavy strand. Among the 13 PCGs, only COX1, COX2, and ND5 uses GTG as the start codon and the rest genes initiate with the start codon of ATG.




Fig. 1. The complete mitogenome structure of *Clangula hyemalis*.

A maximum likelihood analysis was carried out using MEGA X package (Kumar *et al.*, 2016) with GTR+G model and 1,000 bootstrap replicates. We took two *Clangula hyemalis* mtgenomes (one is published with NCBI

accession number MW077850.2 and MW849278.1) and the other eighteen birds from GenBank. The phylogenetic analysis confirms that the Beijing's long-tailed duck close phylogenetic position to North American's one (Fig. 2).

Table 1. Primers used for amplification of *Clangula hyemalis* mitogeno

Primers name	Sequence
<i>co1</i>	F= AATGTAATCGTCACCGCCCA R= TACAGGATTGGGTCTCCCCC
<i>cytb</i>	F= ATCCTGACAGGTCTCTGTCT R= TACGAGGAGGTTGGCCACTA
<i>dloop</i>	F= GCTCCTACTCATACTACCCTGC R= CAGTGTCAAGGTGATTCCCCA
<i>nd2</i>	F= CGCAITGGGCACAACAATCA R= ATAGTGAGGCGAGGATTGCG
<i>nd3-nd4</i>	F= CCCCCGTACGAATGTGGATT R= GGAGGCAGATTGAGCTGGTT
<i>nd1-3</i>	F= AATCTGTGAAAGGAACCTCGGC R= ATCATAGGATTGAGTAGACGGC
<i>16S</i>	F= GTGATAGCTGGTT(A/G)CC(C/T)G R= AAG(C/T)TCCACAGGGTCTTCTCG
<i>CO3</i>	F= ATGGCCCACCAAGCACACTC R= CAATATCAGGCTGCTGCTTC
<i>ND3</i>	F= AAACCT(A/C/G)TCCCC(A/G/C/T)TACGAATG R= ATTCTGCTCATTCTAGGC
<i>ND4</i>	F= TAGCAAGCCAAAACACCT R= GATCAGTGGGTTTGGATTAT
1	F= CAAAGTGGCATCTGTGGAAT R= CTGGCACAAGATTTACCGAC
2	F= AAAGAGCCCATTCGACC(C/T)TC R= GTTGGGTGGTTGAGCTGTGT
3	F= GACTCATCATCCAAGAGCTA R= GGATATACGGTTCAGCCTGT
4	F= AAACCTCCCGCACTCTCACA R= GACAACATCCCGTCATCATT
5	F= GATTCCACGGACTCCACGTA R= GGTTCAGGCGAGGGTTAGTA
6	F= AGA(A/G)CCAAACCCCTCATTC R= AAAATAGCGCTAGTTCTGTG
7	F= CGCTCAATCCCAAACAAACT R= ATGGTATGTCCGCTGTAT

Compared with mitogenome of *Clangula hyemalis* from North American, we found 99.95% identify to the one from Beijing of China. In further, we found 14 SNPs sites and two indels sites between these

two mitogenomes (Table II). In a conclusion, the genetic diversity of *Clangula hyemalis* was lowest, and control region and ND4L would be expected to be as diversity fragments for identifying population-specific marker in *Clangula hyemalis*.




Fig. 2. A phylogenetic tree of *Clangula hyemalis* and the other eighteen species.

Table II. Genetic diversity of *Clangula hyemalis* mitogenome from China and American.

SNP/ InDel	No.	Gene name	Type
INDEL	1	Control region	34bp
SNP	4	Control region	C/T, C/T, G/A, T/C
SNP	2	Control region	A/G
SNP	1	ND1	G/A
SNP	1	tRNA-Cys	T/A
SNP	1	COXI	A/G
INDEL	1	COXI	C
SNP	1	tRNA-Arg	A/R
SNP	3	ND4L	T/C, T/C, A/G
SNP	1	ND6	A/G

DECLARATIONS

Acknowledgement

We thank Xiaoping Wang for picture providing

throughout our project.

Funding

This work was supported by Beijing Financial Project, grant No. 23CB066.

Disclosure statement

No potential conflict of interest was reported by the author(s). The authors comply with the International Union for Conservation of Nature (IUCN) policies research involving species at risk of extinction, the Convention on Biological Diversity and the Convention on the Trade in Endangered Species of Wild Fauna and Flora.

Data availability statement


The genome sequence data by Sangon sequencing that support the findings of this study are openly available in GenBank of NCBI at (<https://www.ncbi.nlm.nih.gov>) (<https://www.ncbi.nlm.nih.gov/>) under the accession no.MW077850.2.

Statement of conflict of interest

The authors have declared no conflict of interest.

References

- Bernt, M., Donath, A. and Jühling F., 2013. *Mol Phylogenet Evol.*, **69**:313-319.
- Bird Life International, 2014. *The IUCN red list of threatened species*. www.iucnredlist.org.
- Day, R.H., Rose, J.R., Prichard, A.K. and Streever, B., 2015. *Island Arctic Alaska*, **68**: 367-379. <https://doi.org/10.14430/arctic4507>
- Dickson, D.L. and Smith, P.A., 2013. *J. Wildl. Manage.*, **77**: 777-790. <https://doi.org/10.1002/jwmg.529>
- Counihan, K.L., Skerratt, L.F., Franson, J.C. and Hollmén, T.E., 2015. *Virology*, **485**: 393-401. <https://doi.org/10.1016/j.virol.2015.07.026>
- Kumar, S., Stecher, G., Li, M., Knyaz, C. and Tamura, K., 2018. *Mol. Biol. Evol.*, **35**: 1547-1549.
- Mackenzie, C.L., Kent-Flora, E.A., Baxter, J.M., Gormley, K.S.G., Cassidy, A.J., Sanderson, W.G. and Porter, J.S., 2022. *Front. Mar. Sci.*, **9**. <https://doi.org/10.3389/fmars.2022.772259>
- Robertson, G.J. and Savard, J.P.L., 2002. Long-tailed duck (*Clangula hyemalis*). In: *The birds of North America Online* A. Poole, Ed. <http://bna.birds.cornell.edu/bna/species/651> <https://doi.org/10.2173/bna.651>
- Wu, J. and Shi, Y., 2016. *Ecol. Inf.*, **31**: 147-155. <https://doi.org/10.13001/1081-3810.3202>
- Xue W., Hou, G., Li, C., Kong, X., Zheng, X., Li, J. and Sun, X., 2013. *Mitochondrial DNA*, **24**: 339-341. <https://doi.org/10.3109/19401736.2012.760081>



Supplementary Material

Characterization and Phylogenetic Analysis of the Mitochondrial Genome of *Clangula hyemalis* (Linnaeus, 1758)

Hengjiu Tian¹, Yan Liu², Rigude Bu³, Yaqian Fan⁴, Lin Li⁴, Shumiao Zhang⁵, Xiangnan Sun¹, Muhammad Younis Laghari⁶ and Meng Meng^{7,8*}

¹Beijing Wildlife Rescue and Rehabilitation Center, Beijing, 100029, China

²Beijing Key Laboratory of Captive Wildlife Technologies, Beijing Zoo, Beijing, 100044, China

³Academy of Inventory and Planning, National Forestry and Grassland Administration, Beijing, 100714, China

⁴Beijing Songshan National Nature Reserve, Beijing, 102115, China

⁵Beijing Milu Ecological Research Center, Beijing, 100076, China

⁶Department of Freshwater Biology and Fisheries, University of Sindh Jamshoro 76080-Pakistan

⁷China Wildlife Conservation Association, Beijing, 100029, China

⁸College of Wildlife and Protected Area, Northeast Forestry University, Harbin, 150040, China

Table 1. Primers used for amplification of *Clangula hyemalis* mitogenome.

Primers name	Sequence
<i>col</i>	F= AATGTAATCGTCACCGCCCA R= TACAGGATTGGGTCTCCCCC
<i>cytb</i>	F= ATCCTGACAGGTCTCCTGCT R= TACGAGGAGGTTGGCCACTA
<i>dloop</i>	F= GCTCCTACTCATACTACCCTGC R= CAGTGTC AAGGTGATTCCCCA
<i>nd2</i>	F= CGCATTGGGCACAACAATCA R= ATAGTGAGGCGAGGATTGCG
<i>nd3-nd4</i>	F= CCCCCGTACGAATGTGGATT R= GGAGGCAGATTGAGCTGGTT

Table continued on next colum.....

Primers name	Sequence
<i>nd1-3</i>	F= AATCTGTGAAAGGAACCTCGGC R= ATCATAGGATTGAGTAGACGGC
<i>16S</i>	F= GTGATAGCTGGTT(A/G)CC(C/T)G R= AAG(C/T)TCCACAGGGTCTTCTCG
<i>CO3</i>	F= ATGGCCCACCAAGCACACTC R= CAATATCAGGCTGCTGCTTC
<i>ND3</i>	F= AAACCT(A/C/G)TCCCC(A/G/C/T)TACGAATG R= ATTCTGCTCATTCTAGGC
<i>ND4</i>	F= TAGCAAGCCAAAACCACTT R= GATCAGTGGGTTTGGATTAT
1	F= CAAAGTGGCATCTGTGGAAT R= CTGGCACAAGATTTACCGAC
2	F= AAAGAGCCCATTCGACC(C/T)TC R= GTTGGGTGGTTGAGCTGTGT
3	F= GACTCATCATCCAAGAGCTA R= GGATATACGGTTCAGCCTGT
4	F= AAACCTCCCGCACTCTCACA R= GACAACATCCCGTCATCATT
5	F= GATTCCACGGACTCCACGTA R= GGTTCAGGCGAGGGTTAGTA
6	F= AGA(A/G)CCAAACCCCTCATTC R= AAAATAGCGCTAGTTCTGTG
7	F= CGCTCAATCCCAAACAACT R= ATGGTATGTCCGCTTGTAT

* Corresponding author: mengmaoxian@163.com
0030-9923/2025/0001-0001 \$ 9.00/0



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Table II. Genetic diversity of *Clangula hyemalis* mitogenome from China and American.

SNP/ InDel	No.	Gene name	Type
INDEL	1	Control region	34bp
SNP	4	Control region	C/T, C/T, G/A, T/C
SNP	2	Control region	A/G
SNP	1	<i>ND1</i>	G/A
SNP	1	<i>tRNA-Cys</i>	T/A
SNP	1	<i>COXI</i>	A/G
INDEL	1	<i>COXI</i>	C
SNP	1	<i>tRNA-Arg</i>	A/R
SNP	3	<i>ND4L</i>	T/C, T/C, A/G
SNP	1	<i>ND6</i>	A/G

Online First Article