



Foraging Behavior and Nutritional Assessment of Selected Forage Species for Captive Blackbuck (*Antelope cervicapra*) at Lal Suhanra National Park, Bahawalpur, Pakistan

Muhammad Azeem Akhter^{1,2}, Muhammad Talha Imtiaz^{1,2}, Ihsan Qadir^{1*}, Mark Edwin Hostetler², Ghulam Yasin¹, Imran Khan^{1,3}, Nasir Qadir⁴, Nazish Iftikhar¹ and Jahanzaib⁵

¹Department of Forestry and Range Management, Bahauddin Zakariya University, Multan, Pakistan

²Wildlife Ecology and conservation, University of Florida, Gainesville, FL, USA

³Department of Forestry and Range Management, Gomal University, Dera Ismail Khan, KP, Pakistan.

⁴Warnell School of Forestry and Natural Resources, University of Georgia, Athens, GA, USA

⁵Department of Renewable Resources, University of Alberta, Canada

ABSTRACT

This study aimed to assess the feeding preference of blackbuck *Antelope cervicapra* and the nutritional analysis in a breeding enclosure at Lal Suhanra National Park, district Bahawalpur, Pakistan, from May 2021 to April 2022. The feeding preference of 180 (60 males, 60 females and 60 fawns) blackbuck was evaluated through cafeteria-type trials with 15 palatable species, including five trees, five shrubs, and five grasses. There were significant differences in feeding preference among different gender groups, with *Salvadora oleoides* and *Prosopis cineraria* being the most preferred tree species correlated with high values of crude fiber (CF) and dry matter (DM). Male blackbuck preferred *Vachellia nilotica*, while female blackbuck preferred *Zizyphus mauritiana* correlated with DM, CF, ether extract (EE), and total ash (TA), neutral detergent fiber (NDF), acid detergent fiber (ADF), hemicellulose (HC) and lignin (L) concentration. *Tamarix aphylla* was the least preferred species for all genders. *Haloxylon salicornicum* was the most preferred shrub species, again correlated with DM, NDF, HC and L concentration but did not have the highest concentration, while *Prosopis cineraria* was the least preferred. *Cenchrus pennisetiformis* was the most preferred species among grasses, correlated with high concentration of DM, HC, and L. The DM content was found to vary significantly among the different plant species studied, with the highest value observed in *Prosopis cineraria* at 94.5%. The highest content of CP was in *Vachellia nilotica* at 10.56%. The maximum content of CF was found in *Prosopis cineraria* at 30.04%. The nutritional content of selected shrub species' DM content ranged from 91.56% to 95.24%, with *Suaeda fruticosa* having the highest value (95.24%). The highest concentration of CP was found in *Suaeda fruticosa* (16.90%) with the highest nutritive value among the selected shrubs. Among the selected grass species, the highest DM content was found in *Cenchrus pennisetiformis*. The highest concentration of CF was found in *Lasiurus scindicus*, while *Panicum antidotale* had the highest value for CP content, ranging from 3.9% to 5.8%. Overall, there was a weak correlation between forage preference and the nutritive potential of selected species. However, blackbuck showed a preference for species that are higher in DM, NDF, HC and L.

Article Information

Received 08 August 2023

Revised 15 April 2025

Accepted 28 April 2025

Available online 22 October 2025
(early access)

Authors' Contribution

Conceptualization: IQ, MTI, MAA. Methodology: MAA, NI, MTI, GY, IK. Investigation: IQ, MAA. Visualization: MTI, MAA, GY. Funding acquisition: MAA, MTI, IQ, J. Supervision: IQ, MEH. Writing original draft: MAA, MTI, GY. Writing review and editing: IQ, GY, MTI, MEH, J.

Key words

Feeding preferences, Nutritional analysis, Dry matter, Ether extract, Neutral detergent fiber, Lignin, Hemicellulose

INTRODUCTION

Large herbivores living in heterogeneous environments face significant challenges in acquiring forage (Dostaler *et al.*, 2011). They must balance the potential for energy and nourishment against competition and predator attack, exposure to plant poisons, or conspecific animosity (Felton *et al.*, 2020). The amount of nutrients in different forage species and plant sections varies substantially (Dostaler *et al.*, 2011). Even if an animal may consume

* Corresponding author: drbhabha@bzu.edu.pk
0030-9923/2025/0001-0001 \$ 9.00/0



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

plants as they are available, larger herbivores require a higher nutritional diet and fewer secondary metabolites (Stutz *et al.*, 2018). Therefore, large herbivores avoid random feeding and use only those plants or parts of plants that best satisfy their metabolic requirements (Balluffi-Fry *et al.*, 2022). The forage species as well as individual parts of the plant may vary greatly in terms of nutrient concentration (Dostaler *et al.*, 2011).

The blackbuck antelope *Antelope cervicapra* was extinct in the wild from Cholistan by 1967 (Aleem, 1978), the only habitat in the wild where it occurred in Pakistan. Despite the minimal impact of urbanization and cultivation due to limited irrigation water and scanty rainfall in that area, the primary cause of its extermination was illegal hunting and poaching (Abid and Tasleem, 2022). However, the worldwide population of blackbuck is the “Least Concern”, but the population is declining under the high pressure of urbanization, commercialization, and over-exploitation (IUCN, 2017). Because of their iconic regional importance, blackbuck was re-introduced from Texas, U.S. in the late 1970s via captive breeding method (Khan *et al.*, 2023). Blackbuck is currently maintained in captivity in several wildlife parks, breeding centers, and zoos, both in private and public sector facilities (Farooq *et al.*, 2022). Breeding animals in captivity to release them back into the wild later is one of the most effective strategies for conserving a species in the wild (Wakechaure and Ganguly, 2016). For this reason, all wildlife and forest departments in Pakistan are working together to restore the iconic species of the Cholistan Desert using the restoration method of captive breeding (Farooq *et al.*, 2022).

The blackbuck is a gazelle-shaped antelope, that belongs to the family Bovidae subfamily Antilopinae, tribe Antilopini (Choudhary and Chisty, 2022). Four subspecies of *Antelope cervicapra* L. have been reported viz. Southeastern blackbuck (*A. cervicapra cervicapra*), Pakistan and northwestern blackbuck (*A. cervicapra rajputanae*), *A. cervicapra centralis* is from central India, and *A. cervicapra rupicapra* found in Nepal and North India (Csurhes and Fisher, 2010; Khalil *et al.*, 2020). The blackbuck is native to South Asia, primarily India, Pakistan, Nepal, and Bangladesh (Choudhary and Chisty, 2022; Tahir, 2022). It has been introduced to other regions, including Texas in the 1930s, where its population grew from 1,000 in 1955 to over 7,000 by 1980 (Wright, 1985). The species’ ability to inhabit various environments, from grasslands to scrublands, contributes to its survival, but continued conservation efforts are crucial for its long-term preservation (Choudhary and Chisty, 2022; Tahir, 2022).

There is noticeable sexual dimorphism in this ungulate species. The term “blackbuck” refers to the male’s dark brown to blackish coat hue (Meena and Saran, 2018). The

length of the whorled horns of males is 79 cm, but none are present in females (Csurhes and Fisher, 2010). The species tail measures 10–17 cm in length, while its body is typically between 100 and 150 cm long (Khalil *et al.*, 2020). The average body weight of males is 19.5–56.7 kg and of females, 19–33 kg. The average recorded lifespan in general is 10 to 15 years (Csurhes and Fisher, 2010). The highest recorded age is 12 to 13 years for males and up to 16 years for females (Tahir, 2022).

Blackbuck was most abundant in the Cholistan Desert in 1950s but due to overexploitation, they became extinct in the wild (Aleem, 1978). Their population was extremely vulnerable to habitat loss, lack of forage, and under considerable stress from overhunting, which may be the reason for their extinction in the wild (Meena and Saran, 2018). In 1970 WWF International arranged a donation of 10 blackbuck, three males and seven females, from Texas ranches (through San Antonio Zoo) where they were introduced from Cholistan in 1930s. Lal Suhanra irrigated forest plantation was selected as the venue for the introduction (Chambers, 1996). The animals were initially kept in a small enclosure covering an area of about one hectare. In 1973 they were transferred to a bigger enclosure, a 3-meter-high wire netting fence 10 cm mesh made of 8-gauge galvanized wire (donated by children of Holland, arranged by WWF International) was erected in a perimeter of about 10 km covering 288 hectares. Wire netting was also extended in the ground up to a depth of 0.6 meter to keep the burrowing animals away from the enclosure (Aleem, 1978).

The blackbuck breeds all the year round but the main rut takes place in February to March and each female bears one to two young (Prater and Barruel, 1971). In the absence of any predators in the enclosure, the population should have greatly increased in seven years in 1978, when the study was conducted; only 22 blackbuck were present in the enclosure (Aleem, 1978). The total population of blackbuck in the Blackbuck Breeding Enclosure RD-25 Lal Suhanra National Park in March 2021 was 324 (Khan *et al.*, 2023). In terms of feeding preference, knowledge of a species’ behavior is crucial for its conservation. Therefore, plant availability and foraging preference are crucial to keeping the population of these animals in the wild (Davies *et al.*, 2012). Since an animal in captivity may exhibit behavior that differs from that in the wild, it is always preferable to examine an animal in its natural habitat (Vats and Bhardwaj, 2009; Jhala and Isvaran, 2016). However, blackbuck in Pakistan still could not be reintroduced in the wild therefore, studying them in captivity is the only choice (Farooq *et al.*, 2022). The primary objective of the current study was to determine the palatability and nutritional potential of different plant species that blackbuck can

access in their native range. This research can help maintain animals in captivity and help design effective restoration of natural habitats for the reintroduction of blackbuck and other ungulates into the wild.

MATERIALS AND METHODS

Study area

The study was conducted in RD-25 Blackbuck Breeding enclosure, Lal Suhanra National Park, Bahawalpur latitude 29°22'49.89" N and longitudes 71°58'6.24"E, at altitudes from 125 to 140 meters above sea level, located in the southern part of Punjab, Pakistan (Fig. 1) from May 2021 to April 2022. Lal Suhanra National Park spans around 65,790.36 hectares (Wariss *et al.*, 2014), whereas the RD- 25 Blackbuck Breeding enclosure is 7.95 hectares in size.




Fig. 1. Map of the study area, RD-25 Blackbuck breeding enclosure, Lal Suhanra National Park, Bahawalpur.

The study site experiences a subtropical continental climate, which is marked by hot temperatures, little to no humidity, strong summer winds, and a high rate of evaporation. The hottest months are May and June, with temperatures ranging from 50°C in the summer to -2°C

in the winter. The annual precipitation range is 90 to 200 mm, and the relative humidity is usually around 60%. The summer wind comes from the southeast, whereas the winter wind comes from the northeast (Wariss *et al.*, 2014).

Feeding preference trials

Forage preference trials were conducted in a cafeteria-type style, and the data were collected from May 2021 to April 2022. For the trials, a temporary pen of 60m² was erected with an adjacent barn. Based on knowledge from literature studied and consulting wildlife keepers, we selected 15 palatable plant species [five species of trees (kikar/babul *Vachellia nilotica*, kandi *Prosopis cineraria*, farsh/okan *Tamarix aphylla*, beri *Zizyphus mauritiana*, wan/peelu *Salvadora oleoides*), five species of shrubs (buee *Suaeda fruticosa*, aak *Calotropis procera*, sajji *Haloxylon recurvum*, sajji *Haloxylon salicornicum*, dela *Capparis decidua*) and five species of grasses (khawi *Cymbopogon jwarancusa*, buffel grass *Cenchrus pennisetiformis*, ghamur/ blue panic grass *Panicum antidotale*, sewan grass *Lasiurus scindicus*, lumb *Aristida depressa*)] that may be present all the year round. We collected these plants growing in the Lal Suhanra National Park. Ten kilograms each of all respective forage species were collected and placed in each pile and fed into a feeder divided into 15 trays made in the pen to conduct the feeding trials. We randomly selected 60 male blackbuck, 60 females, and 60 fawns below the age of three years from the enclosure for the investigation of feeding preference. Each gender class underwent individual tests for each forage species for 60 min, following the protocol by Mirza and Waiz (1973). The number of visits and average time spent on each pile was recorded in each trial using a high-definition camcorder to minimize human disturbance and record precise time measurements. The feeding trials experiment was conducted five times each season. Food preference was computed by the total time spent and the total number of visits to each tray (Kravetz and Suárez, 1998).

Nutritional analysis

The samples for nutritional analysis of selected browse species were collected from the Cholistan Desert in August 2021. Each sample was collected at a fresh weight of 500 grams. Most of the samples were a combination of twigs, leaves, and inflorescence. Following collection, samples were gathered and allowed to air dry in the shade. The samples were pulverized for laboratory examination using a Wiley mill with a 2 mm sieve tube. Samples were ground and then stored in plastic bags for later analysis.

Nutritional contents including dry matter (DM), crude protein (CP), crude fiber (CF), ether extract (EE), and total ash (TA) were analyzed proximally using the procedures

outlined in the Association of Official Analytical Chemists' Standard Protocol (AOAC, 2005). Hemicellulose (HC) was evaluated by computing the difference between neutral detergent fiber (NDF) and acid detergent fiber (ADF), and lignin (L) was computed using the methods provided by Babayemi and Bamikole (2006). The Bahauddin Zakariya University Multan's Institute of Food Science and Nutrition conducted all the aforementioned chemical analyses. All these variables usually determine the nutritional quality of a forage species. A high nutritive forage species has a high percentage of nitrogen, ADF, and NDF, which improves its digestibility as well, however, CP, L and HC determine the enhanced quality of forage (Codron *et al.*, 2007).

Statistical analysis

The data on individual preference were analyzed through a Univariate test (two-way ANOVA) and nutritional parameters and fiber fraction was subjected to analysis of variance (ANOVA) in a completely randomized design, followed by the Latin-Square Design (LSD) method to test the difference (Steel *et al.*, 1997) among means of all forage species using Statistical Analysis System Computer Package (SAS, 2014). All the analyses were considered significant at a 95% confidence interval ($P \leq 0.05$).

RESULTS

Trees: Forage preference

The most preferred tree species overall, based on the time taken to visit for foraging, was *Salvadora oleoides* (21.5 ± 2.5 min), followed by *Prosopis cineraria* (21.1 ± 2.3 min; Fig. 2) seems weekly correlated with the higher concentration of DM and CF. Only, *Tamarix aphylla* had significantly lower ($P \leq 0.05$) palatability when compared to all the other plant species ($P \leq 0.05$; $n = 180$) inversely correlated with the highest value of ash content. The time taken to visit each plant species' results were similar to the number of visits to each species' results. The results showed that *Prosopis cineraria* was the most preferred species with an average of 41.0 ± 5.4 animal visits per pile, followed by *Vachellia nilotica* with 39.3 ± 5.6 animal visits, and *Salvadora oleoides* with 37.2 ± 5.1 animal visits per pile again correlated with the higher concentration of DM and CF.

However, these differences were not significant ($P > 0.05$). *Vachellia nilotica* was significantly different from *Zizyphus mauritiana* with 34.86 ± 5.0 animal visits per pile but not significant from *Prosopis cineraria* 41.0 ± 5.4 ($P > 0.05$). *Tamarix aphylla* was the least preferred species with an average of 22.7 ± 4.4 animal visits per pile, which significantly varies ($P \leq 0.05$) from all other tree species (Fig. 2).




Fig. 2. Preference of tree species for forage by blackbuck based on the average number of animal visits and average time spent per visit of each tree species. Where similar alphabets show the non-significance of the treatment and different alphabets show a significant difference between treatments at 95% Confidence intervals.

In terms of gender-based preference by the time taken to visit for forage, male blackbuck significantly preferred *Vachellia nilotica* (25.6 ± 2.6 min) and *Prosopis cineraria* (20.4 ± 1.9 min) which are enriched with DM, NDF, and ADF over the other three tree species ($P \leq 0.05$; Fig. 3).

Female blackbuck significantly preferred *Zizyphus mauritiana*, (25.2 ± 2.8 min), *Prosopis cineraria*, and *Salvadora oleoides* over *Tamarix aphylla* and *Vachellia nilotica* ($P \leq 0.05$; Fig. 2) but surprisingly only *Prosopis cineraria* had a higher concentration of CF, NDF, and ADF while *Salvadora oleoides* was enriched with CP.

Fawns significantly preferred *Salvadora oleoides* (24.2 ± 3 min) and *Prosopis cineraria* over *Tamarix aphylla* and *Zizyphus mauritiana* ($P \leq 0.05$; Fig. 2) but did not prefer the forage species on their nutritious value. Again, the time taken to visit each plant species' result was similar to the number of visits to each species' results. Gender based analysis on visits per pile demonstrated that male blackbuck visited *Vachellia nilotica* the most with

49.2±6.1 visits per pile, while female blackbuck preferred *Zizyphus mauritiana* with 48.4±5.5 visits per pile, and fawns showed the highest preference for *Salvadora oleoides* with 48.4±5.8 visits per pile, correlated with a high concentration of CP. These preferences were significantly different from *Tamarix aphylla* ($P \leq 0.05$; Fig. 2).

Tree: Nutritional analysis

Different tree species perform differently in terms of each nutrient but overall *Prosopis cineraria* and *Vachellia nilotica* were the best in terms of nutrients. DM, CP, ash content, EE, NDF, ADF, HC and L levels varied amongst tree species. The DM content differed substantially ($P \leq 0.05$) among tree species, ranging from 92.58% to 94.50%. *Prosopis cineraria* had the greatest DM value while *Zizyphus mauritiana* had the lowest (Fig. 3).




Fig. 3. Different nutrients in trees preferred by blackbuck for forage, CP, crude protein; CF, crude fiber; EE, ether extract; NDF, neutral detergent fiber; ADF, acid detergent fiber. Where similar Alphabets show the non-significance of the treatment and different alphabets show a significant difference between treatments at 95% Confidence intervals.

The concentration of CP varied from 8.08 to 10.56%, with *Vachellia nilotica* having the highest ($P \leq 0.05$) and *Prosopis cineraria* having the lowest. *Tamarix aphylla* had the greatest crude ash value ($P \leq 0.05$), whereas *Prosopis*

cineraria had the lowest (9.52). Similarly, *Prosopis cineraria* had the highest CF content and *Tamarix aphylla* had the lowest ($P \leq 0.05$), with fiber concentrations varying from 30.44 to 18.39%. EE varied substantially ($P \leq 0.05$) from 1.67 to 3.55%, with *Vachellia nilotica* having the highest and *Tamarix aphylla* having the lowest percentage (Fig. 3). The NDF varied ($P \leq 0.05$) from 41.51 to 47.18% with the highest percentage in *Vachellia nilotica* and the lowest in *Tamarix aphylla*. The ADF ranged from 26.03 to 30.56% whereas *Prosopis cineraria* was found to have a maximum ($P \leq 0.05$) value and *Zizyphus mauritiana* had the lowest value whereas the concentration of HC significantly varied ($P \leq 0.05$) from 15.27 to 18.43% (Fig. 3).

The highest content of HC was observed in *Vachellia nilotica* (18.43%) and lowest in *Tamarix aphylla* (15.27%). Similarly, the highest L concentration was 10.63% in *Vachellia nilotica* and the lowest in *Tamarix aphylla* (7.28%; Fig. 3).

Shrubs: Forage preference

The study indicated that *Haloxylon salicornicum* was the most preferred shrub species overall, with the optimum time to visit forage of 20.8 ± 5.5 min correlated with a higher concentration of HC and NDF. However, *Capparis decidua* (17.9 ± 6.6 min), with a higher concentration of NDF variation ($P \leq 0.05$) from *Haloxylon salicornicum*, but not from *Suaeda fruticosa* and *Haloxylon recurvum*. Conversely, *Calotropis procera* was significantly ($P \leq 0.05$) the least preferred species, with a mean time taken of 13.5 ± 1.8 min. Surprisingly, a higher concentration of CF was determined in this species. *Haloxylon salicornicum* was mostly preferred by blackbuck in terms of the number of visits, with a mean of 41.5 ± 6.8 animals per pile regardless of its nutritional value. This was followed by *Suaeda fruticosa*, with an average of 38.7 ± 17.8 animals per pile, and *Haloxylon recurvum* with an average of 37.9 ± 0.8 animals per pile correlated with higher concentration of DM and CP. The differences between these three species were not statistically significant. However, *Capparis decidua* had a significantly lower number of visits compared to *Haloxylon salicornicum*, with an average of 34.9 ± 12.5 animals per pile, although it was not significantly different from *Suaeda fruticosa* and *Haloxylon recurvum*. In contrast, *Calotropis procera* had the lowest number of visits, with an average of only 24.3 ± 4.3 animals per pile, and this difference was statistically significant (Fig. 4).

Furthermore, the study also examined gender-based preferences, which revealed that male blackbuck preferred *Haloxylon salicornicum* with an average of 49.2 ± 6.1 animals per pile again in correlation with the higher

concentration of HC and NDF. Female blackbuck, on the other hand, showed a higher preference for *Capparis decidua*, with an average of 48.4 ± 5.5 animals per pile, and this species had a higher concentration of NDF. Finally, fawns showed a strong preference for *Suaeda fruticosa*, with an average of 52.2 ± 4.0 fawns per pile with a correlation to higher values of DM and CP. The detailed chart of feeding preference of shrub species by blackbuck is shown in Figure 4.




Fig. 4. Shrub species preferences for forage by blackbuck based on the average number of animal visits and average time spent per visit of each shrub species. Where similar Alphabets show the non-significance of the treatment and different alphabets show a significant difference between treatments at 95% Confidence intervals.

Shrubs: Nutritional analysis

The DM percentage of shrubs varied considerably ($P \leq 0.05$) among species, ranging from 91.56% to 95.24%. *Suaeda fruticosa* had the greatest DM value while *Calotropis procera* had the lowest (Fig. 5).

The CP concentration ranged from 8.69 to 16.90%, with *Suaeda fruticosa* having the greatest ($P \leq 0.05$) CP content and *Haloxylon salicornicum* having the lowest. The greatest value ($P \leq 0.05$) of crude ash was found in *Suaeda fruticosa* (18.72%) and the lowest in *Capparis decidua* (9.17%). Similarly, *Calotropis procera* had the highest CF

content and *Haloxylon recurvum* had the lowest, with fiber concentration ranging from 13.56 to 33.93% ($P \leq 0.05$). EE varied considerably ($P \leq 0.05$) from 1.07 to 3.10%, with *Suaeda fruticosa* having the highest and *Haloxylon salicornicum* having the lowest. The content of NDFs ranged ($P \leq 0.05$) from 30.65 to 42.84%, with *Capparis decidua* having the greatest proportion and *Haloxylon recurvum* having the lowest. The ADF varied from 11.60 to 23.53%, with *Capparis decidua* having the highest ($P \leq 0.05$) value and *Haloxylon recurvum* having the lowest. The HC, on the other hand, varied considerably ($P \leq 0.05$) from 12.65 to 21.05%. The highest contents of HC were observed in *Haloxylon salicornicum* (21.05%) and lowest in *Suaeda fruticosa* (12.65%). Similarly, the highest concentration of L was 6.91% in *Capparis decidua* and lowest in *Calotropis procera* (5.15%) as shown in Figure 5.




Fig. 5. Different nutrients in shrub species preferred by blackbuck for forage. CP, crude protein; CF, crude fiber; EE, ether extract; NDF, neutral detergent fiber; ADF, acid detergent fiber. Where similar Alphabets show the non-significance of the treatment and different alphabets show a significant difference between treatments at 95% confidence intervals.

Grasses: Forage preference

Cenchrus pennisetiformis was the most preferred grass species for foraging by blackbuck correlated with the

highest value of DM and HC, with an average time taken for a visit of 22.5 ± 2.4 min. This was followed by *Lasiurus scindicus* (22.1 ± 2.8 min) and *Panicum antidotale* (21.3 ± 4.6 min) in correlation with higher values of NDF, CF, and ADF, with no statistically significant differences observed between these three species. However, *Aristida depressa* (16.4 ± 5.0 min) varied ($P \leq 0.05$) from all other species, while *Cymbopogon jwarancusa* was significantly ($P \leq 0.05$) the least preferred species and higher amount of NDF and ADF, with an average time taken for a visit of 13.4 ± 0.9 min. Regarding gender-based preferences, male blackbuck showed a significant preference for *Lasiurus scindicus* which has the highest amount of NDF and CF, with an average time taken for the visit of 25.2 ± 2.8 min. On the other hand, female blackbuck preferred *Panicum antidotale* (25.2 ± 3.0 min) correlated with a higher amount of ADF, CF, and DM while fawn preferred *Cenchrus pennisetiformis* (25.2 ± 2.4 min) the most in correlation with the highest concentration of DM, HC, and Ash. The detailed chart represents the preference of grass species (Fig. 6).




Fig. 6. Grass species preferences for forage by blackbuck based on the average number of animal visits and average time spent per visit of each grass species. Where similar Alphabets show the non-significance of the treatment and different alphabets show a significant difference between treatments at 95% confidence intervals.

The study revealed that *Lasiurus scindicus* was the most preferred grass species, with a mean number of visits per pile of 45.5 ± 4.6 animals, followed closely by *Cenchrus pennisetiformis*, with an average of 42.6 ± 5.2 animals per pile. *Panicum antidotale* also had a high preference among blackbuck, with an average of 40.9 ± 5.6 animals per pile. In contrast, *Aristida depressa* had a significantly lower preference compared to all other grass species, with an average of 30.5 ± 8.3 animals per pile. Additionally, *Cymbopogon jwarancusa* had the lowest ($P \leq 0.05$) preference, with an average of only 25.4 ± 4.4 animals per pile. Furthermore, gender-based preference was observed, with male blackbuck showing a significant preference for *Lasiurus scindicus*, with an average of 50.2 ± 6.1 animals per pile. Female blackbuck, on the other hand, preferred *Panicum antidotale* the most, with an average of 47.4 ± 5.5 animals per pile. Fawn showed a significantly stronger preference for *Cenchrus pennisetiformis*, with an average of 48.4 ± 5.8 fawns per pile.

Grasses: Nutritional analysis

The DM percentage of grasses differed substantially ($P \leq 0.05$) across selected grass species, ranging from 94.28% to 96.5%. *Cenchrus pennisetiformis* had the greatest DM value and *Lasiurus scindicus* had the lowest (Fig. 7).

The concentration of CP ranged from 3.9 to 5.8% and the highest ($P \leq 0.05$) content of CP was observed in *Panicum antidotale* and lowest in *Lasiurus scindicus*. The highest value ($P \leq 0.05$) of crude ash was observed in *Cenchrus pennisetiformis* (18.62%) and the lowest in *Panicum antidotale* (8.54%) which ranged in between these values among selected grasses. Similarly, maximum content of CF was present in *Lasiurus scindicus* and lowest in *Cenchrus pennisetiformis* and the concentration of fiber was significantly ($p \leq 0.05$) varied from 32.56 to 45.54%. EE significantly ($P \leq 0.05$) varied from 1.28 to 2.62% with the maximum in *Cymbopogon jwarancusa* and minimum in *Lasiurus scindicus*. The concentration of NDFs varied ($P \leq 0.05$) from 58.04 to 69.0% with the highest percentage in *Lasiurus scindicus* and the lowest in *Panicum antidotale*.

The ADF ranged from 34.8 to 41.53% where *Cymbopogon jwarancusa* was found to have a maximum ($P \leq 0.05$) value and *Aristida depressa* the lowest value whereas the concentration of HC significantly varied ($P \leq 0.05$) from 23.91 to 34.50%. The highest contents of HC were observed in *Cenchrus pennisetiformis* (34.50%) and lowest in *Aristida depressa* (23.91%). Similarly, the highest concentration of L was 5.59% in *Lasiurus scindicus* and the lowest in *Cymbopogon jwarancusa* (4.5%; Fig. 7).




Fig. 7. Different nutrients in grass species preferred by blackbuck for forage. CP, crude protein; CF, crude fiber; EE, ether extract; NDF, neutral detergent fiber; ADF, acid detergent fiber. Where similar Alphabets show the non-significance of the treatment and different alphabets show a significant difference between treatments at 95% confidence intervals.

DISCUSSION

Trees

Salvadora oleoides and *Prosopis cineraria* were the most favored tree species for blackbuck in the Cholistan rangeland. Previous research has identified *Salvadora oleoides* and *Prosopis cineraria* as preferred food species for other herbivorous animals such as camels and sheep in arid and semiarid environments (Moyo *et al.*, 2011; Gul *et al.*, 2014). A high nutritive forage species has a high percentage of nitrogen, ADF, and NDF, which improves its digestibility as well. However, CP, L, and HC determine the enhanced quality of forage (Codron *et al.*, 2007).

In contrast, *Tamarix aphylla* was the least preferred species for blackbuck, as indicated by both the time taken to visit forage and the number of animal visits. These results are consistent with previous studies showing *Tamarix aphylla* as an unpalatable species by other herbivorous animals i.e., goats and sheep (El-Shaer, 2004).

However, this is contradictory to another study suggesting that *Tamarix aphylla* was highly palatable for camels and overall it was categorized as moderately palatable (Khan and Hussain, 2012). This could be due to the presence of secondary compounds such as tannins and terpenoids that could reduce their palatability and digestibility (McDonald *et al.*, 2011).

Gender-based differences in forage preference were also observed, with male blackbuck showing high preference for *Vachellia nilotica* and female blackbuck showing a higher preference for *Zizyphus mauritiana*. Similar results were observed in various studies that depicted the gender-based differences in forage preference among herbivorous animals such as sheep and deer (Abdullah *et al.*, 2017; Seri *et al.*, 2018; Jesmer *et al.*, 2020). These variations could be due to variation in dietary needs, social behavior, and physiological variation among different gender groups (Arganini *et al.*, 2012).

Our results also revealed considerable differences in the nutritional potential of some tree species. *Prosopis cineraria* had the highest DM content, while *Vachellia nilotica* had the highest concentration of CP. These findings are consistent with previous studies that reported *Prosopis cineraria* and *Vachellia nilotica* as having high nutritional value for herbivorous animals in arid and semi-arid regions (Bhatta *et al.*, 2005; Abdullah *et al.*, 2017). Also, *Prosopis cineraria* and *Vachellia nilotica* are the most palatable species for livestock in rangelands (Fagg and Stewart, 1994). However, the nutritive potential of forage is not the sole determinant of forage preference, as other factors such as availability, accessibility, and palatability could also influence forage selection by all herbivores.

Shrubs

Haloxylon salicornicum was the most preferred shrub species overall by blackbuck in our study in line with previous studies on blackbuck that reported their preference for *Haloxylon salicornicum* (Ashraf *et al.*, 2013). In terms of gender-based preference, male blackbuck showed a significant preference for *Haloxylon salicornicum*, while female blackbuck preferred *Capparis decidua* and fawn preferred *Suaeda fruticosa* among the shrub species. These results are consistent with previous studies that reported gender-based preference in blackbuck (Mirza and Waiz, 1973).

The nutritional analysis of shrub species showed significant variations in DM, CP, crude ash, CF, EE, NDFs, ADF, HC, and L contents among the selected shrubs. The highest value of DM, CP, and crude ash was determined in *Suaeda fruticosa* and the lowest in *Calotropis procera*, *Haloxylon salicornicum*, and

Capparis decidua respectively consistent with previous studies that have examined the nutritional value of various shrub species for blackbuck. For instance, a study reported that the CP content of *Haloxylon salicornicum* ranged from 5.06 to 15.44% (Ashraf *et al.*, 2013) which was lower than the value obtained in this study. Similarly, the CP content of *Suaeda fruticosa* ranged from 10 to 13.5% (Joshi *et al.*, 2018), which is consistent with the value obtained in this study. EE is a lipid component, and animals primarily rely on it for energy generation and maintenance. The presence of high amounts of EE in meal samples indicates that the animals' energy levels are greater (Odedire and Babayemi, 2008). According to our results, the maximum value of EE was observed as 03.55% (*Vachellia nilotica*) and a minimum of 01.07% (*Haloxylon salicornicum*). Our findings were almost comparable with the work of Mahala *et al.* (2009) and Abdullah *et al.* (2017). The results indicated higher EE levels than those found by Towhidi and Zhandi (2007) but slightly lower than those reported by Njidda and Ikhimioya (2010). Elevated EE can enhance feed palatability and provide essential fatty acids, aiding in energy density and vitamin absorption, which is beneficial for blackbuck's growth and health. However, excessive EE may cause digestive issues, nutrient imbalance, and obesity (Katoch, 2022). That's why there is much need for careful dietary balance to optimize the nutritional benefits without adverse effects on the health of blackbuck.

The results of the study suggest that there is a significant correlation between the food preference of blackbuck and the nutritional contents of shrubs. Blackbuck prefers *Haloxylon salicornicum* and *Suaeda fruticosa* because of their high value of HC, DM, and CP, which indicates that they have strong nutritional potential. Whereas, *Calotropis procera*, was the least preferred shrub species, and had the lowest values of DM, CP, and EE. This suggests that the low nutritional value of *Calotropis procera* may be a factor in being unpalatable for blackbuck. These results are in line with earlier studies on herbivorous mammals, which show a preference for plants with more nutrients that are easier to digest (Fritz *et al.*, 2002). Furthermore, a correlation can be observed between those shrubs' palatability and nutritional content. *Haloxylon salicornicum*, having the highest number of visits, also had the highest concentration of HC among the selected shrubs in line with previous research (Ashraf *et al.*, 2013). Similarly, *Suaeda fruticosa*, which had the second-highest number of visits, had the highest values of DM and CP.

Grasses

Our results for grass species preference by blackbuck are in line with earlier research on other herbivorous animals. For instance, preferred grass species like *Cenchrus*

ciliaris and *Pennisetum mezianum* had high protein content (Otieno and Kinyamario, 2018). According to the current study, *Lasiurus scindicus* and *Panicum antidotale* were the most palatable grass species for blackbuck in line with the previous nutritional study of the grasses, which showed that *Panicum antidotale* had the highest CP content and *Cenchrus pennisetiformis* had the highest DM content (Fritz *et al.*, 2002).

There is a correlation between species preference and nutrient content, the grasses selected by herbivorous animals were enriched with CF which is an important nutrient and is helpful for gastrointestinal function (Mena and Bresciani, 2020). Additionally, because they encourage growth and reproduction, high protein grasses were chosen by herbivorous animals (Van Dyne *et al.*, 2019).

CONCLUSION

It can be concluded that blackbuck has distinct preference for different tree, shrub and grass species. The study found that *Salvadora oleoides* and *Prosopis cineraria* were the most preferred tree species, while *Tamarix aphylla* was the least preferred species. Male blackbuck showed a higher preference for *Vachellia nilotica*, while female blackbuck preferred *Zizyphus mauritiana*. The study also found that *Haloxylon salicornicum* was the most preferred shrub species overall, with male blackbuck showing a significant preference for it, while female blackbuck preferred *Capparis decidua* and fawn preferred *Suaeda fruticosa*. Furthermore, male blackbuck prefers *Panicum antidotale* the most, female preferred *Panicum antidotale* and fawn preferred *Cenchrus pennisetiformis*. It is important to consider these preferences when managing the habitats of blackbuck. Preserving the vegetation that blackbuck prefers, may support the plans to reintroduce them in the wild more securely. The nutritional analysis of the selected species showed significant variations in DM, CP, CA, CF, EE, NDF, ADF, HC, and L content among the selected shrubs. The study suggests that palatability could be influenced by different nutrients in different gender groups of blackbuck. Because of the nutritional needs of different gender groups, they may have different requirements.

DECLARATIONS

Acknowledgements

We express our sincere gratitude to the teachers at Bhauddin Zakariya University Multan's Department of Forestry and Range for their kind cooperation and help during the data collection phase. We would also like to extend our sincere gratitude to the administrators and

workers at Lal Suhanra National Park, especially Mr. Muhammad Zahoor (SDFO (Retd.).

Funding

This research project was entirely self-funded by the authors.

Data and materials availability

All the data and material can be provided on request at any of the author's email.

Generative AI and AI-assisted technology statement

The authors have declared that no generative AI or AI-assisted technologies were used to create this manuscript.

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Abdullah, M., Rafay, M., Hussain, T., Ahmad, H., Tahir, U., Rasheed, F., Ruby, T. and Khalil, S., 2017. Nutritive potential and palatability preference of browse foliage by livestock in arid rangelands of Cholistan desert (Pakistan). *J. Anim. Pl. Sci.*, **27**: 1656–1664.
- Abid, S. and Tasleem, M.W., 2022. Population status, distribution, threats and conservation of blackbuck antelope cervicapra in South Asia: An updated. *Int. Peer Rev. Open Access J.*, **12**. <https://doi.org/10.21786/bbrc/15.1.3>
- Aleem, A., 1978. Re-introduction of blackbuck in Pakistan. *Pakistan J. For.*, **28**: 111–115.
- AOAC, 2005. *Official methods of analysis*. Association of Official Analytical Chemists International Gaithersburg (MD).
- Arganini, C., Saba, A., Comitato, R., Virgili, F. and Turrini, A., 2012. Gender differences in food choice and dietary intake in modern western societies. *Publ. Hlth. Soc. Behav. Hlth.*, **4**: 83–102. <https://doi.org/10.5772/37886>
- Ashraf, M.A., Mahmood, K., Wajid, A., Qureshi, A.K. and Gharibreza, M., 2013. Chemical constituents of *Haloxylon salicornicum* plant from Cholistan desert, Bahawalpur, Pakistan. *J. Fd. Agric. Environ.*, **11**: 1176–1182.
- Babayemi, J. and Bamikole, M., 2006. Supplementary value of *Tephrosia bracteolata*, *Tephrosia candida*, *Leucaena leucocephala* and *Gliricidia sepium* Hay for West African dwarf goats kept on range. *J. Cent. Eur. Agric.*, **7**: 323–328.
- Balluffi-Fry, J., Leroux, S.J., Wiersma, Y.F., Richmond, I.C., Heckford, T.R., Rizzuto, M., Kennah, J.L. and Vander, W.E., 2022. Integrating plant stoichiometry and feeding experiments: State-dependent forage choice and its implications on body mass. *Oecologia*, **198**: 579–591. <https://doi.org/10.1007/s00442-021-05069-5>
- Bhatta, R., Vaithyanathan, S., Singh, N.P., Shinde, A.K. and Verma, D.L., 2005. Effect of feeding tree leaves as supplements on the nutrient digestion and rumen fermentation pattern in sheep grazing on semi-arid range of India-I. *Small Rumin. Res.*, **60**: 273–280. <https://doi.org/10.1016/j.smallrumres.2005.01.009>
- Chambers, R., 1996. 4 World wildlife fund and political localism. *Environ. Activism World Civic Polit.*, **4**: 72
- Choudhary, N.L. and Chisty, N., 2022. Behavioural biology and ecology of blackbuck (*Antelope cervicapra*): A review. *Flora Fauna*, **28**: 355–361. <https://doi.org/10.33451/florafauna.v28i2pp355-361>
- Codron, D., Lee-Thorp, J.A., Sponheimer, M., Codron, J., De Ruiter, D. and Brink, J.S., 2007. Significance of diet type and diet quality for ecological diversity of African ungulates. *J. Anim. Ecol.*, **76**: 526–537. <https://doi.org/10.1111/j.1365-2656.2007.01222.x>
- Csurhes, S. and Fisher, P., 2010. *Blackbuck, Antelope cervicapra pest animal risk assessment*. Biosecurity Queensland. Department of Employment, Economic Development and Innovation. pp. 11.
- Davies, N.B., Krebs, J.R. and West, S.A., 2012. *An introduction to behavioural ecology*. John Wiley and Sons.
- Dostaler, S., Ouellet, J.P., Therrien, J.F. and Côté, S.D., 2011. Are feeding preferences of white-tailed deer related to plant constituents? *J. Wildl. Manage.*, **75**: 913–918. <https://doi.org/10.1002/jwmg.118>
- El-Shaer, H.M., 2004. *Potentiality of halophytes as animals fodders under arid conditions of Egypt*. Cahiers Options Méditerranéennes (CIHEAM), pp. 369–374.
- Fagg, C.W. and Stewart, J.L., 1994. The value of *Acacia* and *Prosopis* in arid and semi-arid environments. *J. Arid Environ.*, **27**: 3–25. <https://doi.org/10.1006/jare.1994.1041>
- Farooq, Z., Iqbal, M., Khan, M.S., Asad, S. and Wajid, M., 2022. Circadian variation in activities of blackbuck under captivity in Punjab, Pakistan. *Proc. Zool. Soc.*, **75**: 242–246. <https://doi.org/10.1007/s12595-022-00438-1>
- Felton, A.M., Holmström, E., Malmsten, J., Felton, A., Croomsigt, J.P.G.M., Edenius, L., Ericsson, G., Widemo, F. and Wam, H.K., 2020. Varied diets, including broadleaved forage, are important for a

- large herbivore species inhabiting highly modified landscapes. *Sci. Rep.*, **10**: 1–13. <https://doi.org/10.1038/s41598-020-58673-5>
- Fritz, H., Duncan, P., Gordon, I.J. and Illius, A.W., 2002. Megaherbivores influence trophic guilds structure in African ungulate communities. *Oecologia*, **131**: 620–625. <https://doi.org/10.1007/s00442-002-0919-3>
- Gul, B., Ansari, R., Ali, H., Adnan, M.Y., Weber, D.J., Nielsen, B.L., Koyro, H.W. and Khan, M.A., 2014. The sustainable utilization of saline resources for livestock feed production in arid and semiarid regions: A model from Pakistan. *Emirat. J. Fd. Agric.*, **26**: 1032–1045. <https://doi.org/10.9755/ejfa.v26i12.19102>
- IUCN, S.S.C., 2017. Antelope specialist group (2017). *Antilope cervicapra*. The IUCN red list of threatened species, **2017**: e. T1681A50181949.
- Jesmer, B.R., Kauffman, M.J., Murphy, M.A. and Goheen, J.R., 2020. A test of the niche variation hypothesis in a ruminant herbivore. *J. Anim. Ecol.*, **89**: 2825–2839. <https://doi.org/10.1111/1365-2656.13351>
- Jhala, Y.V. and Isvaran, K., 2016. Behavioural ecology of a grassland antelope, the blackbuck *Antilope cervicapra*: Linking habitat, ecology and behaviour. In: *The ecology of large herbivores in South and Southeast Asia*. Springer. pp. 151–176. https://doi.org/10.1007/978-94-017-7570-0_6
- Joshi, A., Kanthaliya, B. and Arora, J., 2018. Halophytes of thar desert: Potential source of nutrition and feedstuff. *Int. J. Bioassays*, **8**: 5674–5683.
- Katoch, R., 2022. Nutritional and anti-nutritional constituents in forages. In: *Nutritional quality management of forages in the himalayan region*. Springer. pp. 193–223. https://doi.org/10.1007/978-981-16-5437-4_8
- Khalil, S., Safeer, M., Riaz, S., Jamil, H. and Noor, U., 2020. Biology and ecological adaptations of blackbuck (*Antilope cervicapra*): A review. *Agrobiol. Rec.*, **4**: 23–31. <https://doi.org/10.47278/journal.abr/2020.023>
- Khan, B.N., Ehsanullah, T., Bibi, A., Khan, Z.I., Ahmad, M., Bashir, A. and Fahid, A.U.M., 2023. Feeding ecology of reintroduced blackbucks in Lal Suhanra National Park, Bahawalpur. *J. Bioresour. Manage.*, **10**: 17.
- Khan, M. and Hussain, F., 2012. Palatability and animal preferences of plants in Tehsil Takht-e-Nasrati, District Karak, Pakistan. *Afr. J. agric. Res.*, **7**: 5858–5872. <https://doi.org/10.5897/AJAR12.2095>
- Kravetz, F. and Suárez, O., 1998. Transmission of food selectivity from mothers to offspring in *Akodon azarae* (Rodentia, Muridae). *Behaviour*, **135**: 251–259. <https://doi.org/10.1163/156853998793066311>
- Mahala, A.G., Nsahlai, I.V., Basha, N.A.D. and Mohammed, L.A., 2009. Nutritive evaluation of natural pasture at early and late rainfall season in Kordofan and Butana, Sudan. *Austral. J. Basic appl. Sci.*, **3**: 4327–4332.
- McDonald, P., Edwards, R.A., Greenhalgh, J.F.D., Morgan, C.A., Sinclair, L.A. and Wilkinson, R.G., 2011. *Feeding standards for maintenance and growth*. Animal Nutrition, 7th Ed.; Pearson Education Limited: Harlow, UK, pp. 343–383.
- Meena, R. and Saran, R.P., 2018. Distribution, ecology and conservation status of blackbuck (*Antilope cervicapra*): An update. *Int. J. biol. Res.*, **3**: 79–86.
- Mena, P. and Bresciani, L., 2020. Dietary fibre modifies gut microbiota: What's the role of (poly) phenols? *Int. J. Fd. Sci. Nutr. Taylor Francis*, **71**: 783–784. <https://doi.org/10.1080/09637486.2020.1826913>
- Mirza, Z.B. and Waiz, A., 1973. Food availability for blackbuck (*Antilope cervicapra*) at Lal Suhanra Sanctuary, Pakistan. *Biol. Conserv.*, **5**: 119–122. [https://doi.org/10.1016/0006-3207\(73\)90091-8](https://doi.org/10.1016/0006-3207(73)90091-8)
- Moyo, B., Masika, P.J., Hugo, A. and Muchenje, V., 2011. Nutritional characterization of moringa (*Moringa oleifera* Lam.) leaves. *Afr. J. Biotechnol.*, **10**: 12925–12933. <https://doi.org/10.5897/AJB10.1599>
- Njidda, A.A. and Ikhimiya, I., 2010. Nutritional evaluation of some semi-arid browse forages leaves as feed for goats. *Eur. J. appl. Sci.*, **2**: 108–115.
- Odedire, J.A. and Babayemi, O.J., 2008. Comparative studies on the yield and chemical composition of *Panicum maximum* and *Andropogon gayanus* as influenced by *Tephrosia candida* and *Leucaena leucocephala*. *Livest. Res. Rural Dev.*, **20**: 1–8.
- Otieno, D.O. and Kinyamario, J.I., 2018. Grasslands of Eastern Africa: Problems and prospects. In: *Grasslands of the World*. CRC Press. pp. 307–332. <https://doi.org/10.1201/9781315156125-21>
- Prater, S.H. and Barruel, P., 1971. *The book of Indian animals (Bombay Natural History Society, Bombay)*. 3rd ed. Bombay Natural History Society; Oxford University Press.
- SAS Institute Inc., 2014. *The ANOVA Procedure (RCBD)*. In SAS/STAT® 13.1 User's Guide. Cary, NC: SAS Institute Inc.
- Seri, H., Chammem, M., Ferreira, L.M.M., Kechnebou, M., Khorchani, T. and Silva, S.R., 2018. Effects of seasonal variation, group size and sex on the activity budget and diet composition of the *Addax*

- antelope. *Afr. J. Range Forage Sci.*, **35**: 89–100. <https://doi.org/10.2989/10220119.2018.1477831>
- Steel, R.G.D., Torrie, J.H. and Dickey, D., 1997. *Principles and procedure of statistics. A biometrical approach* 3rd Ed. McGraw HillBookCo. Inc., New York, pp. 352–358.
- Stutz, R.S., Bergvall, U.A., Leimar, O., Tuomi, J. and Rautio, P., 2018. Cohesiveness reduces foraging efficiency in a social herbivore. *Anim. Behav.*, **135**: 57–68. <https://doi.org/10.1016/j.anbehav.2017.11.004>
- Tahir, R., 2022. Population status, distribution, threats and conservation of blackbuck *Antilope cervicapra* in South Asia: An updated literature review. *Biosci. Biotechnol. Res. Commun.*, **15**: 12–25. <https://doi.org/10.21786/bbrc/15.1.3>
- Towhidi, A. and Zhandi, M., 2007. Chemical composition, *in vitro* digestibility and palatability of nine plant species for dromedary camels in the province of Semnan, Iran. *Egypt. J. Biol.*, **9**. <https://doi.org/10.1017/S1752756200021360>
- Van Dyne, G.M., Hanson, J.D. and Jump, R.C., 2019. *Seasonal changes in botanical and chemical composition and digestibility of diets of large herbivores on shortgrass prairie*. Proceedings Of The Xiv International Grassland Congress, pp. 684–687.
- Vats, R. and Bhardwaj, C.S., 2009. Investigation in activity time budget and social organization of indian black buck (*Antilope cervicapra*) Linn. (*Mammalia artodactyla*) at MC Zoological Park, Chhatbir (Pb.)(India). *Curr. World Environ.*, **4**: 97. <https://doi.org/10.12944/CWE.4.1.13>
- Wakchaure, R. and Ganguly, S., 2016. Captive breeding in endangered wildlife: A review. *J. biol. Sci. Opin.*, **4**: 186–187. <https://doi.org/10.7897/2321-6328.04544>
- Wariss, H.M., Pirzada, S.A., Alam, K., Anjum, S. and Qureshi, R., 2014. Flora of Lal Suhanra National Park, Bahawalpur, Punjab, Pakistan. *Pakistan J. Bot.*, **46**: 1331–1341.
- Wright, F.C., 1985. *Tricholipeurus balanicus balanicus* (Werneck, 1938) (Malbophaga, Trichodectidae) on blackbuck antelope (*Antilope cervicapra* L.) in Texas. *J. Wildl. Dis.*, **21**: 168169. <https://doi.org/10.7589/0090-3558-21.1.68>