

Evaluation of Reference Gene Expression on Different Tissue in Adults of Bay Snook *Petenia splendida*

Carlos Alfonso Alvarez-González^{1*}, Alejandra del Carmen Castillo-Collado², Gloria Asencio-Alcudia¹, Carlos Alfonso Frías-Quintana³, Vicente Morales-García⁴, Carina Shianya Alvarez-Villagomez¹, Bartolo Concha-Frías⁵, Rafael Martínez-García¹ and Luis Daniel Jimenez-Martinez^{2*}

¹Laboratorio de Acuicultura Tropical, DACBIOL-UJAT, CP. 86150, Villahermosa, México

²Laboratorio de Biología Molecular, DAMJM-UJAT, C.P. 86205, Jalpa de Méndez, Tabasco, México

³Laboratorio de Investigación en Biotecnología Acuicola (LIBA), Tecnológico Nacional de México. Campus Boca del Río (ITBoca), Boca del Río, Veracruz, México.

⁴Laboratorio de Docencia DAMC-UJAT, Ranchería Sur Cuarta Sección, Comalcalco, México

⁵Universidad Popular de la Chontalpa (UPCH). Carretera Cárdenas-Huimanguillo Km 2 S/N, Ranchería, Invitab Paso y Playa, 86556, Cárdenas, Tab. Mexico

Luis Daniel Jimenez-Martinez and Carlos Alfonso Álvarez-González contributed equally to this work.

ABSTRACT

The tenguayaca mojarra *Petenia splendida* is a fish that belongs to the cichlid family and its distribution is located from the southeast of Mexico to Guatemala and Belize. Therefore, molecular biology studies, particularly gene expression, provide information on genes that encode the production of specific proteins and RNA molecules for various functions in living organisms and biological processes biológicos. A very important technique to analyze gene expression levels under different conditions is the qRT-PCR polymerase chain reaction. The objective of the study was to analyze the stability of the reference genes 18S ribosomal RNA (*18S rRNA*), Elongation factor 1-alpha (*efl-a*), Alpha actin (*acta*), Beta-actin (*actb*), Glyceraldehyde-3-phosphate dehydrogenase (*gapdh*) and Alpha tubulin (*α-tubulin*) using the BestKeeper, Normfinder and geNorm software in the expression of different tissues tissues of liver, muscle, spleen, intestine, gill, brain and testis in the case of male organisms and ovary in females of the tenguayaca mojarra (*Petenia splendida*). Finally we can conclude that based on BestKeeper, Normfinder and GeNorm the most stable reference genes in most tissues were 18S rRNA and *actb*, while in the case of the brain it was *α-tubulin* and *gapdh* in the ovary and testes.

Article Information

Received 16 April 2024

Revised 05 November 2025

Accepted 30 November 2025

Available online 28 March 2026 (early access)

Authors' Contribution

ADCCC: Data curation, formal analysis, investigation, methodology, supervision, validation, visualization, writing-original draft, writing-review and editing. CAAG: Conceptualization, investigation, methodology, project administration, resources, supervision, validation, visualization, writing-original draft, writing-review and editing. GA-A: Data curation, formal analysis, methodology, supervision, validation, writing-original draft, writing-review and editing. CAFQ: Data curation, formal analysis, methodology, supervision, validation, writing-original draft, writing-review and editing. VM-G: Advice and data analysis, investigation, methodology, validation, writing-original draft. CSA-V: Conceptualization, formal analysis, investigation, supervision, validation, writing-original draft, writing-review and editing. BC-F: Formal analysis, investigation, methodology, supervision, validation, visualization, writing-original draft, writing-review and editing. RM-G: Writing, reviewing and editing draft and final document. LDJ-M: Conceptualization, investigation, methodology, project administration, resources, supervision, validation, visualization, writing-original draft, writing-review and editing.

Key words

Reference gene, Expression, Gen, Tissue, Bay snook, *Petenia splendida*

* Corresponding author: luisd1984@hotmail.com, alvarez_alfonso@hotmail.com
0030-9923/2026/0001-0001 \$ 9.00/0



Copyright 2026 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

INTRODUCTION

The tenguayaca mojarra *Petenia splendida* is a fish that belongs to the cichlid family and its distribution is located from the southeast of Mexico to Guatemala and Belize. Likewise, *P. splendida* is a carnivorous species that is widely accepted in local markets for its meat quality, mainly as a great source of protein (Arias-Rodriguez *et al.*, 2008; Vidal-López *et al.*, 2009).

Currently, there are various studies carried out on

the tenguayaca mojarra *Petenia splendida*, of which the following stand out: biological and physiological aspects (Álvarez-González *et al.*, 2008; Jiménez-Martínez *et al.*, 2009), taxonomy and ecology (Méndez-García, 2010), while the aquaculture area highlights studies of embryonic development, optimal light and temperature conditions, management of breeders, larval reproduction and population density (Álvarez-González *et al.*, 2008; Contreras-García *et al.*, 2018; Jiménez-Martínez *et al.*, 2009; Pérez-Sánchez and Páramo-Delgadillo, 2008; Treviño *et al.*, 2011; Vidal-López *et al.*, 2009) nutritional and digestive physiology (Álvarez-González *et al.*, 2008; Rodríguez-Estrada *et al.*, 2020; Uscanga-Martínez *et al.*, 2011) and cytogenetics (Arias-Rodríguez *et al.*, 2008). However, in the area of molecular biology, there are few studies carried out in this species, of which the obtaining of the complete mitogenome obtained in adult wildlife organisms and the expression of the Glut2 gene in captive larvae and adults stand out (Castillo-Collado *et al.*, 2022; Del Río-Portilla *et al.*, 2016). Therefore, molecular biology studies, particularly gene expression, provide information on genes that encode the production of specific proteins and RNA molecules for various functions in living organisms and biological processes biológicos (Kozera and Rapacz, 2013). A very important technique to analyze gene expression levels under different conditions is the qRT-PCR polymerase chain reaction. However, to analyze the expression of a gene, two methods are used: absolute relative expression and relative expression, the first is based on a standard curve responsible for quantifying an unknown based on a known amount of the gene and the second quantifies the expression through the comparison of a candidate gene against a reference gene or control gene (Chen *et al.*, 2023; Infante *et al.*, 2008; Kozera and Rapacz, 2013; Tang *et al.*, 2007). In this way, reference genes are those that are expressed constantly and independently of the physiological state of the cell (Jiménez *et al.*, 2022; Pfaffl *et al.*, 2004).

Therefore, the first step in analyzing the expression of a specific gene is to select and identify the best reference genes in our studies to obtain conclusive and reliable data in our qPCR (Kumari *et al.*, 2022; Rassier *et al.*, 2020). The main characteristic that a reference gene must have is to present stable expression in the different tissues, stages of development of the organisms, and experimental conditions studied (Li *et al.*, 2010; Zhang *et al.*, 2021).

In this sense, there are bioinformatics tools that help us validate the performance of reference genes through specific algorithms using the ct values obtained from the samples analyzed by real-time PCR (qPCR), such as BestKeeper (Pfaffl, 2001), Normfinder (Andersen *et al.*, 2004) and GeNorm (Vandesompele *et al.*, 2002).

In the case of fish, there are various works that analyze

the stability of different reference genes using these software through quantitative qPCR in different investigations such as nutrition, development, endocrinology and toxicology in both larvae, juveniles and adults in marine species such as: *Gasterosteus aculeatus* (Hibbeler *et al.*, 2008), *Solea senegalensis* and *Hippoglossus hippoglossus* (Infante *et al.*, 2008; Øvergård *et al.*, 2010), *Salmo salar* (Kortner *et al.*, 2011), *Gadus morhua* (Olsvik *et al.*, 2008), *Danio rerio* (Casadei *et al.*, 2011; McCurley and Callard, 2008), *Paralichthys olivaceus* (Zheng and Sun, 2011), *Anguilla anguilla* (Li *et al.*, 2023), *Scatophagus argus* (Liu *et al.*, 2023), *Takifugu bimaculatus* (Zhong *et al.*, 2022) and in freshwater fish such as *Siniperca chuatsi* (Zhou *et al.*, 2010), *Cyprinus carpio* (Tang *et al.*, 2012), *Oreochromis niloticus* (Yang *et al.*, 2013), *Ctenopharyngodon idella* (Su *et al.*, 2011; Ye *et al.*, 2010) and *Esox lucius* (Jothinarayanan *et al.*, 2023), *Labeo rohita* (Sahoo *et al.*, 2023), *Channa striata* (Kuah *et al.*, 2016; Purohit *et al.*, 2016), *Hucho taimen* (Yang *et al.*, 2022), *monopterus albus* (Hu *et al.*, 2014; Liu *et al.*, 2022), *Atractosteus tropicus* (Jiménez-Martínez *et al.*, 2022). The objective of the study was to analyze the stability of the reference genes 18S ribosomal RNA (*18S rRNA*), elongation factor 1-alpha (*efl-α*), alpha actin (*acta*), beta-actin (*actb*), glyceraldehyde-3-phosphate dehydrogenase (*gapdh*) and alpha tubulin (*α-tubulin*) using the BestKeeper, Normfinder and geNorm software in the expression of different tissues in males and females of the tenguayaca mojarra (*Petenia splendida*). For this reason, it is important to select the best reference gene to normalize the data obtained from qPCR and calculate the relative expression of the problem gene and also to carry out future gene expression studies involved in the aspects of physiology, nutrition, reproduction in the species *P. splendida*.

MATERIALS AND METHODS

To carry out this study, a total of 20 adult specimens were obtained, 10 males (0.40 - 0.52 kg and 32 to 34 cm total length) and 10 females of *P. splendida* (0.40 - 0.52 kg and 32 to 34 cm total length), of which 10 individuals (0.40 - 0.52 kg and 32 to 34 cm total length) which were captured in the El-Horizonte lagoon of the community of Espino, which is located 32 km from the city of Villahermosa, Tabasco, Mexico, with geographical coordinates 18° 14' 50" north latitude and 92° 49' 58" west latitude.

Sampling, RNA extraction, and cDNA synthesis.

The adult individuals of *Petenia splendida* were sacrificed by the heat shock method (-4 °C) following the steps of the methodology of Matthews and Vargas (2012), subsequently, they were dissected to obtain the tissues of

liver, muscle, spleen, intestine, gill, brain and testis in the case of male organisms and ovary in females. Likewise, total RNA was extracted from a pool of tissues per replicate using Trizol reagent (Invitrogen, Carlsbad, CA). To obtain purity, concentration, and integrity of the RNA, they were determined by spectrophotometry using the 260/280 ratio using a NanoDrop 2000 (Thermo Scientific, USA). Subsequently, 1 µg of RNA was used for reverse transcription with iScript™ Select cDNA Synthesis Kit 170–8,896 (Biorad, Hercules, CA).

Primer design

Specific primers were designed for the genes 18S ribosomal RNA (*18S rRNA*), elongation factor 1-alpha (*efl-α*), beta-actin (*actb*), glyceraldehyde-3-phosphate dehydrogenase (*gapdh*) and alpha tubulin (*α-tubulin*) from complete sequences of *Oreochromis niloticus* while alpha actin (*acta*) was designed from the sequence of *Oreochromis aureus* on the GenBank platform using Primer 3 Plus software version 4.0.0 (https://primer3plus.com/primer3web/primer3web_input.htm). The oligonucleotides were analyzed using Oligo Calculator software version 3.27 (<http://biotools.nubic.northwestern.edu/OligoCalc.html>) to corroborate that secondary structures had not formed.

Quantitative polymerase chain reaction (qPCR)

The cDNA resulting from tissues obtained from the adults of both male and female *P. splendida* was diluted in 200 µl of distilled water. Quantitative polymerase chain reactions (qPCR) were performed in a CFX96 Real-Time System Thermal Cycle (model C1000, CA) 96-well thermal cycler. The reaction mixture included 10 µl of Eva Green, 2 µl of cDNA, and 0.2 µM of each primer. Subsequently, the qPCR runs were performed under the conditions of 2 min at 95°C, followed by 38 cycles at 95°C for 10 s,

60°C for 30 s, and extension at 70°C for 5 s. Similarly, the efficiency (E) and the correlation coefficient (R²) of the primers were evaluated using four serial dilutions in triplicate corresponding to cDNA transcribed from 100 to 0.1 ng of total RNA used as liver tissue. Similarly, it was used a standard curve for each primer applying the formula: $E (\%) = (10^{-1/\text{slope}} - 1) \times 100$.

Stability analysis of candidate reference genes

According to the cq obtained from the qPCR using the Bio-Rad CFX-96 Manager thermocycler, the stability of the six candidate reference genes was analyzed using three software packages, including BestKeeper v1 (<https://www.gene-quantification.de/bestkeeper.html#download>), GeNorm integrated into qBasePlus (<https://www.genequantification.de/hkg.html#genorm>) and Normfinder v20 (<https://www.moma.dk/normfinder-software/>). The results from the three software packages were compared and analyzed to determine which gene is the most appropriate reference gene according to the methodology of Li *et al.* (2019).

RESULTS

Analysis of specificity and reliability of qPCR primer

To evaluate the specificity of qPCR, it was observed that the melting curves presented a single product (Fig. 1). Thus, melting curve analysis is used to evaluate the specificity of the quantitative PCR assay when stains, such as EvaGreen or SYBR®, are used. According to the results obtained from the standard curves according to the C_q values of the eight genes of *Petenia splendida*, showing efficiency values of <98 or ≥100% and the R² values were ≤0.999 (Table I).





Fig. 1. Melting curves of six candidate reference genes of *Atractosteus tropicus* for qPCR. (A) 18S ribosomal RNA (*18S rRNA*), (B) Elongation factor 1-alpha (*efl-α*), (C) Alpha actin (*acta*), (D) Beta-actin (*actb*), (E) Glyceraldehyde-3-phosphate dehydrogenase (*gapdh*) and (F) Alpha tubulin (*α-tubulin*).

Table I. Primer sequences and amplification parameters of six candidate reference genes of *Atractosteus tropicus* used in qPCR analysis.

Gene	Primer (5'–3')	Product size (pb)	Amplification efficiency (%)	R ²	GenBank accession numbers
<i>18S rRNA</i>	F: ACTCGTGATAGGGATTGGGG R: CGATCCGAGGACCTCACTAA	155	98	0.99	XR_003216134
<i>efl-α</i>	F: ACGTCAAGAACGTCTCCGTC R: GAGCTCGCTGAATTGTCAGG	194	96	0.98	EU503119
<i>acta</i>	F: TACGCCAACAATGTGCTCTC R: CTGGAAGGTGGACAGGGAAG	180	95	0.95	EF206799
<i>actb</i>	F: ACAACGTCCCAGTCTATGAGG R: CCTTGATGTACGCACGATC	159	98	0.98	KJ126772
<i>gapdh</i>	F: TGACCGTATGCAGAAGGAGA R: CCTCGTCGTACTCCTGCTTG	164	96	0.98	XM_005455438
<i>α-tubulin</i>	F: CAGATTGGGAATGCCTGCTG R: CCTCATCGATCACAGTGGGT	190	92	0.95	XM_019352023

**Fig. 2. BestKeeper analysis results of six candidate reference genes of *Atractosteus tropicus*.**

BestKeeper gene analysis

According to the results obtained from the six reference genes analyzed in BestKeeper, software that

determines stability using the SD value. In this way, the gene with the lowest SD value is considered the most stable, in the different tissues analyzed in both male and female organisms of *P. splendida* (Fig. 2). Thus, in the case of the liver, the most stable gene was *α-tubulin*, followed by *18S rRNA*, *actb*, *efl-α*, *gapdh*, and *acta* in both male and female organisms. Likewise, in the intestine the most stable gene was *18S rRNA*, followed by *actb*, *gapdh*, *efl-α*, *α-tubulin*, and *acta* in both male and female organisms. Similarly, in muscle, it is observed that the most stable gene was *actb*, followed by *18S rRNA*, *efl-α*, *gapdh*, *acta* and *α-tubulin* in the case of male and female organisms. In gills, the most stable gene was *α-tubulin*, followed by *18S rRNA*, *actb*, *gapdh*, *efl-α*, *acta* in male and female organisms. In the case of the brain, the most stable gene was *α-tubulin*, followed by *18S rRNA*, *efl-α*, *actb*, *gapdh*, *α-tubulin* and *acta* in the case of male and female organisms. Similarly, in the spleen the most stable gene was *18S rRNA*, followed by *efl-α*, *gapdh*, *actb*, *α-tubulin* and *acta* in males while in females the most stable gene was *actb*, followed by *18S rRNA*, *efl-α*, *gapdh*, *acta*, and *α-tubulin*. In the heart, the most stable gene was *18S rRNA*, followed by *actb*, *gapdh*, *efl-α*, *α-tubulin*, and *acta* in males, while in females the most stable gene was *actb*, followed by *18S rRNA*, *gapdh*, *efl-α*, *acta* and *α-tubulin*. Finally, the most stable gene was *gapdh*, followed by *18S rRNA*, *actb*, *efl-α*, *acta* and *α-tubulin* in the male testis and in the case of the ovary in females, the most stable gene was *gapdh*, followed by *actb*, *18S rRNA*, *efl-α*, *α-tubulin* and *acta*.

Normfinder gene analysis

Normfinder software determines the stability of reference genes using a mathematical model to estimate intra- and intergroup expression variations. According

to the results evaluated in the eight tissues, the genes are shown from the most stable to the least stable (Fig. 3). In the case of the liver, the most stable gene was *actb*, followed by *ef1- α* , *18S rRNA*, *acta*, *gapdh*, and *α -tubulin* in both male and female organisms. While in the intestine the most stable gene was *18S rRNA*, followed by *actb*, *ef1- α* , *gapdh*, *α -tubulin*, and *acta* in the case of males, while in females the most stable gene was *18S rRNA*, followed by *actb*, *ef1- α* , *gapdh*, *acta*, and *α -tubulin*. Similarly, in the muscle and gill, the most stable gene was *actb*, followed by *18S rRNA*, *ef1- α* , *gapdh*, *acta*, and *α -tubulin* in both male and female organisms. In the case of the brain, the most stable gene was *α -tubulin* followed by *18S rRNA* of *actb*, *ef1- α* , *gapdh*, and *acta* in both male and female organisms. While in the case of the spleen, the most stable gene was *actb*, followed by *18S rRNA*, *ef1- α* , *gapdh*, *acta*, and *α -tubulin* in both male and female organisms. In the heart, the most stable gene was *18S rRNA*, followed by *actb*, *ef1- α* , *gapdh*, *α -tubulin*, and *acta* in both male and female organisms. Finally, in the testis of males and ovaries of females, the most stable gene was *gapdh* followed by *18S rRNA*, *actb*, *ef1- α* , *acta*, and *α -tubulin*.




Fig. 3. NormFinder analysis results of six candidate reference genes of *Atractosteus tropicus*.

geNorm gene analysis

The results obtained in the GeNorm software of the reference genes analyzed in the different tissues in both male and female adults of *Petenia splendida* take into account that M values less than 1.5 are considered the most stable genes (Fig. 4). In agreement with these results, in the case of the liver, the most stable genes were *18S rRNA*, *actb*, and *gapdh* in both males and females. Likewise, in the intestine, the *18S rRNA* and *gapdh* genes are the most stable in both males and females. In the case of muscle and gill, the most stable genes were *18S rRNA*, *actb*, and *gapdh* in both males and females. In this sense, in the brain the most stable gene was *α -tubulin*. In the case of the spleen and heart, the most stable genes were *18S rRNA* and *actb* in both males and females. Finally, in the case of the male testes and ovaries, the most stable genes were *18S rRNA* and *gapdh*.




Fig. 4. geNorm analysis results of six candidate reference genes of *Atractosteus tropicus*.

DISCUSSION

Real-time quantitative polymerase chain reaction (RT-qPCR) is a molecular biology technique that allows the analysis of relative changes in gene expression and is considered the “gold standard” in the field of gene quantification mRNA (Li *et al.*, 2019; Panina *et al.*, 2018; VanGuilder *et al.*, 2008). In this sense, to carry out a relative expression analysis, a process needs to be carried out, a reference gene or control gene is needed to compare it with our problem gene which we need to analyze.

The main characteristic that a reference gene must have is that its expression is constant regardless of the cell, tissues or organ, developmental restrictions, or physiological states of the organism (Kozera and Rapacz, 2013; Yang *et al.*, 2013). To determine the stability of a gene, the three most used software are BestKeeper, Normfinder and GenNorm, each applying different algorithms to select the best reference gene (Radonić *et al.*, 2004; Wang *et al.*, 2015).

According to our studies, the stability of six reference genes, *18S rRNA*, *efl-α*, *acta*, *actb*, *gapdh*, and *α-tubulin*, was analyzed through the three programs geNorm, Normfinder and Bestkeeper in different tissues in adult males and females of *Petenia splendida*. In this way, the reference genes analyzed have different functions such as *18S rRNA* plays an essential role in eukaryotic cells and ribosomes (Hadziavdic *et al.*, 2014; Poletto *et al.*, 2010). On the other hand, the *efl-α* gene is essential and highly conserved and plays an important function in the formation of the cytoskeleton of eukaryotic cells, serving as an essential center in protein networks (Romaus-Sanjurjo *et al.*, 2022; Zhou *et al.*, 2020). In this sense, two actin genes were analyzed, alpha-actin (*acta*) and beta-actin (*actb*), the first is a constituent of the cytoplasmic face of several types of cell adhesion sites (Otey and Carpen, 2004) and the second specifically plays a crucial role in various functions such as; cell migration, involved in blood-brain barrier pathways, cell division, muscle contraction, cell motility, secretion, phagocytosis, and gene expression (Bunnell *et al.*, 2011; Gu *et al.*, 2021). The *gapdh* gene participates in many cellular processes in addition to glycolysis (Lakho *et al.*, 2020; Tristan *et al.*, 2011). Finally, the *α-tubulin* gene is a fundamental element of microtubules that participate in the multiplication and movement of cells (Chen *et al.*, 2021; Janke, 2014).

According to our results obtained in the six reference genes analyzed in the different tissues in adults of *Petenia splendida*, it was not observed between male and female organisms. In the case of the liver and gill, the stable genes were *α-tubulin*, *actb*, *18S rRNA* and *gapdh*. These results agree with studies carried out on species such as

(*Oncorhynchus mykiss*) (Hook *et al.*, 2006), *Oreochromis niloticus* (Yang *et al.*, 2013), *Channa striatus* (Purohit *et al.*, 2016), *Ictalurus punctatus* (Small *et al.*, 2008), *Cynoglossus semilaevis* (Liu *et al.*, 2014), *Scophthalmus maximus* (Gao *et al.*, 2020) and *Danio rerio* (Jaramillo *et al.*, 2018; Rassier *et al.*, 2020), *Atractosteus tropicus* (Jiménez-Martínez *et al.*, 2022), *Paralichthys olivaceus* (Zheng and Sun, 2011), *Salmo salar* (Olsvik *et al.*, 2005). The authors of these works must consider these reference genes ideal for these tissues mainly to carry out gene expression studies according to the functions of each of them, such as the liver in the case of fish is considered the internal organ. largest constitutes 1% of its body mass whose main function is to metabolize all the substances that reach it through the blood, therefore, this organ serves as a histological reference for the analysis of tissue damage caused by inadequate nutrition or polluting effects. from the environment such as pesticides, heavy metals, and others (Li *et al.*, 2010; Olsvik *et al.*, 2007). It is important to highlight that according to the four genes analyzed in these tissues, several authors mention that act can be considered as a reference gene exclusive to the liver (Li *et al.*, 2010; Small *et al.*, 2008; Ye *et al.*, 2010). In contrast, in freshwater species such as *Basilichthys microlepidotus*, 60S ribosomal protein, L8, and EF1 are considered stable genes for the liver (Rojas-Hernandez *et al.*, 2019). In the case of the gill, it is the tissue and respiratory organs that extract oxygen from the water and the excretion of carbon dioxide, considering the main target organ for contaminants, its epithelial tissue is an excellent parameter to evaluate the effects of environmental variables, toxic substances, and water quality (Mazon *et al.*, 2002; Zhang *et al.*, 2021). Finally both the (De Marco *et al.*, 2021; Filby and Tyler, 2007).

In the case of intestine, muscle, spleen, and heart *18S rRNA*, *actb*, *gapdh* our results coincide with the work carried out in species such as *Salmo salar* (Kortner *et al.*, 2011), *Gasterosteus aculeatus* (Hibbeler *et al.*, 2008), *Ictalurus punctatus* (Small *et al.*, 2008) *Siniperca chiasitic*, *Oreochromis niloticus* (Yang *et al.*, 2013), *Ctenopharyngodon idella* (Su *et al.*, 2011), *Astatotilapia burtini* (Mobley *et al.*, 2022) where these tissues are made up of cell cytoskeletons, ribosomes and are involved in the processes of glycolysis. The *18S rRNA* gene is the most used for normalization to evaluate relative expression levels, especially in various studies, especially in cichlids such as *Oreochromis niloticus* (Huang *et al.*, 2012; Poletto *et al.*, 2010; Saranya *et al.*, 2021; Wang *et al.*, 2015; Yang *et al.*, 2013), *Oreochromis mossambicus* (Monjane-Mabuié *et al.*, 2022) and *Astatotilapia burtini* (Mobley *et al.*, 2022) are phylogenetically similar to *Petenia splendida*, which also coincide. with the phylogenetic study through

bioinformatic tools of the *18S rRNA* from the GenBank database for 31 species belonging to 13 genera of cichlid fish (Teleostei: Cichlidae) and conclude that In conclusion, the alignment of the *18S rRNA* genetic sequences between cichlid species also as a phylogenetic tree with bootstrap values revealed a high precision in phylogenetic relationship between species (Azab *et al.*, 2020). Similarly, the *18S rRNA* gene is recommended as a reference gene according to its great stability in various target tissues in marine fish species, the Asian sea bass (*Lates calcarifer*) (Paria *et al.*, 2016). The *actb* gene plays an important role in cellular processes such as cell division and the regulation of gene expression (Dominguez and Holmes, 2011; Ruan and Lai, 2007; Vandekerckhove and Weber, 1978). On the other hand, *actb* is considered the best reference gene in the case of teleost fish (Deloffre *et al.*, 2012; Gao *et al.*, 2020; Gunter *et al.*, 2011). *actb* is used in various studies to normalize molecular expression due to its high conservation as an endogenous maintenance gene and is constantly expressed in different types of cells such as intestinal, muscle, and cardiac cells, it is an excellent reference gene that is stable expressed in different tissues under normal physiological conditions and is used under laboratory conditions applied in experiments where fish are exposed to batteries, stress, osmoregulation, salinity and hypoxia (Jiménez-Martínez *et al.*, 2022; Paria *et al.*, 2016; Small *et al.*, 2008). On the other hand, in species such as *Monopterus albus* they mention that the expression of *actb* can vary and be considered an unstable gene in response to biochemical stimuli, during growth and differentiation, and in pathological states (Hu *et al.*, 2014; Ruan and Lai, 2007). It is important to mention that in the case of cichlids, *actb* is considered a reference gene for the normalization of relative expression in studies of genes involved in pigmentation related to evolution, adaptation, and speciation in cichlid species (Henning *et al.*, 2013). In this way, *gapdh* is a highly conserved gene in most organisms, fulfilling important functions in carbohydrate metabolism and participating in the regulation of mRNA stability (Sahoo *et al.*, 2023). Likewise, it is considered a suitable reference gene for various physiological studies in species such as *Salmo salar*, *Danio rerio*, *Cyprinus carpio*, *Monopterus albus*, and *Salvelinus alpinus* (Ahi *et al.*, 2018; Hu *et al.*, 2014; Jorgensen *et al.*, 2006; Tang *et al.*, 2007). *Gapdh* is a multifunctional gene that participates in various biological processes such as apoptosis and also stands out for being a good reference gene in teleost fish during the egg stage and the development of embryogenesis (Carnevali *et al.*, 2003; Sarropoulou *et al.*, 2011).

Likewise, in the case of testis and ovary, *gapdh* stands out as the most stable gene in adults of *Petenia splendida*, possibly due to the presence of vitellogenin in both female

and male organisms. In this sense, vitellogenin is proteins that contain glycogen and glyconjugates (Groh *et al.*, 2011; Tingaud-Sequeira *et al.*, 2012). Vitellogenin is produced in the liver of females, but amounts of vitellogenin mRNA are also present in the ovary itself (Sun *et al.*, 2023; Wang *et al.*, 2005). Likewise, work has been reported on the presence of mRNA in the testis of fish, mainly in wild fish due to the presence of endocrine disruptors present in the environment (Fang *et al.*, 2021; Pan *et al.*, 2019; Zezza *et al.*, 2020). Authors speculate that the encoded vitellogenin gene may provide nutrients for sperm development (Akasaka *et al.*, 2010, 2013).

In the case of the brain, the most stable gene was α -tubulin compared to the other genes; however, few studies consider α -tubulin as a reference gene specifically for the brain, in this case in species such as goldfish (*Carassius auratus*) and zebra fish (*Danio rerio*) mention that this gene is mainly expressed in the brain, nervous system and plays an important role in neuronal development and regeneration (Hieber *et al.*, 1998; Ilies *et al.*, 2012; Larson *et al.*, 2010; Senut *et al.*, 2004). Likewise, α -tubulin are components of microtubules where their adequate regulation depends on the processes of proliferation, differentiation, migration, growth and guidance of axons and dendrites, and formation of synapses in the different stages of development until adulthood. in the brain (Aiken *et al.*, 2017; Buscaglia *et al.*, 2020; Xie *et al.*, 2021).

Finally we can conclude that based on BestKeeper, Normfinder and GeNorm the most stable reference genes in most tissues were *18S rRNA* and *actb*, while in the case of the brain it was α -tubulin and finally *gapdh* in the ovary and testes, Likewise, no differences were shown in the expression in the analyzed tissues of females and males of *Petenia splendida*. Likewise, the reference gene can vary depending on the organism, tissue, and many other factors.

DECLARATIONS

Acknowledgments

Thanks to the Laboratory of Physiology in Aquatic Resources (LAFIRA) of the academic division of biological sciences of the Juarez University of Tabasco (DACBIOL-UJAT).

Funding

The study received no external funding.

Ethical approval

All applicable international, national, and/or institutional guidelines for the care and use of animals were followed by authors.

Generative AI and AI-assisted technology statement
.....?

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Ahi, E.P., Singh, P., Lecaudey, L.A., Gessl, W. and Sturmbauer, C., 2018. Maternal mRNA input of growth and stress-response-related genes in cichlids in relation to egg size and trophic specialization. *EvoDevo*, **9**: 23. <https://doi.org/10.1186/s13227-018-0112-3>
- Aiken, J., Buscaglia, G., Bates, E.A. and Moore, J.K., 2017. The α -tubulin gene TUBA1A in brain development: A key ingredient in the neuronal isotype blend. *J. Dev. Biol.*, **5**: 8. <https://doi.org/10.3390/jdb5030008>
- Akasaka, M., Harada, Y. and Sawada, H., 2010. Vitellogenin C-terminal fragments participate in fertilization as egg-coat binding partners of sperm trypsin-like proteases in the ascidian *Halocynthia roretzi*. *Biochem. biophys. Res. Commun.*, **392**: 479–484. <https://doi.org/10.1016/j.bbrc.2010.01.006>
- Akasaka, M., Kato, K.H., Kitajima, K. and Sawada, H., 2013. Identification of novel isoforms of vitellogenin expressed in ascidian eggs. *J. exp. Zool. B: Mol. Dev. Evol.*, **320**: 118–128. <https://doi.org/10.1002/jez.b.22488>
- Alvarez-González, C.A., Márquez-Couturier, G., Arias-Rodríguez, L., Contreras-Sánchez, W.M., Uscanga-Martínez, A., Perales-García, N., Moyano-López, F.J., Hernández-Jiménez, R., Civera-Cerecedo, R. and Goytortua-Bores, E., 2008. Avances en la fisiología digestiva y nutrición de la mojarra tenguayaca *Petenia splendida*. Avances En Nutrición Acuicola. IX Simposio Internacional de Nutrición Acuicola, Noviembre 24 – 27, Universidad Autónoma de Nuevo León, Nuevo León, pp. 135-235.
- Andersen, C.L., Jensen, J.L. and Ørntoft, T.F., 2004. Normalization of real-time quantitative reverse transcription-PCR data: A model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res.*, **64**: 5245–5250. <https://doi.org/10.1158/0008-5472.CAN-04-0496>
- Arias-Rodríguez, L., Ibarra-Castro, L. and Páramo-Delgadillo, S., 2008. Los cromosomas mitóticos y meióticos del pez tropical *Petenia splendida* (Cichlidae). *Rev. Biol. Trop.*, **56**: 895–907.
- Azab, M.S., Ahmed, S.H., El-Shaarawy, H.I., Kasem, N.R.A. and Hamza, D.S., 2020. Molecular phylogenetic correlation among cichlid fishes (Teleostei: Cichlidae) based on 18S rRNA Gene sequencing analysis. *Egypt. Acad. J. biol. Sci. C, Physiol. mol. Biol.*, **12**: 229–239. <https://doi.org/10.21608/eajbsc.2020.159710>
- Bunnell, T.M., Burbach, B.J., Shimizu, Y. and Ervasti, J.M., 2011. β -Actin specifically controls cell growth, migration, and the G-actin pool. *Mol. Biol. Cell*, **22**: 4047–4058. <https://doi.org/10.1091/mbc.e11-06-0582>
- Buscaglia, G., Northington, K.R., Moore, J.K. and Bates, E.A., 2020. Reduced TUBA1A tubulin causes defects in trafficking and impaired adult motor behavior. *Eneuro*, **7**. <https://doi.org/10.1523/ENEURO.0045-20.2020>
- Carnevali, O., Polzonetti, V., Cardinali, M., Pugnali, A., Natalini, P., Zmora, N., Mosconi, G. and Polzonetti-Magni, A.M., 2003. Apoptosis in sea bream *Sparus aurata* eggs. *Mol. Reprod. Dev. Incorporat. Gamete Res.*, **66**: 291–296. <https://doi.org/10.1002/mrd.10356>
- Casadei, R., Pelleri, M.C., Vitale, L., Facchin, F., Lenzi, L., Canaider, S., Strippoli, P. and Frabetti, F., 2011. Identification of housekeeping genes suitable for gene expression analysis in the zebrafish. *Gene Express. Patterns*, **11**: 271–276. <https://doi.org/10.1016/j.gexp.2011.01.003>
- Castillo-Collado, A. del C., Frías-Quintana, C.A., Morales-García, V., Alvarez-Villagomez, C.S., Asencio-Alcudia, G., Peña-Marín, E.S., Martínez-Bautista, G., Jiménez-Martínez, L.D. and Alvarez-González, C.A., 2022. Characterization and expression of the gene glucose transporter 2 (GLUT2) in embryonic, larval and adult Bay snook *Petenia splendida* (Cichliformes: Cichlidae). *Neotrop. Ichthyol.*, **20**. <https://doi.org/10.1590/1982-0224-2021-0171>
- Chen, J., Kholina, E., Szyk, A., Fedorov, V.A., Kovalenko, I., Gudimchuk, N. and Roll-Mecak, A., 2021. α -tubulin tail modifications regulate microtubule stability through selective effector recruitment, not changes in intrinsic polymer dynamics. *Dev. Cell*, **56**: 2016–2028. <https://doi.org/10.1016/j.devcel.2021.05.005>
- Chen, Z., Halford, N.G. and Liu, C., 2023. Real-time quantitative PCR: Primer design, reference gene selection, calculations and statistics. *Metabolites*, **13**: 806. <https://doi.org/10.3390/metabo13070806>
- Contreras-García, M.J., Contreras-Sánchez, W.M., Mcdonal-Vera, A. and Hernández-Vidal, U.,

2018. Masculinization of the native bay snook, *Petenia splendida*, using oral administration of the synthetic steroid 17 α -methyltestosterone. *J. World Aquacult. Soc.*, **49**: 889–896. <https://doi.org/10.1111/jwas.12449>
- De Marco, G., Brandão, F., Pereira, P., Pacheco, M. and Cappello, T., 2021. Organ-specific metabolome deciphering cell pathways to cope with mercury in wild fish (golden grey mullet *Chelon auratus*). *Animals*, **12**: 79. <https://doi.org/10.3390/ani12010079>
- Del Río-Portilla, M.A., Vargas-Peralta, C.E., Farfán, C., Barriga-Sosa, I. de los A. and García-De-León, F.J., 2016. The complete mitochondrial DNA of the bay snook, *Petenia splendida*, a native Mexican cichlid. *Mitochond. DNA Part A*, **27**: 1381–1382. <https://doi.org/10.3109/19401736.2014.947590>
- Deloffre, L.A.M., Andrade, A., Filipe, A.I., Canario, A.V.M., 2012. Reference genes to quantify gene expression during oogenesis in a teleost fish. *Gene*, **506**: 69–75. <https://doi.org/10.1016/j.gene.2012.06.047>
- Dominguez, R. and Holmes, K.C., 2011. Actin structure and function. *Annu. Rev. Biophys.*, **40**: 169–186. <https://doi.org/10.1146/annurev-biophys-042910-155359>
- Fang, G.Z., Huang, G.Y., Ying, G.G., Qiu, S.Q., Shi, W.J., Xie, L., Yang, Y.Y. and Ma, D.D., 2021. Endocrine disrupting effects of binary mixtures of 17 β -estradiol and testosterone in adult female western mosquitofish (*Gambusia affinis*). *Ecotoxicol. Environ. Saf.*, **208**: 111566. <https://doi.org/10.1016/j.ecoenv.2020.111566>
- Filby, A.L. and Tyler, C.R., 2007. Appropriate housekeeping genes for use in expression profiling the effects of environmental estrogens in fish. *BMC mol. Biol.*, **8**: 1–13. <https://doi.org/10.1186/1471-2199-8-10>
- Gao, Y., Gao, Y., Huang, B., Meng, Z. and Jia, Y., 2020. Reference gene validation for quantification of gene expression during ovarian development of turbot (*Scophthalmus maximus*). *Sci. Rep.*, **10**: 823. <https://doi.org/10.1038/s41598-020-57633-3>
- Groh, K.J., Nesatyy, V.J., Segner, H., Eggen, R.I.L. and Suter, M.J.F., 2011. Global proteomics analysis of testis and ovary in adult zebrafish (*Danio rerio*). *Fish Physiol. Biochem.*, **37**: 619–647. <https://doi.org/10.1007/s10695-010-9464-x>
- Gu, Y., Tang, S., Wang, Z., Cai, L., Lian, H., Shen, Y. and Zhou, Y., 2021. A pan-cancer analysis of the prognostic and immunological role of β -actin (ACTB) in human cancers. *Bioengineered*, **12**: 6166–6185. <https://doi.org/10.1080/21655979.2021.1973220>
- Gunter, H.M., Clabaut, C., Salzburger, W. and Meyer, A., 2011. Identification and characterization of gene expression involved in the coloration of cichlid fish using microarray and qRT-PCR approaches. *J. mol. Evol.*, **72**: 127–137. <https://doi.org/10.1007/s00239-011-9431-x>
- Hadziavdic, K., Lekang, K., Lanzen, A., Jonassen, I., Thompson, E.M. and Troedsson, C., 2014. Characterization of the 18S rRNA Gene for designing universal eukaryote specific primers. *PLoS One*, **9**: e87624. <https://doi.org/10.1371/journal.pone.0087624>
- Henning, F., Jones, J.C., Franchini, P. and Meyer, A., 2013. Transcriptomics of morphological color change in polychromatic Midas cichlids. *BMC Genom.*, **14**: 1–14. <https://doi.org/10.1186/1471-2164-14-171>
- Hibbeler, S., Scharsack, J.P. and Becker, S., 2008. Housekeeping genes for quantitative expression studies in the three-spined stickleback *Gasterosteus aculeatus*. *BMC Mol. Biol.*, **9**: 1–10. <https://doi.org/10.1186/1471-2199-9-18>
- Hieber, V., Dai, X., Foreman, M. and Goldman, D., 1998. Induction of α 1-tubulin gene expression during development and regeneration of the fish central nervous system. *J. Neurobiol.*, **37**: 429–440. [https://doi.org/10.1002/\(SICI\)1097-4695\(19981115\)37:3<429::AID-NEU8>3.0.CO;2-N](https://doi.org/10.1002/(SICI)1097-4695(19981115)37:3<429::AID-NEU8>3.0.CO;2-N)
- Hook, S.E., Skillman, A.D., Small, J.A. and Schultz, I.R., 2006. Gene expression patterns in rainbow trout, *Oncorhynchus mykiss*, exposed to a suite of model toxicants. *Aquat. Toxicol.*, **77**: 372–385. <https://doi.org/10.1016/j.aquatox.2006.01.007>
- Hu, Q., Guo, W., Gao, Y., Tang, R. and Li, D., 2014a. Reference gene selection for real-time RT-PCR normalization in rice field eel (*Monopterus albus*) during gonad development. *Fish Physiol. Biochem.*, **40**: 1721–1730. <https://doi.org/10.1007/s10695-014-9962-3>
- Huang, C.W., Li, Y.H., Hu, S.Y., Chi, J.R., Lin, G.H., Lin, C.C., Gong, H.Y., Chen, J.Y., Chen, R.H. and Chang, S.J., 2012. Differential expression patterns of growth-related microRNAs in the skeletal muscle of Nile tilapia (*Oreochromis niloticus*). *J. Anim. Sci.*, **90**: 4266–4279. <https://doi.org/10.2527/jas.2012-5142>
- Ilies, I., Zupanc, M.M. and Zupanc, G.K.H., 2012. Proteome analysis reveals protein candidates involved in early stages of brain regeneration of

- teleost fish. *Neuroscience*, **219**: 302–313. <https://doi.org/10.1016/j.neuroscience.2012.05.028>
- Infante, C., Matsuoka, M.P., Asensio, E., Cañavate, J.P., Reith, M. and Manchado, M., 2008. Selection of housekeeping genes for gene expression studies in larvae from flatfish using real-time PCR. *BMC Mol. Biol.*, **9**: 1–12. <https://doi.org/10.1186/1471-2199-9-28>
- Janke, C., 2014. The tubulin code: Molecular components, readout mechanisms, and functions. *J. Cell Biol.*, **206**: 461–472. <https://doi.org/10.1083/jcb.201406055>
- Jaramillo, M.L., Pereira, A.G., Davico, C.E., Nezzi, L., Ammar, D., Müller, Y.M.R. and Nazari, E.M., 2018. Evaluation of reference genes for reverse transcription-quantitative PCR assays in organs of zebrafish exposed to glyphosate-based herbicide, Roundup. *Animal*, **12**: 1424–1434. <https://doi.org/10.1017/S1751731117003111>
- Jiménez M.L.D., Morales, G.V., Frias, Q.C.A., Carmen, C.C.A., Asencio, A.G.G., Alvarez, V.C.S., Peña Marín, E.S., Concha, F.B. and Alvarez-Gonzalez, C.A., 2022. Quality evaluation of reference gene expression on different tissues in adults of tropical gar *Atractosteus tropicus*. *Pakistan J. Zool.*, **54**: 363–372. <https://doi.org/10.17582/journal.pjz/20200913180928>
- Jiménez-Martínez, L.D., Alvarez-González, C.A., Contreras-Sánchez, W.M., Márquez-Couturier, G., Arias-Rodríguez, L. and Almeida-Madrigal, J.A., 2009. Evaluation of larval growth and survival in Mexican mojarra, *Cichlasoma urophthalmus*, and Bay Snook, *Petenia splendida*, under different initial stocking densities. *J. World Aquacult. Soc.*, **40**: 753–761. <https://doi.org/10.1111/j.1749-7345.2009.00295.x>
- Jorgensen, S.M., Kleveland, E.J., Grimholt, U. and Gjoen, T., 2006. Validation of reference genes for real-time polymerase chain reaction studies in Atlantic salmon. *Mar. Biotechnol.*, **8**: 398–408. <https://doi.org/10.1007/s10126-005-5164-4>
- Jothinarayanan, N., Karlsen, F. and Roseng, L.E., 2023. Comparative evaluation of loop-mediated isothermal amplification and PCR for detection of *Esox lucius* housekeeping genes for use in on-site environmental monitoring. *J. Fish Biol.*, **103**: 897–905. <https://doi.org/10.1111/jfb.15476>
- Kortner, T.M., Valen, E.C., Kortner, H., Marjara, I.S., Krogdahl, Å. and Bakke, A.M., 2011. Candidate reference genes for quantitative real-time PCR (qPCR) assays during development of a diet-related enteropathy in Atlantic salmon (*Salmo salar* L.) and the potential pitfalls of uncritical use of normalization software tools. *Aquaculture*, **318**: 355–363. <https://doi.org/10.1016/j.aquaculture.2011.05.038>
- Kozera, B. and Rapacz, M., 2013. Reference genes in real-time PCR. *J. appl. Genet.*, **54**: 391–406. <https://doi.org/10.1007/s13353-013-0173-x>
- Kuah, M.K., Jaya-Ram, A. and Shu-Chien, A.C., 2016. A fatty acyl desaturase (fads2) with dual $\Delta 6$ and $\Delta 5$ activities from the freshwater carnivorous striped snakehead *Channa striata*. *Compr. Biochem. Physiol. A: Mol. Integr. Physiol.*, **201**: 146–155. <https://doi.org/10.1016/j.cbpa.2016.07.007>
- Kumari, R., Srivastava, P.P., Mohanta, K.N., Kumar, R., Das, P., Sahoo, L., Das, G., Nandanpawar, P.C., Debbarma, J., Krishna, G. and Siddaiah, G.M., 2022. Evaluation of candidate reference genes for RT-qPCR analysis during developmental stages in striped murrel (*Channa striata*). *Aquacult. Res.*, **53**: 6868–6877. <https://doi.org/10.1111/are.16152>
- Lakho, S.A., Haseeb, M., Huang, J., Yang, Z., Hasan, M.W., Aleem, M.T., Naqvi, M.A.H., Memon, M.A., Song, X. and Yan, R., 2020. Glyceraldehyde-3-phosphate dehydrogenase from *Eimeria acervulina* modulates the functions of chicken dendritic cells to boost Th1 type immune response and stimulates autologous CD4+ T cells differentiation *in-vitro*. *Vet. Res.*, **51**: 1–14. <https://doi.org/10.1186/s13567-020-00864-z>
- Larson, T.A., Gordon, T.N., Lau, H.E. and Parichy, D.M., 2010. Defective adult oligodendrocyte and Schwann cell development, pigment pattern, and craniofacial morphology in puma mutant zebrafish having an alpha tubulin mutation. *Dev. Biol.*, **346**: 296–309. <https://doi.org/10.1016/j.ydbio.2010.07.035>
- Li, J.Y., Chen, W.Z., Yang, S.H., Xu, C.L., Huang, X., Chen, C. and Xie, H., 2019. Screening of reference genes in real-time PCR for *Radopholus similis*. *PeerJ*, **7**: e6253. <https://doi.org/10.7717/peerj.6253>
- Li, Y.Y., Chen, X., Yang, J.X., Chen, Q., Song, T.Y. and Ge, J.Q., 2023. Evaluation of housekeeping genes as references for quantitative real-time PCR analysis of *Eun eelropea*, *Anguilla anguilla*. *J. Fish Biol.*, **102**: 141–154. <https://doi.org/10.1111/jfb.15247>
- Li, Z., Yang, L., Wang, J., Shi, W., Pawar, R.A., Liu, Y., Xu, C., Cong, W., Hu, Q. and Lu, T., 2010. β -Actin is a useful internal control for tissue-specific gene expression studies using quantitative real-time PCR in the half-smooth tongue sole *Cynoglossus semilaevis* challenged with LPS or

- Vibrio anguillarum*. *Fish Shellf. Immunol.*, **29**: 89–93. <https://doi.org/10.1016/j.fsi.2010.02.021>
- Liu, C., Xin, N., Zhai, Y., Jiang, L., Zhai, J., Zhang, Q. and Qi, J., 2014. Reference gene selection for quantitative Real-Time RT-PCR normalization in the half-smooth tongue sole (*Cynoglossus semilaevis*) at different developmental stages, in various tissue types and on exposure to chemicals. *PLoS One*, **9**: e91715. <https://doi.org/10.1371/journal.pone.0091715>
- Liu, Y., Wu, S., Jiang, N., Liu, W., Zhou, Y., Zeng, L., Zhong, Q., Li, Z. and Fan, Y., 2022. Characterization of reference genes for qRT-PCR normalization in rice-field eel (*Monopterus albus*) to assess differences in embryonic developmental stages, the early development of immune organs, and cells infected with rhabdovirus. *Fish Shellfish Immunol.*, **120**: 92–101. <https://doi.org/10.1016/j.fsi.2021.11.021>
- Liu, Z., Wang, T., Liu, P., Jiang, D., Liu, X., Deng, S., Wu, T., Huang, Y., Zhu, C., Li, G. and Jiang, M., 2023. Selection of reference gene for expression studies in the ovary and pituitary of spotted scat (*Scatophagus argus*) at different ovarian stages. *Fishes*, **8**. <https://doi.org/10.3390/fishes8020120>
- Matthews, M. and Varga, Z.M., 2012. Anestesia y eutanasia en pez cebra. *Revista ILAR.*, **53**: 192–204. <https://doi.org/10.1093/ilar.53.2.192>
- Mazon, A.F., Cerqueira, C.C.C. and Fernandes, M.N., 2002. Gill cellular changes induced by copper exposure in the South American tropical freshwater fish *Prochilodus scrofa*. *Environ. Res.*, **88**: 52–63. <https://doi.org/10.1006/enrs.2001.4315>
- McCurley, A.T. and Callard, G.V., 2008. Characterization of housekeeping genes in zebrafish: Male-female differences and effects of tissue type, developmental stage and chemical treatment. *BMC Mol. Biol.*, **9**: 102. <https://doi.org/10.1186/1471-2199-9-102>
- Méndez-García, C.A., 2010. *Revisión sistemática del complejo Petenia splendida (Teleostei: Cichlidae) en áreas selectas de Guatemala, Centro América*. *Rev. Biol. Trop.*, **59**: 1205–1216. <https://doi.org/10.15517/rbt.v0i0.3392>
- Mobley, R.B., Ray, E.J. and Maruska, K.P., 2022. Expression and localization of neuronal nitric oxide synthase in the brain and sensory tissues of the African cichlid fish *Astatotilapia burtoni*. *J. Comp. Neurol.*, **530**: 2901–2917. <https://doi.org/10.1002/cne.25383>
- Monjane-Mabuie, A., Mondlane-Milisse, A., Pedro, O., Leão-Buchir, J. and Correia, D., 2022. Mercury pollution assessment and metallothionein gene expression in tilapia (*Oreochromis mossambicus*): A case study of revuê river in manica, mozambique. *Rendiconti Lincei. Sci. Fisiche Natur.*, **33**: 513–526. <https://doi.org/10.1007/s12210-022-01092-7>
- Olsvik, P.A., Lie, K.K., Jordal, A.E.O., Nilsen, T.O. and Hordvik, I., 2005. Evaluation of potential reference genes in real-time RT-PCR studies of Atlantic salmon. *BMC Mol. Biol.*, **6**: 1–9. <https://doi.org/10.1186/1471-2199-6-21>
- Olsvik, P.A., Lie, K.K., Sæle, Ø. and Sanden, M., 2007. Spatial transcription of CYP1A in fish liver. *BMC Physiol.*, **7**: 12. <https://doi.org/10.1186/1472-6793-7-12>
- Olsvik, P.A., Sjøteland, L. and Lie, K.K., 2008. Selection of reference genes for qRT-PCR examination of wild populations of Atlantic cod *Gadus morhua*. *BMC Res. Notes*, **1**: 1–9. <https://doi.org/10.1186/1756-0500-1-47>
- Otey, C.A. and Carpen, O., 2004. Alpha-actinin revisited: A fresh look at an old player. *Cell Motil. Cytosk.*, **58**: 104–111. <https://doi.org/10.1002/cm.20007>
- Øvergård, A.C., Nerland, A.H. and Patel, S., 2010. Evaluation of potential reference genes for real time RT-PCR studies in Atlantic halibut (*Hippoglossus hippoglossus* L.), during development, in tissues of healthy and NNV-injected fish, and in anterior kidney leucocytes. *BMC Mol. Biol.*, **11**: 36. <https://doi.org/10.1186/1471-2199-11-36>
- Pan, X., Liu, Y., Zhou, K., Mu, X., Zheng, S., Liu, C. and Hu, Y., 2019. Tissue expression and bioinformatics analysis of the vitellogenin gene of Asian arowana (*Scleropages formosus*). *J. appl. Ichthyol.*, **35**: 970–977. <https://doi.org/10.1111/jai.13927>
- Panina, Y., Germond, A., Masui, S. and Watanabe, T.M., 2018. Validation of common housekeeping genes as reference for qPCR Gene expression analysis during iPS reprogramming process. *Sci. Rep.*, **8**: 8716. <https://doi.org/10.1038/s41598-018-26707-8>
- Paria, A., Dong, J., Babu, P.P., Makesh, M., Chaudhari, A., Thirunavukkarasu, A.R., Purushothaman, C.S. and Rajendran, K.V., 2016. Evaluation of candidate reference genes for quantitative expression studies in Asian seabass (*Lates calcarifer*) during ontogenesis and in tissues of healthy and infected fishes. *Indian J. Exp. Biol.*, **54**: 597–605.
- Pérez-Sánchez, E. and Páramo-Delgadillo, S., 2008. The culture of cichlids of southeastern Mexico. *Aquacult. Res.*, **39**: 777–783. <https://doi.org/10.1111/j.1365-2109.2008.01929.x>
- Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucl.*

- Acids Res.*, **29**: e45–e45. <https://doi.org/10.1093/nar/29.9.e45>
- Pfaffl, M.W., Tichopad, A., Prgomet, C. and Neuvians, T.P., 2004. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper–Excel-based tool using pair-wise correlations. *Biotechnol. Lett.*, **26**: 509–515. <https://doi.org/10.1023/B:BILE.0000019559.84305.47>
- Poletto, A.B., Ferreira, I.A. and Martins, C., 2010. The B chromosomes of the African cichlid fish *Haplochromis obliquidens* harbour 18S rRNA gene copies. *BMC Genet.*, **11**: 1–8. <https://doi.org/10.1186/1471-2156-11-1>
- Purohit, G.K., Mahanty, A., Mohanty, B.P. and Mohanty, S., 2016. Evaluation of housekeeping genes as references for quantitative real-time PCR analysis of gene expression in the murrelet *Channa striatus* under high-temperature stress. *Fish Physiol. Biochem.*, **42**: 125–135. <https://doi.org/10.1007/s10695-015-0123-0>
- Radonić, A., Thulke, S., Mackay, I.M., Landt, O., Siebert, W. and Nitsche, A., 2004. Guideline to reference gene selection for quantitative real-time PCR. *Biochem. biophys. Res. Commun.*, **313**: 856–862. <https://doi.org/10.1016/j.bbrc.2003.11.177>
- Rassier, G.T., Silveira, T.L.R., Remião, M.H., Daneluz, L.O., Martins, A.W.S., Dellagostin, E.N., Ortiz, H.G., Domingues, W.B., Komninou, E.R., Kütter, M.T., Marins, L.F.F. and Campos, V.F., 2020. Evaluation of qPCR reference genes in GH-overexpressing transgenic zebrafish (*Danio rerio*). *Sci. Rep.*, **10**: 12692. <https://doi.org/10.1038/s41598-020-69423-y>
- Rodríguez-Estrada, U., Méndez-Marín, O., Pérez-Morales, A., Martínez-García, R., Peña-Marín, E., Civera-Cerecedo, R., Goytortua-Bores, E.E., Martínez-Yáñez, R. and Álvarez-González, C.A., 2020. Lipid requirement of bay snook (*Petenia splendida* Günther, 1862) juveniles. *Latin Am. J. aquat. Res.*, **48**: 674–685. <https://doi.org/10.3856/vol48-issue4-fulltext-2429>
- Rojas-Hernandez, N., Véliz, D. and Vega-Retter, C., 2019. Selection of suitable reference genes for gene expression analysis in gills and liver of fish under field pollution conditions. *Sci. Rep.*, **9**: 3459. <https://doi.org/10.1038/s41598-019-40196-3>
- Romaus-Sanjurjo, D., Saikia, J.M., Kim, H.J., Tsai, K.M., Le, G.Q. and Zheng, B., 2022. Overexpressing eukaryotic elongation factor 1 alpha (eEF1A) proteins to promote corticospinal axon repair after injury. *Cell Death Discov.*, **8**: 390. <https://doi.org/10.1038/s41420-022-01186-z>
- Ruan, W. and Lai, M., 2007. Actin, a reliable marker of internal control? *Clin. Chim. Acta*, **385**: 1–5. <https://doi.org/10.1016/j.cca.2007.07.003>
- Sahoo, P.K., Parida, S., Parida, S., Parida, P. and Paul, A., 2023. Stability evaluation and validation of appropriate reference genes for real-time PCR expression analysis of immune genes in the rohu (*Labeo rohita*) skin following argulosis. *Sci. Rep.*, **13**: 2660. <https://doi.org/10.1038/s41598-023-29325-1>
- Salem, M., Paneru, B., Al-Tobasei, R., Abdouni, F., Thorgaard, G.H., Rexroad, C.E. and Yao, J., 2015. Transcriptome assembly, gene annotation and tissue gene expression atlas of the rainbow trout. *PloS One*, **10**: e0121778. <https://doi.org/10.1371/journal.pone.0121778>
- Saranya, K., Manivasagan, V., Anbalagan, S., Sathasivam, J., Sureshbabu, K., Gopi, K., Raj, C.L., Sreethaar, S. M. and Patil, R., 2021. Molecular systematic investigations of three fin fish cichlid species of *Oreochromis niloticus* (Nile tilapia), genetically improved farmed tilapia (GIFT) and *Astronotus ocellatus* (Oscar Cichlid). *Let. appl. NanoBioSci.*, **10**: 2850 - 2860. <https://doi.org/10.33263/LIANBS104.28502860>
- Sarropoulou, E., Nousdili, D., Kotoulas, G. and Magoulas, A., 2011. Functional divergences of GAPDH isoforms during early development in two perciform fish species. *Mar. Biotechnol.*, **13**: 1115–1124. <https://doi.org/10.1007/s10126-011-9375-6>
- Senut, M.C., Gulati-Leekha, A. and Goldman, D., 2004. An element in the α 1-tubulin promoter is necessary for retinal expression during optic nerve regeneration but not after eye injury in the adult zebrafish. *J. Neurosci.*, **24**: 7663–7673. <https://doi.org/10.1523/JNEUROSCI.2281-04.2004>
- Small, B.C., Murdock, C.A., Bilodeau-Bourgeois, A.L., Peterson, B.C. and Waldbieser, G.C., 2008. Stability of reference genes for real-time PCR analyses in channel catfish (*Ictalurus punctatus*) tissues under varying physiological conditions. *Compr. Biochem. Physiol. B: Biochem. Mol. Biol.*, **151**: 296–304. <https://doi.org/10.1016/j.cbpb.2008.07.010>
- Su, J., Dong, J., Huang, T., Zhang, R., Yang, C. and Heng, J., 2011. Myeloid differentiation factor 88 gene is involved in antiviral immunity in grass carp *Ctenopharyngodon idella*. *J. Fish Biol.*, **78**: 973–979. <https://doi.org/10.1111/j.1095-8649.2011.02910.x>
- Su, J., Zhang, R., Dong, J. and Yang, C., 2011.

- Evaluation of internal control genes for qRT-PCR normalization in tissues and cell culture for antiviral studies of grass carp (*Ctenopharyngodon idella*). *Fish Shellf. Immunol.*, **30**: 830–835. <https://doi.org/10.1016/j.fsi.2011.01.006>
- Sun, S.X., Liu, Y.C., Limbu, S.M., Li, D.L., Chen, L.Q., Zhang, M.L., Yin, Z., Du, Z.Y., 2023. Vitellogenin 1 is essential for fish reproduction by transporting DHA-containing phosphatidylcholine from liver to ovary. *Biochim. biophys. Acta Mol. Cell Biol. Lipids*, **1868**: 159289. <https://doi.org/10.1016/j.bbalip.2023.159289>
- Tang, R., Dodd, A., Lai, D., McNabb, W.C. and Love, D.R., 2007. Validation of zebrafish (*Danio rerio*) reference genes for quantitative real-time RT-PCR normalization. *Acta Biochim. Biophys. Sin.*, **39**: 384–390. <https://doi.org/10.1111/j.1745-7270.2007.00283.x>
- Tang, Y., Yu, J., Xu, P., Li, J., Li, H. and Ren, H., 2012. Identification of housekeeping genes suitable for gene expression analysis in Jian carp (*Cyprinus carpio* var. jian). *Fish Shellf. Immunol.*, **33**: 775–779. <https://doi.org/10.1016/j.fsi.2012.06.027>
- Tingaud-Sequeira, A., Knoll-Gellida, A., André, M. and Babin, P.J., 2012. Vitellogenin expression in white adipose tissue in female teleost fish. *Biol. Reprod.*, **86**: 31–38. <https://doi.org/10.1095/biolreprod.111.093757>
- Treviño, L., Alvarez-González, C.A., Perales-García, N., Arévalo-Galán, L., Uscanga-Martínez, A., Márquez-Couturier, G., Fernández, I. and Gisbert, E., 2011. A histological study of the organogenesis of the digestive system in bay snook *Petenia splendida* Günther, 1862 from hatching to the juvenile stage. *J. appl. Ichthyol.*, **27**: 73–82. <https://doi.org/10.1111/j.1439-0426.2010.01608.x>
- Tristan, C., Shahani, N., Sedlak, T.W. and Sawa, A., 2011. The diverse functions of GAPDH: Views from different subcellular compartments. *Cell. Signal.*, **23**: 317–323. <https://doi.org/10.1016/j.cellsig.2010.08.003>
- Uscanga-Martínez, A., Perales-García, N., Alvarez-González, C.A., Moyano, F.J., Tovar-Ramírez, D., Gisbert, G.E., Márquez-Couturier, G., Contreras-Sánchez, W.M., Arias-Rodríguez, L. and Indy, J.R., 2011. Changes in digestive enzyme activity during initial ontogeny of bay snook *Petenia splendida*. *Fish Physiol. Biochem.*, **37**: 667–680. <https://doi.org/10.1007/s10695-011-9467-2>
- Vandekerckhove, J. and Weber, K., 1978. At least six different actins are expressed in a higher mammal: An analysis based on the amino acid sequence of the amino-terminal tryptic peptide. *J. mol. Biol.*, **126**: 783–802. [https://doi.org/10.1016/0022-2836\(78\)90020-7](https://doi.org/10.1016/0022-2836(78)90020-7)
- Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A. and Speleman, F., 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.*, **3**: 1–12. <https://doi.org/10.1186/gb-2002-3-7-research0034>
- VanGuilder, H.D., Vrana, K.E. and Freeman, W.M., 2008. Twenty-five years of quantitative PCR for gene expression analysis. *BioTechniques*, **44**: 619–626. <https://doi.org/10.2144/000112776>
- Vidal-López, J.M., Alvarez-González, C.A., Contreras-Sánchez, W.M. and Hernández-Vidal, U., 2009. Masculinización del ciclido nativo Tenhuayaca, *Petenia splendida* (Günther, 1862), usando nauplios de *Artemia* como vehículo del esteroide 17- α metiltestosterona. *Hidrobiológica*, **19**: 211–216.
- Wang, E., Wang, K., Chen, D., Wang, J., He, Y., Long, B., Yang, L., Yang, Q., Geng, Y. and Huang, X., 2015. Evaluation and selection of appropriate reference genes for real-time quantitative PCR analysis of gene expression in Nile tilapia (*Oreochromis niloticus*) during vaccination and infection. *Int. J. mol. Sci.*, **16**: 9998–10015. <https://doi.org/10.3390/ijms16059998>
- Wang, H., Tan, J.T.T., Emelyanov, A., Korzh, V. and Gong, Z., 2005. Hepatic and extrahepatic expression of vitellogenin genes in the zebrafish, *Danio rerio*. *Gene*, **356**: 91–100. <https://doi.org/10.1016/j.gene.2005.03.041>
- Xie, L., Huang, J., Dai, L., Luo, J., Zhang, J., Peng, Q., Sun, J. and Zhang, W., 2021. Loss-of-Function plays a major role in early neurogenesis of tubulin α -1 a (TUBA1A) mutation-related brain malformations. *Mol. Neurobiol.*, **58**: 1291–1302. <https://doi.org/10.1007/s12035-020-02193-w>
- Yang, C.G., Wang, X.L., Tian, J., Liu, W., Wu, F., Jiang, M. and Wen, H., 2013. Evaluation of reference genes for quantitative real-time RT-PCR analysis of gene expression in Nile tilapia (*Oreochromis niloticus*). *Gene*, **527**: 183–192. <https://doi.org/10.1016/j.gene.2013.06.013>
- Yang, X., Tong, G., Dong, L., Yan, T., Xu, H., Tang, G., Zhang, Y., Ma, K., Yin, J. and Kuang, Y., 2022. Evaluation of qPCR reference genes for taimen (*Hucho taimen*) under heat stress. *Sci. Rep.*, **12**: 313. <https://doi.org/10.1038/s41598-021-03872-x>
- Ye, X., Zhang, L., Dong, H., Tian, Y., Lao, H., Bai, J. and Yu, L., 2010. Validation of reference genes

- of grass carp *Ctenopharyngodon idellus* for the normalization of quantitative real-time PCR. *Biotechnol. Lett.*, **32**: 1031–1038. <https://doi.org/10.1007/s10529-010-0258-0>
- Zezza, D., Bisegna, A., Angelozzi, G., Merola, C., Conte, A., Amorena, M. and Perugini, M., 2020. Impact of endocrine disruptors on vitellogenin concentrations in wild brown trout (*Salmo trutta trutta*). *Bull. Environ. Contamin. Toxicol.*, **105**: 218–223. <https://doi.org/10.1007/s00128-020-02916-8>
- Zhang, F., Xu, J., Wang, X., Jabeen, K. and Li, D., 2021. Microplastic contamination of fish gills and the assessment of both quality assurance and quality control during laboratory analyses. *Mar. Pollut. Bull.*, **173**: 113051. <https://doi.org/10.1016/j.marpolbul.2021.113051>
- Zheng, W. and Sun, L., 2011. Evaluation of housekeeping genes as references for quantitative real time RT-PCR analysis of gene expression in Japanese flounder (*Paralichthys olivaceus*). *Fish Shellfish Immunol.*, **30**: 638–645. <https://doi.org/10.1016/j.fsi.2010.12.014>
- Zhong, Z., Ao, L., Zhao, L., Zhang, Z. and Jiang, Y., 2022. Screening and validation of reference genes for qPCR analysis in gonads and embryos of *Takifugu bimaculatus*. *Aquacult. Fish.*, **7**. <https://doi.org/10.1016/j.aaf.2020.10.002>
- Zhou, H., Guan, Y., Feng, M., Fu, Y., Tachibana, H. and Cheng, X., 2020. Evaluation on elongation factor 1 alpha of *Entamoeba histolytica* interaction with the intermediate subunit of the Gal/GalNAc lectin and actin in phagocytosis. *Pathogens*, **9**: 702 <https://doi.org/10.3390/pathogens9090702>
- Zhou, R.X., Meng, T., Meng, H.B., Cheng, D.X., Bin, S.Y., Cheng, J., Fu, G.H., Chu, W.Y. and Zhang, J.S., 2010. Selection of reference genes in transcription analysis of gene expression of the Mandarin fish, *Siniperca chuatsi*. *Zool. Res.*, **31**: 141–146. <https://doi.org/10.3724/SP.J.1141.2010.02141>