




Supplementary Material

Genetic Analysis of Antimicrobial Resistance Genes in *Salmonella* Isolated from Diseased Broilers in Egypt

Mona F. Shousha^{1*}, Aml M. Ragab² and Salwa M. Helmy¹

¹Department of Bacteriology, Mycology and Immunology, Faculty of Veterinary Medicine, Kafr El-Sheikh University, Kafr El-Sheikh 33516, Egypt

²Unit of Bacteriology, Animal Health Research Institute, Tanta Branch, Agriculture Research Center (ARC), Egypt



Supplementary Fig. 1. Agarose gel electrophoretic PCR pattern for detecting *qnrA*, *qnrB*, *qnrS*, *bla*_{TEM}, *bla*_{SHV} and *bla*_{CMY-2} at 543 bp, 562 bp, 491 bp, 516 bp, 392 bp and 462 bp respectively L: Ladder from 100 bp to 1000 bp P: Positive control N: Negative control: field isolate tested and confirmed by PCR to be negative for the related genes Lane 1 to 10: Negative amplification of *qnrA*, *qnrB* and *bla*_{CMY-2} and Positive amplification of *qnrS*, *bla*_{TEM} and *bla*_{SHV}

* Corresponding author: mona_shousha95@yahoo.com
0030-9923/2024/0006-2763 \$ 9.00/0



Copyright 2024 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).