



Supplementary Material

Occupancy Patterns of *Acanthodactylus cantoris* and *Eutropis dissimilis* in Khyber Pakhtunkhwa, Pakistan

Muhammad Shehzad¹, Muhammad Rais², Syed Mohsin Shah¹, Tauheed Ullah Khan³, Waseem Ahmed², Sanaullah Khan⁴ and Abdul Majid^{1*}

¹Department of Zoology, Kohat University of Science and Technology, Kohat, Khyber Pakhtunkhwa, Pakistan

²Herpetology lab, Department of Zoology, Wildlife, and Fisheries, PMAS-Arid Agriculture University, Rawalpindi, Pakistan

³Guangdong Key Laboratory of Animal Conservation and Resource Utilization, Guangdong Public Laboratory of Wild Animal Conservation and Utilization Institute of Zoology, Guangdong Academy of Sciences, Guangzhou, China

⁴Department of Zoology, University of Peshawar, Peshawar, Khyber Pakhtunkhwa, Pakistan



Supplementary Fig. 1. Indian fringe-fingered lizard (*Acanthodactylus cantoris*) (A) and Striped Grass Skink (*Eutropis dissimilis*) (B).

* Corresponding author: abdulmajid@kust.edu.pk
0030-9923/2025/0005-2307 \$ 9.00/00



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access  article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).