



## Supplementary Material

# Seasonal and Phenological Relationship Between Spiders and Sucking Insect Pests in Spring and Fall Maize Crops

Nisar Ahmad<sup>1\*</sup>, Sidra Wajid<sup>1,2,4\*</sup>, Donghui Wu<sup>2,3,4</sup>, Liang Chang<sup>3,4</sup>, Ahmad Nawaz<sup>5</sup>, Muhammad Aslam<sup>6</sup> and Syed Makhdoom Hussain<sup>7</sup>

<sup>1</sup>Department of Zoology, University of Jhang, Jhang 35200, Pakistan

<sup>2</sup>School of Environment, Northeast Normal University, Changchun City, Jilin Province, China

<sup>3</sup>State Key Laboratory of Black Soil Conservation and Utilization, Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun, China

<sup>4</sup>Key Laboratory of Wetland Ecology and Environment, Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun 130012, China

<sup>5</sup>Department of Plant Sciences, College of Agricultural and Marine Sciences, Sultan Qaboos University, Muscat, Oman

<sup>6</sup>Department of Plant Breeding and Genetics, University of Agriculture Faisalabad 38040, Pakistan


<sup>7</sup>Department of Zoology, Government College University Faisalabad 38000, Pakistan

\* Corresponding author: [drnisarahmad@uoj.edu.pk](mailto:drnisarahmad@uoj.edu.pk), xidi888@nenu.edu.cn  
0030-9923/2026/0002-0817 \$ 9.00/0



Copyright 2026 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access  article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).



Supplementary Fig. 1. Maize fields investigated during spring and fall seasons to determine the seasonal and phenological relationship of spider attacking sucking insect pests in maize field of Department of Plant Breeding and Genetics Research Area, University of Agriculture, Faisalabad, Pakistan.