

Antibiofilm Activity of Iron Nanoparticles Synthesised from *Camellia sinensis* against Bacterial and Fungal Pathogens

Tehreem Nasir, Aisha Waheed Qurashi*, Syeda Shazia Bokahri, Uzma Rafi and Roheela Yasmeen

Department of Biology, Lahore Garrison University, Sector C Phase 6 Avenue 4 DHA Lahore, Pakistan

Supplementary Table I. Cultural characteristic of bacterial isolates.

Iso-lates	Culture characteristics				Morphological characteristics				Biochemical characteristics						
	Colony color	Form	Margin	Elevation	Cell shape	G staining	Capsule staining	Spore staining	Catalase test	Methyl red test	Voges proskauer test	Citrate utilization test	Urease test	H ₂ S production test	Motility testing
TBT	OW	Irregular	Undulate	Flat	Rods	+	+	+	+	+	+	+	-	-	+
TBK	OW	Circular	Entire	Raised	Rods	+	+	-	+	+	+	+	-	-	+
TBC	OW	Irregular	Undulate	Flat	Rods	+	+	-	+	+	+	+	-	-	+
TBO	OW	Circular	Entire	Raised	Rods	+	-	-	+	+	+	+	+	-	+
TBA	OW	Circular	Entire	Raised	Rods	+	-	-	+	-	+	+	-	-	+
TBL	OW	Circular	Entire	Raised	Cocci	+	-	-	+	+	+	+	+	-	-
TBR	OW	Irregular	Undulate	Flat	Rods	+	-	-	+	+	+	+	-	-	+


OW, Offwhite; + positive; -.Negative

* Corresponding author: aieshawaheed@yahoo.com,
aishawaheedqureshi@lgu.edu.pk
 0030-9923/2026/0005-2125 \$ 9.00/0



Copyright 2026 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).



Supplementary Fig. 1. Characteristics of iron nanoparticles.