



Research Article

Outbreaks of PPR-FMD among Sheep and Goats in Hail, Saudi Arabia

Ahmed Zein Elabdeen Mahmoud¹, Muaz Magzob Abdellatif^{2*} and Mohamed Abdelsalam Abdalla³

¹Veterinary Laboratories, Ministry of Environment, Water and Agriculture, Saudi Arabia; ²Department of Biology (Microbiology), Faculty of Arts and Science, Northern Borders University, Saudi Arabia; ³Sudan University of Sciences and Technology, Veterinary Medicine and Surgery, Sudan.

Abstract | *Peste des petits ruminants* (PPR) is a viral disease which primarily affects small ruminants, causing significant economic losses for the livestock industry. Clinical investigation was carried out during the suspected outbreaks of PPR/FMD in sheep and goats in Hail, Saudi Arabia. During the outbreaks 271 (20.6%) died out of 648 (49.3%) of affected animals, the overall morbidity in goats (33.3%) was higher as compared to sheep (12.3%), with a case fatality rate in goats reached 74.69% in the 2nd outbreak. Affected animals exhibited fever, stomatitis, mucopurulent to bloody nasal/ocular discharges, watery to bloody diarrhea and lameness. The major gross lesions were erosive stomatitis, hemorrhages in lung, liver, abomasum, small and large intestine. PPRV NP was detected in sheep (n=20) and goats (n=18), anti-PPRV NP was also detected among sheep and goats (n=162) by ELISA, 3ABC-FMD enzyme immunoassay were positive (n=6) when ovine sera were tested. The study revealed that PPR-FMD viruses were circulating in the region and warrants proper control measures.

Editor | Muhammad Abubakar, National Veterinary Laboratories, Park Road, Islamabad, Pakistan.

Received | April 17, 2017; **Accepted** | June 19, 2017; **Published** | June 25, 2017

***Correspondence** | Muaz Magzob Abdellatif, Department of Biology (Microbiology), Faculty of Arts and Science, Northern Borders University, Saudi Arabia; **Email:** muazm20@gmail.com

Citation | Mahmoud, A.Z.E., M.M. Abdellatif and M.A. Abdalla. 2017. Outbreaks of PPR-FMD among sheep and goats in Hail, Saudi Arabia. *Veterinary Sciences: Research and Reviews*, 3(2): 38-44.

DOI | <http://dx.doi.org/10.17582/journal.vsr/2017/3.2.38.44>

Key words: Outbreaks, PPR, FMD, ELISA, Hail, Saudi Arabia

Introduction

Peste des petits ruminants (PPR) is a highly contagious often fatal viral disease of domestic and wild ruminants caused by PPR virus (PPRV) of the family *Paramyxoviridae* (Venkataramanan et al., 2005; Diallo et al., 2007; Banyard et al., 2010). PPR is transmitted by direct contact with infectious animals shedding the virus in both ocular-nasal discharges and in fecal matter (Munir et al., 2013). Fomite contamination with the virus from infected animals such as feed troughs and bedding is an additional source of infection (Lefèvre et al., 1990). The disease is char-

acterized by pyrexia, purulent mucous nasal/ocular discharges, necrotizing and erosive stomatitis, gastroenteritis, diarrhea and bronchopneumonia (Balamurugan et al., 2014). Geographically, PPR has been reported in the Middle East, South Asia, China and sub-Saharan Africa (Shaila et al., 1996), serological evidence has been documented in Uganda, Sudan, Tanzania and Ethiopia (Waret-Szkuta et al., 2008; Luka et al., 2011). In Saudi Arabia, it was first reported in 1990 (Abu Elzein et al., 1990), the disease has since spread to different parts of the country (Housawi et al., 2004; Al-Afaleq et al., 2004; Boshra et al., 2015; Mahmoud et al., 2016, 2017).

Foot and mouth disease (FMD) is one of the most serious transboundary animal diseases. It is a highly contagious viral disease, and may have rapid and unanticipated national and international spread and is characterized by fever and vesicular eruption in the mouth and on the feet and teats (Radostits et al., 2000; Alexandersen, 2005; Wernery and Kinne, 2012). The virus was classified under the genus *aphthovirus* of the family *Picornaviridae* (Radostits et al., 2000). Mild or unapparent infections in sheep and goats may hinder control programs (Kitching and Hughes, 2002). Outbreaks of FMD was repeatedly documented among sheep and goats in various regions of Saudi Arabia (Hafez et al., 1993a; Hafez et al., 1993b; Hamada et al., 2008; Alsayeqh and Fathi, 2012).

The current investigation was aimed at describing the pathology and epidemiology of PPR-FMD infections in four mixed flocks (sheep, goats and camels) in Hail district. Saudi Arabia.

Materials and Methods

Epidemiological investigation

Outbreaks occurred among four mixed farms (sheep, goats and camels) in Hail district. Saudi Arabia were investigated during February to May 2016 (Table 1).

Sample collection

Nasopharyngeal, ocular and rectal swabs (n=46) were

collected from affected goats (n=18), sheep (n=20) and clinically healthy camels (n=8). Sera were collected from goats (n=38), sheep (n=124) and camels (n=8), samples were preserved at 4°C until tested.

P.M Examination

Autopsy were performed on recently dead sheep (n=26) and goats (n=20), gross pathology was recorded.

PPR-NP ELISA

Immune capture ELISA kits for detection of PPR-NP antigen was used (ID Screen®PPR Antigen Capture- CIRAD, Montpellier, France) according to the manufacturer's protocol. The absorbance was read at 450nm with an ELISA reader (BioTek ELX800). The results were calculated as sample positivity (S/P%) and the level of PPRV was expressed as percentage of the positive control. If S/P >20% it was considered negative, S/P ≥20% it was considered positive.

Anti-NP ELISA

The NP-epitopes based competitive ELISA kits (ID screen®PPR competition, ID vet, Grabels, FRANCE) was used for screening of anti-PPRV nucleoprotein antibodies in serum according to the manufacturer's protocol. The absorbance was read at 450nm with an ELISA reader (BioTek ELX800). Samples were presented as S/N%, S/N ≤ 50% - positive, >50 - ≥60%-doubtful and >60% were considered negative.

Table 1: Epidemiological data of outbreaks in Hail, Saudi Arabia

Outbreak (Date)	Species	Vaccination	No. affected (%)	No. dead (%)	Case fatality rate (%)	Total
1st Outbreak (February 2016)	Goats	X	130 (65)	57 (29)	43.8	200
	Sheep	X	150 (50)	37 (12)	24.7	300
	Camels	X	0	0	0	0
	Total		280	94	33.6	500
2nd Outbreak (April 2016)	Goats	X	83 (63.8)	62 (47.7)	74.69	130
	Sheep	√	0	0	0	70
	Camels	X	0	0	0	9
	Total		83	62	24.53	209
3rd Outbreak (April 2016)	Goats	X	35 (70)	7 (14)	20	50
	Sheep	X	150 (50)	47 (16)	31.3	300
	Camels	X	0	0	0	4
	Total		185	54	29.19	354
4th Outbreak (May 2016)	Goats	X	0	0	0	0
	Sheep	X	100 (40)	61 (24.4)	61	250
	Camels	X	0	0	0	0
	Total		100	61	61	250

Table 2: *Morbidity& mortality rates according to age*

	Age group	No. affected (%)	No. dead (%)	Case fatality rate (%)	Total
1 st Outbreak	Young	150 (73.5)	54 (26.5)	36	204
	Adult	130 (76.5)	40 (23.5)	30.8	170
	Total	280 (74.9)	94 (25.1)	33.8	374
2 nd Outbreak	Young	44 (52.4)	40 (47.6)	90.9	84
	Adult	39 (63.9)	22 (36.1)	56.4	61
	Total	83 (57.2)	62 (42.8)	74.7	145
3 rd Outbreak	Young	75 (80.6)	18 (19.4)	24	93
	Adult	110 (75.3)	36 (24.7)	32.7	146
	Total	185 (77.4)	54 (22.6)	29.2	239
4 th Outbreak	Young	100 (62.1)	61 (37.9)	61	161
	Adult	0	0	0	0
	Total	100 (62.1)	61 (37.9)	61	161

Table 3: *Morbidity& mortality rates according to sex*

		Male		Female		Total
		No. affected (%)	No. dead (%)	No. affected (%)	No. dead (%)	
1 st Outbreak	Goats	37 (19.8)	13 (6.9)	93 (49.7)	44 (23.5)	187
	Sheep	23 (12.3)	7 (3.7)	127 (67.9)	30 (16)	187
	Camels	0	0	0	0	0
	Total	60 (16)	20 (5.3)	220 (58.8)	74 (19.9)	374
2 nd Outbreak	Goats	35 (24.1)	24 (16.6)	48 (33.1)	38 (26.2)	145
	Sheep	0	0	0	0	0
	Camels	0	0	0	0	0
	Total	35 (24.1)	24 (16.6)	48 (33.1)	38 (26.2)	145
3 rd Outbreak	Goats	3 (7.1)	0	32 (76.2)	7 (16.7)	42
	Sheep	23 (11.7)	4 (2)	127 (64.5)	43 (21.8)	197
	Camels	0	0	0	0	0
	Total	26 (10.9)	4 (1.7)	159 (66.5)	50 (20.9)	239
4 th Outbreak	Goats	0	0	0	0	0
	Sheep	100 (62.1)	61 (37.9)	0	0	161
	Camels	0	0	0	0	0
	Total	100 (62.1)	61 (37.9)	0	0	161

3ABC-FMD ELISA

Enzyme immunoassay kits (IDEXX FMD 3ABC Ab test. IDEXX Laboratories, Inc. Westbrook, Maine 04092.USA) was used for detection of nonstructural polyprotein (NSP) 3ABC of FMD antigen, it was performed in accordance to the manufacturers' instructions. Samples with percentage values >30% - positive, >20% -negative and samples between 20 and 30% were considered suspicious.

Results

Epidemiological investigation

During the four outbreaks 271 (20.6%) died out of

648 (49.3%) of affected animals, with a case fatality reached 41.8%, no record of vaccination, the overall morbidity in goats (33.3%) was higher as compared to sheep (12.3%) (Figure 1 and Table 1). The disease affected all age groups (Table 2), morbidity and mortality rates were high among female animals (Table 3).

Clinical signs

Affected animals exhibited fever (up to 41°C), loss of appetite, stomatitis, hyper salivation, respiratory distress, mucopurulent to bloody nasal/ocular discharges, watery to bloody diarrhea and lameness (Figure 2).




Figure 1: Morbidity, Mortality and case fatality rates during PPR-FMD outbreaks

Necropsy findings

Dissection of animals revealed erosions on the soft and hard palates, hemorrhages in lung (evidence of

pneumonia), iver, abomasum, small and large intestine (Figure 3).

Sandwich ELISA

The presence of PPRV-antigen was detected in 38 swabs (82.6%) from sheep (n=20) and goats (n=18). Camel swabs were non-reactive.

c-ELISA

Screened sheep and goat sera were positive, camels were seronegative for anti-PPRV nucleoprotein antibodies.

FMD ELISA

Six ovine sera (30%) were positive for anti-NP antibodies while goats were negative.



Figure 2: Infected animals showed A-B. Salivation C-D. dried mucopurulent discharges E-F. erosions on lips, gum, soft and hard palates, tongue and cheeks G. white nodule on gum H. ocular discharges I. reddened eye J-K. watery to bloody diarrhea.




Figure 3: Postmortem examination of infected animals revealed **A.** False membranes; erosions on the soft and hard palates. Hemorrhage in **B-C.** lung (evidence of pneumonia) and liver **D.** Abomasum **E-F.** small & large intestine

Discussion

PPRV is one of the most prevalent infectious diseases among sheep and goats and is considered an emerging economically important disease in Saudi Arabia (Abu-Elzein et al., 1990; Al-Dubaib., 2009; Banyard et al., 2010; Mahmoud et al., 2016, 2017). In the present study, outbreaks occurred in four mixed farms were investigated, PPRV-NP, anti-PPRV-NP were detected from sheep and goats and FMD-NSP were positive from sheep sera. During the outbreaks 271 (20.6%) died out of 648 (49.3%), the overall morbidity in goats (33.3%) was higher as compared to sheep (12.3%), a case fatality rate was high in goats (74.69%), other authors found a mortality of 23 – 100% (Abu Elzein et al., 1990; Taylor et al., 1990; Chowdhury et al., 2014). Mahmoud and Galbat (2017) reported high susceptibility of sheep flocks to the infection with FMD and PPR viruses, with positivity rates of 76% for FMD and 64% for PPR antigen, the susceptibility of sheep and goats can vary with the breed of animal and strain of virus (Kitching and Hughes, 2002). Previous studies indicated that more severe disease results from mixed infection of bacteria and viruses (Osman et al., 2009). We did not take any attempt to isolate bacteria from these cases.

Young animals (6–12 months) were more susceptible than the aged, which are inconsistent with the findings of Taylor et al. (1990). This may be due to maternally derived PPR antibody that persists up to 4 months (Libeau et al., 1992). It is likely that antibody derived from sub-clinical infection in aged goats

protected themselves as well as their young progenies. This observation was in accordance with reports of Taylor et al. (1979).

Foot and mouth disease (FMD) is considered one of the enzootic diseases in Saudi Arabia (Al-Mezaini et al., 1985; Aidros, 2002; Yousef et al., 2012). Sheep infections are frequently mild or in apparent (Geering, 1967; Donaldson and Sellers, 2000), often been implicated as disseminators of the virus (Krystynak, 1987; Barnett and Cox, 1999). Saudi Arabia imports annually millions of live ruminants for slaughter. The majority of these animals are imported from countries where FMD is enzootic, importing either carrier animals which might act as potential source of infection or subclinical infected animals which might actively excrete FMD virus (Hafez et al. 1994).

The present investigation detected mixed PPR-FMD infections among small ruminants in Hail. Saudi Arabia, large-scale sero-prevalence and molecular identification of the circulating strains will help in implementing proper control measures.

Authors' Contribution

Mahmoud, A.Z., M. Abdellatif investigate the clinical picture of the outbreaks, test samples and wrote the manuscript. M. Abdalla supervised the study and revised the manuscript, all read and approved it for submission.

References

- Abu Elzein, E.M.E., Hassanien, M.M., Alafaleq, A.I., Abdelhadi, M.A., Housawi F.M.T. 1990. Isolation of PPR virus from goats in Saudi Arabia. *Vet. Record*, 127: 309-310.
- Al-Dubaib, M.A. 2009. *Peste des petits ruminants* morbillivirus infection in lambs and young goats at Qassim region, Saudi Arabia. *Trop. Anim. Health Prod.*, 41(2): 217-220. <https://doi.org/10.1007/s11250-008-9178-6>
- Alsayeqh, A.F and Fat'hi, S.M. 2012. Recurrent appearance of foot and mouth disease virus (FMDV) in Saudi Arabia. *Sci. J. Microbiol.*, 1(3): 63-70.
- Aidros, H.A. 2002. Regional status and approaches to control and eradication of foot and mouth disease in the Middle East and North Africa. *OIE Sci. Tech. Rev.*, 21: 45-458. <https://doi.org/10.20506/rst.21.3.1348>
- Al-Afaleq, A., Abu-Elzein, E., Al-Naeem, A. and Mahmoud, A. 2004. Serosurveillance for *peste des petits ruminants* (PPR) and rinderpest antibodies in naturally exposed Saudi sheep and goats. *Vet. Arch.*, 74 (6): 459-465.
- Alexandersen, S. and Mowat, N. 2005. Foot-and-mouth disease: Host range and pathogenesis. *Curr. Topics Microbiol. Immunol.*, 288:9-42. https://doi.org/10.1007/3-540-27109-0_2
- Almezaini, S., Sinousi, Y., Chang, S.H., Hawari, and Hafez, S.M. 1985. Some epizootiological aspects of foot and mouth disease in Saudi Arabia. *Proc. Saudi Biol. Soc.*, 8: 267-279.
- Balamurugan, V., Hemadri, D., Gajendragad, M.R., Singh, R.K and Rahman, H. 2014. Diagnosis and control of *peste des petits ruminants*: A comprehensive review. *Virus Dis.* 25(1): 39-56. <https://doi.org/10.1007/s13337-013-0188-2>
- Banyard, A.C., Parida, S., Batten, C., Oura, C., Kwiatek, O and Libeau, G. 2010. Global distribution of *peste des petits ruminants* virus and prospects for improved diagnosis and control. *J. Gen. Virol.*, 91: 2885-2897. <https://doi.org/10.1053/tvj.1998.0338>
- Barnett, P.V and Cox, S.J. 1999. The role of small ruminants in the epidemiology and transmission of foot-and-mouth disease. *Vet. J.*, 158: 6-13.
- Boshra, H., Truong, T., Babiuk, S. and Hemida, M.G. 2015. Seroprevalence of sheep and goat pox, *Peste des petits ruminants* and Rift Valley fever in Saudi Arabia. *PLOS One*. 13:1-12. <https://doi.org/10.1371/journal.pone.0140328>
- Diallo, A., Minet, C., Le Goff, C., Berhe, G., Albina, E., Libeau, G. and Barrett, T. 2007. The threat of *Peste des petits ruminants*: Progress in vaccine development for disease control. *Vaccine*. 25: 5591-5597. <https://doi.org/10.1016/j.vaccine.2007.02.013>
- Donaldson, A. I and Sellers, R. F. 2000. Foot-and-mouth disease. Chapter in *Diseases of Sheep*, 3rd edition. W B Martin and I D Aitken (Editors). Blackwell science, Oxford, 2000.
- Geering, W.A. 1967. Foot and mouth disease in sheep. *Australian Vet. J.*, 43: 485-489. <https://doi.org/10.1111/j.1751-0813.1967.tb04774.x>
- Chowdhury, E.H., Bhuiyan, A.R., Rahman, M.M., Siddique, M.S.A. and Islam, M.R. 2014. Natural peste des petits ruminants virus infection in Black Bengal goats: Virological, pathological and immunohistochemical investigation. *BMC Vet. Res.* 10:263. <https://doi.org/10.1186/s12917-014-0263-y>
- Hafez, S.M., Farag, M.A., Mazloun, K.S. and Al-Bokmy, A.M. 1994. Serological survey of foot and mouth disease in Saudi Arabia. *Revue scientifique et technique (International Office of Epizootics)*. 13(3): 711-9. <https://doi.org/10.20506/rst.13.3.791>
- Hafez, S.M., Farag, M.A., al-Sukayran, A. and al-Mujalli, D.M. 1993a. Epizootiology of foot and mouth disease in Saudi Arabia: I. Analysis of data obtained through district field veterinarians. *Revue scientifique et technique (International Office of Epizootics)*. 12(3): 807-16. <https://doi.org/10.20506/rst.12.3.715>
- Hafez, S.M., Farag, M.A. and Al-Sukayran, A. 1993b. Epizootiology of foot and mouth disease in Saudi Arabia: II. Current status on dairy farms and control measures in operation. *Revue scientifique et technique (International Office of Epizootics)*. 12(3): 817-830. <https://doi.org/10.20506/rst.12.3.720>
- Hamada, K.Z., Habesh, A.R., Saad, M.A. and Aleisa, A.A. 2008. Detection and Characterization of Foot and Mouth Disease Virus, Serotype A in Saudi Arabia in 2005. *Vet. Med. J.*, 56: 2.
- Housawi, F., Abu Elzein, E., Mohamed, G., Gameel, A., Al-Afaleq, A., Hegazi, A. and Al Bishr, B. 2004. Emergence of *peste des petits ruminants* in sheep and goats in Eastern Saudi Arabia. *The Revue d'élevage et médecine vétérinaire des pays tropicaux*. 57 (1-2): 31-34. <https://doi.org/10.1371/journal.pone.0140328>

[org/10.19182/remvt.9901](https://doi.org/10.19182/remvt.9901)

- Kitching, R.P. and Hughes, G.J. 2002. Clinical variation in foot and mouth disease: Sheep and goats. *Revue scientifique et technique* (International Office of Epizootics). 21(3): 505-12. <https://doi.org/10.20506/rst.21.3.1367>
- Krystynak, R.H.E. 1987. Canadas experience with foot-and-mouth disease. *Can. Vet. J.*, 28: 540-542.
- Libeau, G., Diallo, A., Calvez, D. and Lefevre, P.C. 1992. Competitive ELISA using anti- N monoclonal antibodies for specific detection of RP in cattle and ruminants. *Vet. Microbiol.* 31:147-160. [https://doi.org/10.1016/0378-1135\(92\)90073-3](https://doi.org/10.1016/0378-1135(92)90073-3)
- Lefèvre, P. C. and Diallo A. 1990. Peste des petites ruminants. *Revue scientifique et technique*. 9: 951-65.
- Luka, P.D., Erume, J., Mwiine, F.N. and Ayebazibwe, C. 2011. Sero-prevalence of *Peste des petits ruminants* antibodies in sheep and goats after vaccination in Karamoja, Uganda: Implication on Control. *Int. J. Anim. Vet. Adv.*, 3(1): 18-22.
- Mahmoud, A.Z., Abdellatif and Shazali, M. 2016. Prevalence of PPR-virus antibodies in sheep, goats and camels in Hail, Saudi Arabia. *Brit. J. Virol.*, 3(3s): 86-89. <https://doi.org/10.17582/journal.bjv/2016.3.3s.86.89>
- Mahmoud, A.Z., M. Abdellatif and Abdalla, A. 2017. High seroprevalence of PPRV-antibodies among sheep and goats in Hail, Saudi Arabia. *Vet. Sci. Res. Rev.*, 3(1): 1-5. <https://doi.org/10.17582/journal.vsr/2017.3.1.1.5>
- Mahmoud, M.A. and Galbat, S.A. 2017. Outbreak of foot and mouth disease and peste des petits ruminants in sheep flock imported for immediate slaughter in Riyadh. *Vet. World.* 10(2): 238-243. <https://doi.org/10.14202/vetworld.2017.238-243>
- Munir, M., Zohari, S. and Berg, M. 2013. Molecular biology and pathogenesis of *Peste des Petits ruminants virus*. *Springer Briefs Anim. Sci.* 151.
- Osman, N.A., Ali, A.S., Rahman, M.E.A. and Fadol, M.A. 2009. Pathological, serological and virological findings in goats experimentally infected with Sudanese *peste des petits ruminants* virus isolates. *J. Gen. Virol.* 1:1-6.
- Radostits, O.M., Gay, C.C., Blood, D.C. and Hinchkiff, K.W. 2000. *A Text Book of Veterinary Medicine*. 9th Edn, ELBS and Baillier Tindall. PP. 563-618.
- Shaila, M.S., Shamaki, D., Foryth, M.A., Diallo, A., Goatley, L. and Kitching, R.P. 1996. Geographical distribution and epidemiology of PPR viruses. *Virus Res.*, 43:149-53. [https://doi.org/10.1016/0168-1702\(96\)01312-3](https://doi.org/10.1016/0168-1702(96)01312-3)
- Taylor, W.P., Abusaidy, S. and Barret, T. 1990. The epidemiology of PPR in the sultanate of Oman. *Vet. Microbiol.*, 22:341-352. [https://doi.org/10.1016/0378-1135\(90\)90021-M](https://doi.org/10.1016/0378-1135(90)90021-M)
- Taylor, W.P. 1979. Serological studies with the virus of peste des petits ruminants in Nigeria. *Res. Vet. Sci.* 26:236-242.
- Yousef, M.R, Al-Eesa, A.A and Al-Blowi, M.H. Antigenic and genetic analysis of foot and mouth disease virus isolates from 2005 to 2009 in Saudi Arabia for Selection of Candidate vaccine strains. *J. Adv. Vet. Res.*, 2: 73-79.
- Waret-Szkuta, A., Roger, F., Chavernac, D., Yigezu, L., Libeau, G. and Pfeiffer, D.U. 2008. *Peste des Petits Ruminants* (PPR) in Ethiopia: Analysis of a national serological survey. *BMC Vet. Res.*, 4: 34.
- Wernery, U. and Kinne, J. 2012. Foot and mouth disease and similar virus infections in camelids: A review. *Revue scientifique et technique* (International Office of Epizootics). 31(3): 907-918.
- Venkataramanan, R., Bandyopadhyay, S.K. and Oberoi, M.S. Present status and strategies for the control of transboundary and other economically important animal diseases in India: A review. *Indian J. Anim. Sci.*, 75: 456-464. <https://doi.org/10.20506/rst.31.3.2160>