



Review Article

An Overview of Production and Industrial Exploitation of Bacterial Laccases

Fehmina Ashraf^{1*}, Muhammad Irfan², Hafiz Abdullah Shakir¹, Shaukat Ali³, Muhammad Khan¹

¹Department of Zoology, University of the Punjab new campus, Lahore, Pakistan

²Department of Biotechnology, University of Sargodha, Sargodha, Pakistan

³Department of Zoology, Government College University Lahore, Pakistan

Article History

Received: June 13, 2019

Revised: March 26, 2020

Accepted: June 15, 2020

Published: June 29, 2020

Authors' Contributions

FA reviewed the literature and wrote the manuscript. MI supervised the study. HAS helped in manuscript writing. SA did literature survey. MK critically reviewed the final draft.

Keywords

Bacteria, Laccase, Production, Application

Abstract | Laccases (p-diphenol: oxygen oxidoreductase, EC 1.10.3.2) are multi-copper polyphenolic oxidases that oxidize many phenolic and non-phenolic aromatic compounds such as amino or methoxy monophenols, syringaldazine, ABTS and amino or methoxy monophenols. They have been depicted in various species of bacteria, plants and in different genera of fungi. Laccases have been purified by various methods. It involves in di-oxygen to water reduction and 1e⁻ oxidation of phenolic and its allied parts. Laccases are mostly used in different industries like food, paper and pulp, pharmaceutical and textile etc. Both laccases and mediators have been used in different process like delignification of pulp. Laccases from bacterial species are used in dye decolouration and biobleaching processes. This review article describes the whole overview of laccases.

Novelty Statement | This paper reported the updated knowledge about bacterial laccases.

To cite this article: Ashraf, F., Irfan, M., Shakir, H.A., Ali, S. and Khan, M., 2020. An overview of production and industrial exploitation of bacterial laccases. *Punjab Univ. J. Zool.*, 35(1): 147-156. <https://dx.doi.org/10.17582/journal.pujz/2020.35.1.147.156>

Introduction

Laccases (p-diphenol: oxygen oxidoreductase, EC 1.10.3.2) are the top abundant adherents of multi-copper protein family that also encompass monooxygenases, dioxygenases, tyrosinases. Their phylogenetic development is from microbial blue copper proteins azurins to ceruloplasmin, the plasma proteins of eukaryotes (Mayer and Harrel, 1979; Harvey and Walker, 1999; Solomon *et al.*, 1996; Claus, 2003). Yoshida firstly discovered laccase enzyme in *Rhus vernicifera* (Japanese tree) sap in 1883. Whereas, in 1985 laccase was characterized as metal

-containing oxidase by Bertrand (Solomon *et al.*, 1996; Mayer and Harrel, 1979; Harvey and Walker, 1999; Giardina *et al.*, 2010).

Enzymes comprise about 15% to 30% of carbohydrates and 60 to 90 kDa molecular mass. These coppers containing laccase molecules are EC 1.10.3.2 (1,4-benzenediol: oxygen oxidoreductases) and are reported in micro-organisms as well as in higher plants. These are the glycosylated poly-phenols oxidases that hold four Cu²⁺ (copper ions) per molecule and execute one electron (e⁻) of the phenolic oxidation and its allied compound as well as oxygen reduction to water. (Couto and Herrera, 2006; Gianfreda *et al.*, 1999). Mediators (e⁻ shuttle) support the laccase enzyme to oxidize the non-phenolic substrates (Baiocco *et al.*, 2003).

Corresponding author: Fehmina Ashraf

irfan.biotechnologist@gmail.com

Laccase enzymes are described based on having three prosthetic groups of copper categorized on electron paramagnetic resonance ((EPR) signals and light absorbance (Claus, 2003). T1 (Type-1) copper is the site of first oxidation and exhibit absorption band at about 600 nm. The intense electronic absorption via covalent bond of cysteine-copper resulted in blue shade (color) of T1 copper. Whereas, T2 (Type-2) copper and T3 (Type-3) copper jointly form trinuclear center of laccases and appear as a site of dioxygen reduction (Roberts *et al.*, 2003). T2 copper exhibit paramagnetic attributes in EPR but unable to show absorption band in visible range. T3 copper exhibit band at 330nm (Decker and Terwilliger, 2000). Reduction of oxygen molecule resulted in water liberation. Two histidines show linkage with T2 copper and similarly six histidines show linkage with T3 copper. Hydroxyl bridge amid two copper (Cu) atoms of T3 upholds the robust anti-ferromagnetical connection (Kumar *et al.*, 2003) Figure 1.




Figure 1: Scheme of T1 and T2/T3 copper sites of laccase (CotA) from *Bacillus subtilis* (modified from Enguita *et al.*, 2003).

Laccase enzymes show catalytic action on phenolic components of lignin by oxidizing Cα and hence cleaving

Cα- Cβ as well as aryl-alkyl. It is supposed that laccase catalyze the following: firstly, reducing substrate leads T1 copper reduction, secondly; internal electron shifts from T1 Cu to T2 Cu and T3 Cu, Thirdly, at T2 Cu and T3 Cu molecular oxygen reduces to water. Reducing substrate oxidizes by laccase enzyme resulted in loss of solo electron and free unstable radical formation. The radical further undergoes enzymatic (phenol to quinone, oxidation) or non-enzymatic like polymerization, disproportion or hydration reactions (Xu, 1999).




Figure 2: Laccase action on lignin, phenolic sub-units oxidation (adapted from Archibald *et al.*, 1997).

Distribution of laccase enzyme was reported in insects, bacteria, higher plants and fungi. Plants containing turnips, potatoes, apples, pears, cabbages and many other vegetables have laccases. Deuteromycetes, Basidiomycetes and Ascomycetes as well as 60 other fungi have laccase enzyme. (Gianfreda *et al.*, 1999). Laccase based bacterial strains were found such as *Bacillus sp.* HR03, *B. subtilis* SF, *B. halodurans*, *B. subtilis* WP1, *B. pumilus*, *P. desmolyticum* NCIM 2112, *P. putida* and *Azospirillum lipoferum* etc (Narayanan and Murugan, 2014).

Sources

The first microbial laccase was reported in rhizospheric bacterial strain, *Azospirillum lipoferum* accompanying plant root (Givaudan *et al.*, 1993; Sharma *et al.*, 2007; Sharma and Kuhad, 2008). Where its function was found to be melanin pigment formation (Sharma and Kuhad, 2008; Faure *et al.*, 1994, 1995). *Yersinia pestis*, *Pseudomonas maltophilia*, *Xanthomonas campestris* (copA), *Thermus thermophilus* HB27, *Streptomyces psammoticus* MTCC 7334, *S. lavendulae*, *Streptomyces griseus* (epoA), *S. cyaneus*, *Streptomyces antibioticus*, *Pseudomonas fluorescens* GB-1, *Bacillus subtilis*, *Pseudomonas syringae*, *Pseudomonas putida* GB1(cumA), *Caulobacter crescentus*, *Pseudomonas aeruginosa*, *Mycobacterium tuberculosis*, *Pseudomonas aerophilum* (pae1888), *Bordetella compestris*, *Oceanobacillus heyensis* (cotA), *Marinomonas mediterranea*, *Escherichia coli*, *Alpha-proteobacterium* SD21 and *Aquifex aeolicus* are some bacterial strains in which

laccase enzyme has been reported (Alexandre and Zulin, 2000; Enguita *et al.*, 2003; Sharma *et al.*, 2007; Arora and Sharma, 2010). Laccase enzyme that was obtained from *Marinomonas mediterranea*, a marine bacterium had six Cu binding sites but no functional assignment (Amat *et al.*, 2001; Sharma and Kuhad, 2008). Laccase obtained from CotA (constituent of endospore coat) of *Bacillus subtilis* is best studied. CotA takes part in bio-generation of brown coloured spore pigment (Driks, 2004). This pigment act as protective cover against harmful hydrogen peroxide (H_2O_2) and Ultra violet (UV) rays (Martins *et al.*, 2002).

Production

Solid state fermentation (SSF)

The process occurring in near absence or absence of free-flowing water is denoted as solid-state fermentation (SSF). Bacterial laccase synthesis under SSF appeared as economical (Couto and Herrera, 2006). To run SSF about 15% moisture or wetness is indispensable. The following substrates are mostly utilized in SSF such as wheat bran, wood shavings, cereal grains, sawdust and various other materials of animal and plant too (Galzer and Nikaido, 2007). Murugesan *et al.* (2007) studied that in SSF, the conditions for growth of microorganisms are almost compatible to their natural habitation. By employing SSF conditions, phenol oxidase (a laccase-type) was derived from *Streptomyces cyaneus* bacterium and was proposed that enzyme phenol oxidase would be suitable for solubilization as well as mineralization of the lignin contents from wheat straw (Berrocal *et al.*, 2000).

Submerged fermentation (SMF) /liquid fermentation (LF)

Fermentation taking place in excessive water or liquid is denoted as submerged fermentation and it came in strong focus in 1940s (Singhania *et al.*, 2010). By employing SmF, a significant amount of laccase can be achieved because of having set configurations of bioreactor and ease in controlling parameters (Thiruchelvam and Ramsay, 2007). The SmF established procedures that have been optimized and cultivated for decades for enzymes synthesis by bacterial strains or others cannot be substituted by SSF (Hölker, 2005). According to Téllez-Jurado *et al.* (2005) liquid batch culture in SmF presented more growth and synthesis of laccase than that of SSF. Agricultural wastes appeared as cheap substrates for laccase synthesis and acted as ligninolytic enzymes inducers because of having cellulose, hemicellulose and lignin contents. Wheat bran has been considered the most suitable substrate for laccase enzyme synthesis with SmF and SSF as well (Papinutti *et al.*, 2003; Marques de Souza *et al.*, 2002).

Factors affecting laccase production

Influence of nitrogen and carbon source

An ideal and defined growth medium for organism comprises 0.1% and 1% (w/v) of yeast extract (nitrogen source) and many other sources (carbon and nitrogen)

respectively. Sources of carbon such as mannose, fructose, lactose, maltose and glucose are usually used. Superfluous concentration of sucrose as well as glucose act as inhibitors for initiation thus lessens the laccase synthesis. Polymeric substrates just like cellulose are capable enough to tackle such problems (Lee *et al.*, 2004). Peptone, $(NH_4)_2SO_4$, $NaNO_3$, yeast extract and urea are sources of nitrogen and are usually used. However, exhaustion of nitrogen leads to trigger of production of laccase (Keyser *et al.*, 1978).

Influence of incubation time and temperature

On laccase synthesis, there is a narrow effect of temperature but optimum or ideal temperature changes from strain to strain. Optimal temperature for laccase synthesis changes with correspondence to the presence of dark (30°C) and light (25°C) (Thurston, 1994; Pointing *et al.*, 2000). Activity of laccase enzyme significantly augmented by pre-incubating at the temperatures of 40°C and then 50°C (Farnet *et al.*, 2000). According to Palmieri *et al.* (1993), an unchanged in activity was observed even after such a long incubation period beyond 4 hours at 40 °C. The spore-coat of strain *Bacillus subtilis* has CotA component that is a gene product. CotA is a bacterial laccase is best studied enzyme (Hullo *et al.*, 2001). The exception of CotA is its thermo-stability with half-life of 2 hours at 80°C and 75°C optimal temperature (Martins *et al.*, 2002).

Influence of pH

Thurston (1994) observed the limited or narrow effect of pH on laccase synthesis. However, reactions of substrate for laccase enzyme cause effect on variation of optimal pH. Reactions of substrate for laccase enzyme cause effect on variation of optimal pH. According to several reports, laccase activity exhibits its profile as bell-shaped. At optimal pH, the enhancement in substrate oxidation occurs because of potency difference amid T1 Cu and phenolic substrate whereas, hydroxyl anion (OH^-) makes bond to T2/T3 Cu centre of laccase (Kunamneni *et al.*, 2007). For syringaldazine substrate, the activity of enzyme was observed in 3.0–8.0 pH range. Optimal pH 4.0 and 5.0 was determined for L1 (laccase isozyme) and L2 respectively (Cordi *et al.*, 2007).

Influence of inducer

Inducers such as guaiacol, ferulic acid, veratryl alcohol, ethanol and gallic acid enhance the synthesis of laccase enzyme. On attaining the growth phase, synthesis of laccase enzyme in γ -proteobacterium JB augmented 13 times after adding $CuSO_4$ to media. Malachite green, Ethidium bromide, Thymol blue and phenol red have also stimulated the synthesis of laccase by 19, 17, 2 and 4 folds respectively (Kanam *et al.*, 2004). Alcohol boosted laccase activity comparatively to xylinine. Boost of laccase by alcohol appeared costly method (Lee *et al.*, 1999).

Influence of metal ions

From alkaline medium, the obtainable laccase lessens 50 times than that of cultivation media having low copper ion concentration (Palmieri *et al.*, 2000; Assavanig *et al.*, 1992). Zhang *et al.* (2010) studied that EDTA, Mg^{2+} , Cu^{2+} and Zn^{2+} at 6.25–50mM proved to be un-affecting agents for the activity of laccase enzyme. However, a decline in laccase activity was observed due to Fe^{2+} and Ca^{2+} ions at 6.25–50mM and 25–50mM concentrations respectively. Molecular and Biochemical approaches were employed for identification of soil bacterial strain *Stenotrophomonas maltophilia* AAP56. Without any substrate, the influence of Zn^{2+} , Mn^{2+} , Cu^{2+} , Mg^{2+} , Ca^{2+} , Fe^{2+} , urea, EDTA and sodium azide by 5 minutes incubation period at 4°C was studied. A significant increase in enzyme synthesis about 2.6 folds with 275UL-1 activity of laccase enzyme was noted Galai *et al.* (2009).

Purification of laccase

Enzymes purification employing ammonium sulphate has been carried out for several years. However, protein precipitation via buffer/desalt exchange of protein, ammonium sulphate, gel filtration and anion-exchange chromatography proved to be more effectual methods (Grotewold *et al.*, 1998). Purification of bacterial laccase derived from soil bacterial strain *Pseudomonas putida* F6 had done by employing the coalition of gel filtration and anion-exchange chromatography and consequently observed an escalation in activity of laccase enzyme by 518 and 747 Umg^{-1} experimented by McMahon *et al.*, 2007. Partial purification of laccase derived from cell of *Streptomyces psammoticus* strain was done by applying an ordinary method ammonium sulphate for precipitation whereas, copper and calcium alginate beads were applied for immobilization of laccase in entrapment method. However, alginate beads using calcium-based laccase immobilization hold just 42.5% of the activity of laccase whereas in contrast, alginate bead utilizing copper substantiated better for immobilization of laccase by holding 61% of the laccase activity (Niladevi and Prema, 2008). Purification of bacterial laccase from a strain *Azospirillum lipoferum* was carried out by dialysis method in which ammonium sulphate was expended for precipitation of protein from supernatant (Diamantidis *et al.*, 2000). Purified bacterial laccase derived from strain *Streptomyces lavendulae* REN-7 exhibited 73kDa molecular mass and solo band of protein after applying 10% of SDS-PAGE (Suzuki *et al.*, 2003).

Catalytic activity

Catalytic activity of laccase enzyme upswings by the grouping of laccase enzyme and mediators. These mediators are very tiny (little molecular weight) and act as electron carrier amid laccase enzyme and substrate (Duran *et al.*, 2002; Galli and Gentili, 2004) Laccase enzymes have been widely used to oxidize or catalyze substrates

for many years. The following phenolic and non-phenolic substrates have been oxidized or catalyzed are methoxy- or amino-monophenols, and non-phenols such as ABTS, hydroxyindoles, syringaldazine, aromatic diamines etc. (Mayer, 1987; Cai *et al.*, 1993). The efficiency of laccase along with ABTS rises many folds, so it oxidizes laccase substrates and non-laccase substrates both (Bourbonnais and Paice, 1992). Bacterial laccase with broader pH appears more effective as compared to other laccases like fungal laccase when employed industrially such as bio-bleaching of textiles stuffs, paper pulp processing etc. (Singh *et al.*, 2009b). Singh *et al.* (2007) and Ye *et al.* (2010) reported the activity and stability of bacterial laccases at extensive pH range probably 6.0 to 8.5. As the laccase derivative of Metagenomec worked best at 9.0 pH for a non-phenolic substrate “syringaldazine” (Ye *et al.*, 2010). Singh *et al.* (2007) and Ye *et al.* (2010) reported the activity and stability of bacterial laccases at extensive pH range probably 6.0 to 8.5. As laccase derivative of Metagenomec worked best at 9.0 pH with a non-phenolic substrate “syringaldazine” Ye *et al.* (2010). Whereas, *Bacillus halodurans* C-125 derived alkaline bacterial laccase Lbh1 activity at optimum pH range 7.5–8.0 was observed with syringaldazine as confirmed by Ruijsenaars and Hartmans, 2004. Bacterial strain c-proteobacterium JB laccase showed 70% stability at both acidic 3.0 pH and alkaline 10.6 pH and 80% stability at broad 4.0 to 9.0 pH range with 48 hours incubation period at normal 37°C temperature. The absence of protease production may be a reason of stability (Singh *et al.*, 2007).

Applications

Laccases are used biotechnologically because they have capability to oxidize a variety of phenolic compounds as well as non-phenolic compounds (Figure 3) (Mohammadian *et al.*, 2010). Laccases are exploited in the following industries including textile, petrochemical, paper, pulp, food, wood, pharmaceutical (medicines) etc. (Arora and Sharma, 2010). Couto and Herrera (2006) found that Laccases are not only confine to water distillation systems but they are also being involved in anticancer medicines (drugs) preparation as well as reducing the harmfulness of cosmetics. Now researchers are trying to work for the enzymatically organic compounds production and utility of laccases for the development of biosensor, biotransformation and bio-oxidation. (Arora and Sharma, 2010).

Pulp and paper industry

Formation of paper at industrial level is acquired by following conventional method by using chemical oxidants having oxygen or chlorine for pre-treatment e.g. separation of lignin and delignification (Luisa *et al.*, 1996). In last few years, industrially the traditional and polluting methods of pre-treatment i.e. chlorine-based chemical methods have changed into oxygen-based chemical methods.

However, use of laccase is safer method for pre-treatment or delignification of wood pulp along with protection of cellulose integrity. (Barreca *et al.*, 2003; Gamelas *et al.*, 2005; Shi, 2006). Mediators are the chief chemical agents that support laccases in the process of delignification of wood pulp (Bourbonnais *et al.*, 1997). Laccases cannot oxidize lignin directly. Mediator (e^- shuttle) plays an important and intermediate role in delignification. The oxidation of mediator is done by laccases whereas it (mediator) then oxidizes lignin in wood (Bourbonnais *et al.*, 1997; Svendsen and Xu, 2001). This indirect delignification is supposed to be impossible because of huge structure of laccase and its access towards lignin present in plant cell (Sharma *et al.*, 2007; Kandioller and Christov, 2001; Singh *et al.*, 2008). Laccases obtaining from two bacterial strains *P. stutzeri* (soil bacterium) (Kumar *et al.*, 2005) and *S. cyaneus* (Arias *et al.*, 2003) along with mediators HOBt and ABTS have been used in the process of bio-bleaching of eucalyptus kraft pulps (Held *et al.*, 2005). Laccases have competency to functionalize and cross-link the lignocellulosic fibers (Call, 2005; Guebitz and Cavaco, 2003; Xu *et al.*, 2006).

to decolourize wastewater as well as bleaching of textiles and artificial dyes (Setti *et al.*, 1999). Microbial laccases from bacterial strains *Stenotrophomonas maltophilia* and *Streptomyces psammoticus* are competent enough to decolorize numerous synthetic dyes (Romero *et al.*, 2006; Niladevi *et al.*, 2007). Paszczynski *et al.* (1991) has worked on the improvement of degradation of azo dyes by the application of *Streptomyces* (bacterial sp.). These azo dyes are approximately half of total synthetic dyes (Selvam and Swaminathan, 2003).

Food industry

In food processing industry, laccases are being employed in different areas i.e. juice processing, wine stabilization, baking as well as bioremediation of industrial effluents because of its capability to remove unwanted phenolic compounds (Couto and Herrera, 2006). Yague *et al.* (2000) found that the effluents of beer factory are somehow hazardous for environment because of its black colour (dark brown) and holding polyphenolic compounds. Laccase plays role in removing these phenols and bond between enzyme-substrate by using membrane filter method (Minussi, 2002). Phenolic contents and oxidative products already existing in juices provide flavor or taste, aroma and colour to it. Enzymatic darkening is a condition in which polymerization and phenolic as well as polyphenolic oxidation cause change in odor and colour of juice. So, in order to keep juices in original colour and taste or high quality, laccases are used in food processing industries (Ribeiro *et al.*, 2010). Laccases are used in bakery products too. Laccases are supplemented to dough to enhance or improve gluten structure of dough, product volume, crumb structure and baked products softness (Minussi *et al.*, 2002).

Immobilization of laccases

Glass ceramic was taken as decolourizing agent in textile industry after modified by chemicals aminopropyltriethoxysilane/ glutaraldehyde as well as carbodiimide/glutaraldehyde. Besides this montmorillonite was also used to decolourize textiles after modification by aminopropyltriethoxysilane/ glutaraldehyde chemicals (Peralta-Zamora *et al.*, 2003). Laccases immobilized on various substrates i.e. ceramics support and the best one pyrolytic graphite (Minussi *et al.*, 2007). Alumina pellet substrates were employed for immobilization of laccase-based spores of bacterial strain "Bacillus SF". Immobilized spores as well as free spores had ability to decolourize the common dyes used in textile industry like Acid Blue 74, Mordant Black 9, Mordant Brown 15, Mordant Brown 96. The immobilization took about 90 minutes. However, at 60°C half-life increased by 66-80 hours after immobilization of laccase-based "Bacillus SF" on alumina pellet substrates (Held *et al.*, 2005).




Figure 3: Scheme of applications of laccase (adapted from Morozova *et al.*, 2007).

Textile industry

Artificial dyes are being used extensively in industries, most probably in textile, food, paper printing, leather and cosmetic industries for coloration purposes (Forgacs *et al.*, 2004). Tavares *et al.* (2009) found that dyes are actually dyed or shaded molecules and give colour to cellulose fibers. These dyes become responsible for the assembly of large amount of stained effluent. As a result, the synthetic colours cause hindrance in biodegrading process (Wesenberg *et al.*, 2003; Moilanen *et al.*, 2010). Problem in using artificial dyes is its carcinogenicity because of having aromatic compounds (Baughman and Perenich, 1988). Laccases exploitation is rising rapidly,

Pharmaceutical industry

In pharmaceutical or medicine industry, laccase molecules have been employed in generation of detoxifying agent and antimicrobial agent. Anti-inflammatory, sedatives, anesthetics and antibiotics are also synthesized with the use of laccases (Arora and Sharma, 2010; Nicotra *et al.*, 2004; Haught *et al.*, 2001; Juelich *et al.*, 2001; Johansen, 1996). By the expense of laccases, anticancer drug “actinocin” has been made from a chemical 4-methyl-3-hydroxyanthranilic acid. The drug “actinocin” cures the cancer by causing hindrance in DNA transcription from tumor (cancer) cell (Burton, 2003). The oxidative conversion of both aromatic compounds and aliphatic amines into acid “3-(3, 4-dihydroxyphenyl)-propionic” can be done by laccases and the resultants act as anti-viral agents (Ncanana *et al.*, 2007). Laccases also show inhibitory action for the reverse transcriptase of HIV-1 reported by Wang and Ng (2004). Roggen *et al.* (2001) found another function of laccase that is to lessen the allergenicity.

Conclusion

Laccases are not only present in bacteria but also in plants and fungi. Laccases of bacteria (bacterial Laccases) are enzymes secreted by different bacterial strains such as *Streptomyces sp.*, *Pseudomonas sp.*, *Bacillus sp.* etc. These catalyze redox reactions. Bacterial laccases have different biotechnological applications like food industry, textile industry, dye decolouration, pharmaceuticals. These are also involved in biodegradation/ detoxification the industrial effluents as well as in bioremediation. In paper and pulp industry it is used as biobleaching agent.

Conflict of interest

The authors have declared no conflict of interest.

References

- Alexandre, G., and Zhulin, I.B., 2000. Laccases are wide spread in bacteria. *Trends Biotech.* 2000; **18**: 41– 42. [https://doi.org/10.1016/S0167-7799\(99\)01406-7](https://doi.org/10.1016/S0167-7799(99)01406-7)
- Amat, A.S., Elio, P.L., Fernandez, E., Borron, J.C.G. and Solano, F., 2001. Molecular cloning and functional characterization of a unique multipotent polyphenol oxidase from *Marinomonas mediterranea*. *Biochem. Biophys. Acta*, 1547: 104–116. [https://doi.org/10.1016/S0167-4838\(01\)00174-1](https://doi.org/10.1016/S0167-4838(01)00174-1)
- Archibald, F.S., Bourbonnais, R., Jurasek, L., Paice, M.G., and Reid, I.D., 1997. Kraft pulp bleaching and delignification by *Trametes versicolor*. *J. Biotechnol.*, **53**: 215–336. [https://doi.org/10.1016/S0168-1656\(97\)01675-1](https://doi.org/10.1016/S0168-1656(97)01675-1)
- Arias, M.E., Arenas, M., Rodríguez, J., Soliveri, J., Ball, S.A. and Hernández, M., 2003. Kraft pulp biobleaching and mediated oxidation of a nonphenolic substrate by laccase from *Streptomyces cyaneus* CECT 3335. *Appl. Environ. Microbiol.*, **69**: 1953–1958. <https://doi.org/10.1128/AEM.69.4.1953-1958.2003>
- Arora, D.S. and Sharma, R.K., 2010. Ligninolytic Fungal Laccases and their biotechnological applications. *Appl. Biochem. Biotechnol.*, **160**: 1760–1788. <https://doi.org/10.1007/s12010-009-8676-y>
- Assavanig, A., Amornkitticharoen, B., Ekpaial, N., Meevootisom, V., Flegel, T.W., 1992. Isolation, characterization and function of laccase from *Trichoderma*. *Appl. Microbiol. Biotechnol.*, **38**: 198–202. <https://doi.org/10.1007/BF00174468>
- Baiocco, P., Barreca, A.M., Fabbrini, M., Galli, C. and Gentili, P., 2003. Promoting laccase activity towards non-phenolic substrates: a mechanistic investigation with some laccase-mediator systems. *Org. Biomol. Chem.*, **1**: 191–197. <https://doi.org/10.1039/B208951C>
- Barreca, A.M., Fabbrini, M., Galli, C., Gentili, P. and Ljunggren, S., 2003. Laccase/mediated oxidation of a lignin model for improved delignification procedures. *J. Mol. Catal. B-Enzym.*, **26**: 105–110. <https://doi.org/10.1016/j.molcatb.2003.08.001>
- Baughman, G.L., Perenich, T.A., 1988. Fate of dyes in aquatic systems: I solubility and partitioning of some hydrophobic dyes and related compounds. *Environ. Toxicol. Chem.*, **7**: 183–199. <https://doi.org/10.1002/etc.5620070302>
- Berrocal, M., Ball, A.S., Huerta, S. and Arias, J.M., 2000. Biological upgrading of wheat straw through solid-state fermentation with *Streptomyces cyaneus*. *Appl. Microbiol. Biotechnol.*, **54**: 764–771. <https://doi.org/10.1007/s002530000454>
- Bourbonnais, R., Paice, M.G., Freiermuth, B., Bodie, E. and Borneman, S., 1997. Reactivities of various mediators and laccases with kraft pulp and lignin model compounds. *Appl. Environ. Microbiol.*, **12**: 4627–4632. <https://doi.org/10.1128/AEM.63.12.4627-4632.1997>
- Bourbonnais, R. and Paice, M.G., 1996. Enzymatic delignification of Kraft pulp using laccase and a mediator. *J. Technol. Assoc. Pap. Pulp. Ind.*, **79**: 199–204.
- Bourbonnais, R.E. and Paice, M.G., 1992. Demethylation and delignification of kraft pulp by *Trametes versicolor* laccase in the presence of 2,2'-azinobis (3-ethylbenzthiazoline-6-sulphonate). *Appl. Microbiol. Biotechnol.*, **36**: 823–827. <https://doi.org/10.1007/BF00172202>
- Burton, S., 2003. Laccases and phenol oxidases in organic synthesis. *Curr. Org. Chem.*, **7**: 1317–1331. <https://doi.org/10.2174/1385272033486477>
- Cai, W., Martin, R., Lemaure, B., Leuba, J.L. and Petiard, V., 1993. Hydroxy-indoles: a new class of laccase substrates. *Pl. Physiol. Biochem.*, **31**: 441–445.
- Call, H.P., 2005. WO2005103372 A2.
- Cha, J., Cooksey, D.A., 1991. Copper resistance

- in *Pseudomonas syringae* by periplasmic and outer membrane proteins. *Proc. Natl. Acad. Sci. USA.*, **88**: 8915–8919. <https://doi.org/10.1073/pnas.88.20.8915>
- Claus, H. and Filip, Z., 1997. The evidence of a laccase-like activity in a *Bacillus sphaericus* strain. *Microbiol. Res.*, **152**: 209–215. [https://doi.org/10.1016/S0944-5013\(97\)80014-6](https://doi.org/10.1016/S0944-5013(97)80014-6)
- Claus, H., 2003. Laccases and their occurrence in prokaryotes. *Arch. Microbiol.*, **179**: 145–150. <https://doi.org/10.1007/s00203-002-0510-7>
- Cordi, L., Minussi, R.C., Freire, R.S. and Dur'an, N., 2007. Fungal laccase: copper induction, semi-purification, immobilization, phenolic effluent treatment and electrochemical measurement. *Afr. J. Biotechnol.*, **6**: 1255–1259.
- Couto, S.R. and Toca-Herrera, J.L., 2006. Industrial and biotechnological applications of laccases: A review. *Biotechnol. Adva.*, **24**: 500–513. <https://doi.org/10.1016/j.biotechadv.2006.04.003>
- Decker, H. and Terwilliger, N., 2000. Cops and robbers: putative evolution of copper oxygen binding proteins. *J. Exp. Biol.*, **203**: 1777–1782.
- Diamantidis, G., Effosse, A., Potier, P. and Bally, R., 2000. Purification and characterization of first bacterial laccase in the rhizospheric bacterium *Azospirillum lipoferum*. *Soil Biol. Biochem.*, **32**: 919–927. [https://doi.org/10.1016/S0038-0717\(99\)00221-7](https://doi.org/10.1016/S0038-0717(99)00221-7)
- Dom'inguez, A., Couto, S.R. and Sanrom'an, M.A., 2005. Dye decolorization by *Trametes hirsuta* immobilized into alginate beads. *World J. Microbiol. Biotechnol.*, **21**: 405–409. <https://doi.org/10.1007/s11274-004-1763-x>
- Driks, A., 2004. The *Bacillus subtilis* spore coat. *Phytopathology*, **94**: 1249–1251. <https://doi.org/10.1094/PHYTO.2004.94.11.1249>
- Duran, N., Rosa, M.A., Annibale, A. and Gianfreda, L., 2002. Applications of laccases and tyrosinases (phenoloxidases) immobilized on different supports: A review. *Enzyme Microb. Technol.*, **31**: 907–931. [https://doi.org/10.1016/S0141-0229\(02\)00214-4](https://doi.org/10.1016/S0141-0229(02)00214-4)
- Enguita, F.J., Martins, L.O., Henriques, A.O. and Carrondo, M.A., 2003. Crystal structure of a bacterial endospore coat component. A laccase with enhanced thermostability properties. *J. Biol. Chem.*, **278**: 19416–19425. <https://doi.org/10.1074/jbc.M301251200>
- Farnet, A.M., Criquet, S., Tagger, S., Gil, G. and Le Petit, J., 2000. Purification, partial characterization, and reactivity with aromatic compounds of two laccases from *Marasmius quercophilus* strain 17. *Can. J. Microbiol.*, **46**: 189–194. <https://doi.org/10.1139/cjm-46-3-189>
- Faure, D., Bouillant, M. and Bally, R., 1995. Comparative study of substrates and inhibitors of *Azospirillum lipoferum* and *Pyricularia oryzae* laccases. *Appl. Environ. Microbiol.*, **61**: 1144–1146. <https://doi.org/10.1128/AEM.61.3.1144-1146.1995>
- Faure, D., Bouillant, M.L. and Bally, R., 1994. Isolation of *Azospirillum lipoferum* 4T Tn5 mutants affected in melanization and laccase activity. *Appl. Environ. Microbiol.*, **60**: 3413–3415. <https://doi.org/10.1128/AEM.60.9.3413-3415.1994>
- Forgacs, E., Cserhatia, T. and Oros, G., 2004. Removal of synthetic dyes from wastewaters: A review. *Environ. Int.*, **30**: 953–971. <https://doi.org/10.1016/j.envint.2004.02.001>
- Francis, C.A. and Tebo, B.M., 2001. CumA multicopper oxidase genes from diverse Mn(II)-oxidizing and non-Mn(II)-oxidizing *Pseudomonas* strains. *Appl. Environ. Microbiol.*, **67**: 4272–4278. <https://doi.org/10.1128/AEM.67.9.4272-4278.2001>
- Francis, C.A. and Tebo, B.M., 2002. Enzymatic manganese (II) oxidation by metabolically dormant spores of diverse *Bacillus* species. *Appl. Environ. Microbiol.*, **68**: 874–880. <https://doi.org/10.1128/AEM.68.2.874-880.2002>
- Galai, S., Limam, F. and Marzouki, M., 2009. A new *Stenotrophomonas maltophilia* strain producing laccase, use in decolourization of synthetic dyes. *Appl. Biochem. Biotech.*, **158**: 416–431. <https://doi.org/10.1007/s12010-008-8369-y>
- Galli, C. and Gentili, P., 2004. Chemical messengers: mediated oxidations with the enzyme laccase. *J. Phys. Org. Chem.*, **17**: 973–977. <https://doi.org/10.1002/poc.812>
- Galzer, A.N. and Nikaido, H., 2007. *Microbial Biotechnology: Fundamentals of Applied Microbiology*. 2nd Edn., Cambridge, UK. ISBN-13: 9780521842105, pp. 554. <https://doi.org/10.1017/CBO9780511811227>
- Gamelas, J.A.F., Tavares, A.P.M., Evtuguin, D.V. and Xavier, A.M.B., 2005. Oxygen bleaching of kraft pulp with polyoxometalates and laccase applying a novel multi-stage process. *J. Mol. Catal. B-Enzym.*, **33**: 57–64. <https://doi.org/10.1016/j.molcatb.2005.03.001>
- Gianfreda, L., Xu, F., Bollag, J.M., 1999. Laccases: A useful group of oxidoreductive enzymes. *Biorem. J.*, **3**: 1–25. <https://doi.org/10.1080/10889869991219163>
- Giardina, P., Faraco, V., Pezzella, C., Piscitelli, A., Vanhulle, S. and Sannia, G., 2010. Laccases: A never-ending story. *Cell. Mol. Life Sci.*, **67**: 369–385. <https://doi.org/10.1007/s00018-009-0169-1>
- Givaudan, A., Effosse, A., Faure, D., Potier, P., Bouillant, M.L. and Bally, R., 1993. Polyphenol oxidase in *Azospirillum lipoferum* isolated from rice rhizosphere: Evidence for laccase activity in nonmotile strains of *Azospirillum lipoferum*. *FEMS Microbiol. Lett.*, **108**: 205–210. <https://doi.org/10.1111/j.1574-6968.1993.tb06100.x>
- Grotewold, E., Taccioli, G.E., Aisemberg, G.O. and

- Judewicz, N.D., 1998. Purification of an extracellular fungal laccase. *Mircen J. Appl. Microbiol. Biotechnol.*, **4**: 357–363. <https://doi.org/10.1007/BF01096141>
- Guebitz, G.M., Cavaco and Paulo, A., 2003. New substrates for reliable enzymes: enzymatic modification of polymers. *Curr. Opin. Biotechnol.*, **14**: 577–582. <https://doi.org/10.1016/j.copbio.2003.09.010>
- Harvey, B.M. and Walker, J.R.K., 1999. Studies with plant laccases: I. Comparison of plant and fungal laccases. *Biochem. Mol. Biol. Biophys.*, **3**: 45–51.
- Haight, J.C., Miracle, G.S. and Convents, A.C., 2001. WO2001060157 A2.
- Held, C., Kandelbauer, A., Schroeder, M., Cavaco-Paulo, A. and Guebitz, G.M., 2005. Biotransformation of phenolics with laccase containing bacterial spores. *Environ. Chem. Lett.*, **3**: 74–77. <https://doi.org/10.1007/s10311-005-0006-1>
- Hölker, U. and Lenz, J., 2005. Solid-state fermentation are there any biotechnological advantages? *Curr. Opin. Microbiol.*, **8**: 301. <https://doi.org/10.1016/j.mib.2005.04.006>
- Hou, H., Zhou, J., Wang, J., Du, C. and Yan, B., 2004. Enhancement of laccase production by *Pleurotus ostreatus* and its use for the decolorization of anthraquinone dye. *Process. Biochem.*, **39**: 1415–1419. [https://doi.org/10.1016/S0032-9592\(03\)00267-X](https://doi.org/10.1016/S0032-9592(03)00267-X)
- Hullo, M.F., Moszer, I., Danchin, A. and Martin-Verstraete, I., 2001. CotA of *Bacillus subtilis* is a copper-dependent laccase. *J. Bacteriol.*, **183**: 5426–5430. <https://doi.org/10.1128/JB.183.18.5426-5430.2001>
- Isono, Y. and Hoshino, M., 1989. Laccase-like activity of nucleoside oxidase in the presence of nucleosides. *Agric. Biol. Chem.*, **53**: 2197–2203. <https://doi.org/10.1271/bbb1961.53.2197>
- Johansen, C., 1996. WO9606532 A1.
- Juelich, W.-D., Schauer, F., Lindequist, U., Hammer, E., Schaefer, A. and Jonas, U., 2001. WO2001098518 A2.
- Kanam, M., Prince, S. and Neena, C., 2004. Copper and dyes enhance laccase production in γ -proteobacterium JB. *Bioethanol. Lett.*, **26**: 1047–1050. <https://doi.org/10.1023/B:BILE.0000032959.10370.18>
- Kandioller, G. and Christov, L., 2001. Evaluation of the delignification and bleaching abilities of selected laccases with HBT on different pulps. In: Argyropoulos DS (ed) *Oxidative delignification chemistry fundamentals and catalysis*. ACS symposium series, 750. *Oxford Univ. Press, USA*, pp. 427–443. <https://doi.org/10.1021/bk-2001-0785.ch027>
- Keyser, P., Kirk, T.K. and Zeikus, J.G., 1978. Ligninolytic enzyme system of *Phanerochaete chrysosporium*: synthesized in the absence of lignin in response to nitrogen starvation. *J. Bacteriol.*, **135**: 790–797. <https://doi.org/10.1128/JB.135.3.790-797.1978>
- Kumar, A., Vanamala, A. and Kumar, R., 2005. Exploration of bacterial laccase in *Pseudomonas stutzeri* and its application in bleaching the wood pulp. *FEBS J.*, **272**(s1): N6–008P.
- Kumar, S.V.S., Phale, P.S., Durani, S. and Wangikar, P.P., 2003. Combined sequence and structure analysis of the fungal laccase family. *Biotechnol. Bioeng.*, **83**: 386–394. <https://doi.org/10.1002/bit.10681>
- Kunamneni, A., Ballesteros, A., Plou, F.J. and Alcalde, M., 2007. Fungal laccase—a versatile enzyme for biotechnological applications, in communicating current research and educational topics and trends in applied microbiology. 2007; A. Mendez-Vilas, Ed., Formex, Badajoz, Spain. **1**: 233–245.
- Lee, I.Y., Jung, K.H., Lee, C.H. and Park, Y.H., 1999. Enhanced production of laccase in *Trametes versicolor* by the addition of ethanol. *Biotechnol. Lett.*, **21**: 965–968. <https://doi.org/10.1023/A:1005646812381>
- Lee, K.H., Wi, S.G., Singh, A.P. and Kim, Y.S., 2004. Micromorphological characteristics of decayed wood and laccase produced by the brown-rot fungus *Coniophora puteana*. *J. Wood Sci.*, **50**: 281–284. <https://doi.org/10.1007/s10086-003-0558-2>
- Lee, Y., Henderson, M., Panopoulos, N.J. and Schroth, M.N., 1994. Molecular cloning, chromosomal mapping, and sequence analysis of copper resistance genes from *Xanthomonas campestris* pv. juglandis: homology with small copper proteins and multicopper oxidases. *J. Bacteriol.*, **176**: 173–188. <https://doi.org/10.1128/JB.176.1.173-188.1994>
- Luisa, M., Goncalves, F.C. and Steiner, W., 1996. Purification and characterization of laccase from a newly isolated wood-decaying fungus. In: *Enzymes for pulp and paper processing*. Eds.: Jeffries TW, Viikari IL. *Am. Chem. Soc. Washington, USA*, pp. 258–266. <https://doi.org/10.1021/bk-1996-0655.ch020>
- Marques de Souza, C.G., Zilly, A. and Peralta, R.M., 2002. Production of laccase as the sole phenoloxidase by a Brazilian strain of *Pleurotus pulmonarius* in solid state fermentation. *J. Basic Microb.*, **42**: 83. [https://doi.org/10.1002/1521-4028\(200205\)42:2<83::AID-JOBM83>3.0.CO;2-Z](https://doi.org/10.1002/1521-4028(200205)42:2<83::AID-JOBM83>3.0.CO;2-Z)
- Martins, L.O., Soares, C.M., Pereira, M.M., Teixeira, M., Costa, T., Jones, G.H. and Henriques, A.O., 2002. Molecular and biochemical characterization of a highly stable bacterial laccase that occurs as a structural component of the *Bacillus subtilis* endospore coat. *J. Biol. Chem.*, **277**: 18849–18859. <https://doi.org/10.1074/jbc.M200827200>
- Mayer, A.M. and Harel, E., 1979. Polyphenol oxidases in plants. *Phytochemistry*, **18**: 193–215. [https://doi.org/10.1016/0031-9422\(79\)80057-6](https://doi.org/10.1016/0031-9422(79)80057-6)
- Mayer, A.M., 1987. Polyphenol oxidases in plants—recent progress. *Phytochemistry*, **26**: 11–20. [https://doi.org/10.1016/S0031-9422\(00\)81472-7](https://doi.org/10.1016/S0031-9422(00)81472-7)
- McMahon, A.M., Doyle, E.M., Brooks, S. and Connor, K.E.O., 2007. Biochemical characterization of the coexisting tyrosinase and laccase in the soil

- bacterium *Pseudomonas putida* F6. *Enzyme Microb. Technol.*, **40**: 1435–1441. <https://doi.org/10.1016/j.enzmictec.2006.10.020>
- Minussi, R.C., Miranda, M.A. and Silva, J.A., 2007. Purification, characterization and application of laccase from *Trametes versicolor* for colour and phenolic removal of olive mill wastewater in the presence of 1-hydroxybenzotriazole. *Afr. J. Biotechnol.*, **6**: 1248–1254.
- Minussi, R.C., Pastore, G.M. and Dur'an, N., 2002. Potential applications of laccase in the food industry. *Trends Fd. Sci. Technol.*, **13**: 205–216. [https://doi.org/10.1016/S0924-2244\(02\)00155-3](https://doi.org/10.1016/S0924-2244(02)00155-3)
- Mohammadian, M., Roudsari, M.F., Mollania, N., Dalfard, A.B. and Khajeh, K., 2010. Enhanced expression of a recombinant bacterial laccase at low temperature and microaerobic conditions: Purification and biochemical characterization. *J. Ind. Microbiol. Biotechnol.*, **5**: 41–45.
- Moilanen, U., Oasma, J.F., Winquist, E., Leisola, M. and Couto, S.R., 2010. Decolorization of simulated textile dye baths by crude laccases from *Trametes hirsute* and *Cerrena unicolor*. *Eng. Life Sci.*, **10**: 1–6. <https://doi.org/10.1002/elsc.200900095>
- Morozova, O.V., Shumakovich, G.P., Gorbacheva, M.A., Shleev, S.V. and Yaropolov, A.I., 2007. “Blue” Laccases. *J. Biochem.*, **72**: 1136–1150. <https://doi.org/10.1134/S0006297907100112>
- Murugesan, K., Dhamija, A., Nam, I.H., Kim, Y.M. and Chang, Y.S., 2007. Decolourization of reactive black 5 by laccase: Optimization by response surface methodology. *Dyes. Pig.* **75**: 176–184. <https://doi.org/10.1016/j.dyepig.2006.04.020>
- Narayanan, P.M. and Murugan, S., 2014. Production, purification and application of bacterial laccase: A review. *Biotechnology*, **13**: 196–205. <https://doi.org/10.3923/biotech.2014.196.205>
- Ncanana, S., Baratto, L., Roncaglia, L., Riva, S. and Burton, S.G., 2007. Laccase mediated oxidation of totarol. *Adv. Synth. Catal.*, **349**: 1507–1513. <https://doi.org/10.1002/adsc.200700005>
- Nicotra, S., Cramarossa, M.R., Mucci, A., Pagnoni, U.M., Riva, S. and Forti, L., 2004. Biotransformation of resveratrol: synthesis of *trans*-dehydrodimers catalyzed by laccases from *Myceliophthora thermophyla* and from *Trametes pubescens*. *Tetrahedr.*, **60**: 595–600. <https://doi.org/10.1016/j.tet.2003.10.117>
- Niladevi, K.N. and Prema, P., 2008. Immobilization of laccase from *Streptomyces psammotus* and its application in phenol removal using packed bed reactor. *World J. Microbiol. Biotechnol.*, **24**: 1215–1222. <https://doi.org/10.1007/s11274-007-9598-x>
- Niladevi, K.N., Sukumaran, R.K. and Prema, P., 2007. Utilization of rice straw for laccase production by *Streptomyces psammotus* in solid-state fermentation. *J. Ind. Microbiol. Biotechnol.*, **34**: 665–674. <https://doi.org/10.1007/s10295-007-0239-z>
- Palmieri, G., Giardina, P., Bianco, C., Fontallella, B. and Sannina, G., 2000. Copper induction of laccase isoenzyme in the lignolytic fungus *Pleurotus ostreatus*. *Appl. Microbiol. Biotechnol.*, **66**: 920–924. <https://doi.org/10.1128/AEM.66.3.920-924.2000>
- Palmieri, G., Giardina, P. and Marzullo, L., 1993. Stability and activity of a phenol oxidase from the ligninolytic fungus *Pleurotus ostreatus*. *Appl. Microbiol. Biotechnol.*, **39**: 632–636. <https://doi.org/10.1007/BF00205066>
- Papinutti, V.L., Diorio, L.A. and Forchiassin, F., 2003. Production of laccase and manganese peroxidase by *Fomes sclerodermeus* grown on wheat bran. *J. Ind. Microb. Biotechnol.*, **30**: 157. <https://doi.org/10.1007/s10295-003-0025-5>
- Paszczynski, A., Paszczynski, A., Pasti, M., Goszczynski, B.S., Crawford, D.L. and Crawford, R.L., 1991. New approach to improve degradation of recalcitrant azo dyes by *Streptomyces* spp. and *Phanerochaete chrysosporium*. *Enzyme Microbiol. Technol.*, **13**: 378–384. [https://doi.org/10.1016/0141-0229\(91\)90198-J](https://doi.org/10.1016/0141-0229(91)90198-J)
- Pedersen, L.S., Felby, C. and Munk, N., 2001. US 6187136.
- Peralta-Zamora, P., Pereira, C.M. and Tiburtius, E.R.L., 2003. Decolorization of reactive dyes by immobilized laccase. *Appl. Catal. B.*, **42**: 131–144. [https://doi.org/10.1016/S0926-3373\(02\)00220-5](https://doi.org/10.1016/S0926-3373(02)00220-5)
- Pointing, S.B., Jones, E.B.G. and Vrijmoed, L.L.P., 2000. Optimization of laccase production by *Pycnoporus sanguineus* in submerged liquid culture. *Mycologia*, **92**: 139–144. <https://doi.org/10.1080/00275514.2000.12061138>
- Ribeiro, D.S., Henrique, S.M.B., Oliveira, L.S., Macedo, G.A. and Fleuri, L.F., 2010. Enzymes in juice processing: A review. *Int. J. Fd. Sci. Technol.*, **45**: 635–641. <https://doi.org/10.1111/j.1365-2621.2010.02177.x>
- Roberts, S.A., Weichsel, A., Grass, G., Thakali, K., Hazzard, J.T., Tollin, G., Rensing, C. and Montfort, W.R., 2003. Crystal structure and electron transfer kinetics of CueO: a multicopper oxidase required for copper homeostasis in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA*, **99**: 2766–2771. <https://doi.org/10.1073/pnas.052710499>
- Roggen, E.L., Ernst, S., Svendsen, A., Friis, E.P. and Von Der Osten, C., 2001. WO2001083559 A2.
- Romero, S., Bl'aquez, P. and Caminal, G., 2006. Different approaches to improving the textile dye degradation capacity of *Trametes versicolor*. *Biochem. Eng. J.*, **31**: 42–47. <https://doi.org/10.1016/j.bej.2006.05.018>
- Ruijsenaars, H.J. and Hartmans, S., 2004. A cloned *Bacillus halodurans* multicopper oxidase exhibiting alkaline laccase activity. *Appl. Microbiol. Biotechnol.*, **65**: 177–182. <https://doi.org/10.1007/s00253-004-1571-0>
- Selvam, K., Swaminathan, K. and Chae, K.S., 2003.

- Decolourization of azo dyes and a dye industry effluent by a white rot fungus *Thelephora* sp. *Bioresour. Technol.*, **88**: 115–119. [https://doi.org/10.1016/S0960-8524\(02\)00280-8](https://doi.org/10.1016/S0960-8524(02)00280-8)
- Setti, L., Giuliani, S., Spinozzi, G. and Pifferi, P.G., 1999. Laccase catalyzed oxidative coupling of 3-methyl 2-benzothiazolinone hydrazone and methoxyphenols. *Enzyme Microb. Technol.*, **25**: 285–289. [https://doi.org/10.1016/S0141-0229\(99\)00059-9](https://doi.org/10.1016/S0141-0229(99)00059-9)
- Sharma, K.K. and Kuhad, R.C., 2008. Laccase: Enzyme revisited and function redefined. *Ind. J. Microbiol.*, **48**: 309–316. <https://doi.org/10.1007/s12088-008-0028-z>
- Sharma, P., Goel, R. and Caplash, N., 2007. Bacterial laccases. *World J. Microbiol. Biotechnol.*, **23**: 823–832. <https://doi.org/10.1007/s11274-006-9305-3>
- Shi, J., 2006. CN1844572 A.
- Singh, G., Ahuja, N., Batish, M., Capalash, N. and Sharma, P., 2008. Biobleaching of wheat straw rich soda pulp with alkalophilic laccase from *γ-Proteobacterium* JB: optimization of process parameters using response surface methodology. *Bioresour. Technol.*, **99**: 7472–7479. <https://doi.org/10.1016/j.biortech.2008.02.023>
- Singh, G., Capalash, N., Goel, R. and Sharma, P., 2007. A pH-stable laccase from alkali-tolerant *γ-proteobacterium* JB: purification, characterization and indigo carmine degradation. *Enzyme Microb. Technol.*, **41**: 794–799. <https://doi.org/10.1016/j.enzmictec.2007.07.001>
- Singh, G., Capalash, N. and Sharma, P., 2009b. Performance of an alkalophilic and halo tolerant laccase from *γ-proteobacterium* JB in the presence of industrial pollutants. *J. Gen. Appl. Microbiol.*, **55**: 283–289. <https://doi.org/10.2323/jgam.55.283>
- Singhania, R.R., Sukumarana, R.K., Patel, A.K., Larroche, C. and Pandey, A., 2010. Advancement and comparative profiles in the production technologies using solid-state and submerged fermentation for microbial cellulases. *Enzyme Microb. Tech.*, **46**: 541. <https://doi.org/10.1016/j.enzmictec.2010.03.010>
- Solomon, E.I., Sundaram, U.M. and Machonkin, T.E., 1996. Multicopper oxidases and oxygenases. *Chem. Rev.*, **96**: 2563–2605. <https://doi.org/10.1021/cr950046o>
- Suzuki, T., Endo, K., Ito, M., Tsujibo, H., Miyamoto, K., Inamori, Y., 2003. A thermostable laccase from *Streptomyces lavendulae* REN-7: Purification, characterization, nucleotide sequence and expression. *Biosci. Biotechnol. Biochem.*, **76**: 2167–2175. <https://doi.org/10.1271/bbb.67.2167>
- Svendsen, A. and Xu, F., 2001. US 6184015.
- Tavares, A.P.M., Cristovao, R.O., Gamelas, J.A.F., Loureiro, J.M., Boaventuraa, R.A.R. and Macedoa, E.A., 2009. Sequential decolourization of reactive textile dyes by laccase mediator system. *J. Chem. Technol. Biotechnol.*, **84**: 442–446. <https://doi.org/10.1002/jctb.2060>
- Téllez-Jurado, A., Arana-Cuenca, A. and González-Becerra, A.E., 2005. Expression of a heterologous laccase by *Aspergillus niger* cultured by solid-state and submerged fermentations. *Enzyme Microb. Tech.*, **38**: 665. <https://doi.org/10.1016/j.enzmictec.2005.07.021>
- Thiruchelvam, A.T. and Ramsay, J.A., 2007. Growth and laccase production kinetics of *Trametes versicolor* in a stirred tank reactor. *Applied Microbial. Biotechnology*, **74**: 547–554. <https://doi.org/10.1007/s00253-006-0695-9>
- Thurston, C.F., 1992. The structure and function of fungal laccases. *Microbiology*, **140**: 19–26. <https://doi.org/10.1099/13500872-140-1-19>
- Thurston, C., 1994. The structure and function of fungal lacases. *Microbiology*, **140**: 19–26.
- Wang, H.X., Ng, T.B., 2004. A novel laccase with fair thermostability from the edible wild mushroom (*Albatrella dispansus*). *Biochem. Biophys. Res. Commun.*, **315**: 450–454. [https://doi.org/10.1016/S0006-291X\(04\)00967-2](https://doi.org/10.1016/S0006-291X(04)00967-2)
- Wesenberg, D., Kyriakides, I. and Agathos, N., 2003. White-rot fungi and their enzymes for the treatment of industrial dye effluents. *Biotechnol. Adv.*, **22**: 161–187. <https://doi.org/10.1016/j.biotechadv.2003.08.011>
- Xu, F., 1999. Laccase, In Flickinger, M.C. and Drew, S.W. (eds.), *Encyclopedia of bioprocess technology: Fermentation, Biocatalysis, Bioseparation* John Wiley and Sons Inc., New York, pp. 1545–1554.
- Xu, H., Bloomfield, K. and Lund, H., 2006. WO2006126983 A1.
- Xu, Q., Fu, Y., Qin, M. and Li, Z., 2006. CN1763305 A.
- Yague, S., Terron, M.C., Gonzalez, T., Zapico, E., Bocchini, P., Galetti, G.C. and Gonzalez, A.E., 2000. Biotreatment of tannin rich beer factory waste water with white rot basidiomycete *Coriolopsis gallica* monitored by pyrolysis/gas chromatography/mass spectrometry. *Rapid. Commun. Mass Spectrom.*, **14**: 905–910. [https://doi.org/10.1002/\(SICI\)1097-0231\(20000530\)14:10<905::AID-RCM963>3.0.CO;2-7](https://doi.org/10.1002/(SICI)1097-0231(20000530)14:10<905::AID-RCM963>3.0.CO;2-7)
- Ye, M., Li, G., Liang, W.Q. and Liu, Y.H., 2010. Molecular cloning and characterization of a novel metagenome-derived multicopper oxidase with alkaline laccase activity and highly soluble expression. *Appl. Microbiol. Biotechnol.*, **87**: 1023–1031. <https://doi.org/10.1007/s00253-010-2507-5>
- Zang, G.Q., Wang, Y.F., Zang, X.Q., Ng, T.B. and Wang, H.X., 2010. Purification and characterization of a novel laccase from the edible mushroom *Clitocybe maxima*. *Process. Biochem.*, **45**: 627–633. <https://doi.org/10.1016/j.procbio.2009.12.010>