



Research Article

Field Evaluation of Promising Botanical Extracts, Plant Essential Oils and Differential Chemistry Insecticides against Subterranean Termites *Odontotermes obesus* (Isoptera: Termitidae)

Muhammad Shahzad Akbar, Farrukh Sajjad, Muhammad Afzal, Muhammad Luqman, Muhammad Asam Riaz and Muhammad Zeeshan Majeed*

College of Agriculture, University of Sargodha, Sargodha 40100, Pakistan.

Abstract | Subterranean termites are destructive pests of agricultural crops, forest and orchard plantations and wooden infrastructures. A wide range of persistent synthetic chemicals are employed to prevent and control the infestations of subterranean termites. Most of these chemicals have high mammalian toxicity and environmental hazards. This study was, therefore, aimed to evaluate the efficacy of some promising botanical extracts (*Dodonaea viscosa*, *Gardenia jasminoides* and *Nerium indicum*), plant essential oils (*Allium sativum*, *Citrus aurantium* and *Cymbopogon citratus*) and differential-chemistry insecticides (chlorantraniliprole, chlorfenapyr, emamectin, indoxacarb, pyriproxyfen and triflumuron) against subterranean termites (*Odontotermes obesus*) attacking sugarcane crop. Setts of sugarcane variety HSF555 were sown on the ridges under RCB design after treating them with 20% botanical extracts, 2.0% essential oils and differential-chemistry insecticides with their field recommended dose rates. Results revealed that as compared to control plots, all treatment plots exhibited reduced bud and shoot damage with maximum bud germination and minimum termite infestation. Maximum bud germination was recorded in chlorantraniliprole (92.99%) and chlorfenapyr (92.03%), followed by *C. aurantium* (92.84%) and *A. sativum* (85.78%) as compared to minimum bud germination in control (40.85%). Highest bud and shoot damage was recorded in control (61.92 and 48.15%, respectively), while the lowest was recorded in *C. aurantium* (16.18 and 15.56%, respectively). Similarly, minimum per sett termite counts was recorded in chlorantraniliprole (16.25) treated plots, followed by chlorfenapyr (22.83), pyriproxyfen (30.04), *A. sativum* (28.50) and *C. aurantium* (29.00), while maximum termite infestation was recorded in control (68.50 individuals per sett). In brief, chlorantraniliprole, chlorfenapyr and pyriproxyfen among differential-chemistry insecticides and *A. sativum*, *C. aurantium* and *G. jasminoides* among botanicals were the most effective at preventing the upsurge of termite infestation till 90 days of sett sowing, and are recommended to be used as eco-friendly options for controlling subterranean termites in sugarcane.

Received | August 12, 2020; **Accepted** | December 08, 2020; **Published** | February 06, 2021

***Correspondence** | Muhammad Zeeshan Majeed, College of Agriculture, University of Sargodha, Sargodha 40100, Pakistan; **Email:** zeeshan.majeed@uos.edu.pk

Citation | Akbar, MS, F. Sajjad, M. Afzal, M. Luqman, M.A. Riaz and M.Z. Majeed. 2021. Field evaluation of promising botanical extracts, plant essential oils and differential chemistry insecticides against subterranean termites *Odontotermes obesus* (Isoptera: Termitidae). *Sarhad Journal of Agriculture*, 37(1): 120-127.

DOI | <http://dx.doi.org/10.17582/journal.sja/2021/37.1.120.127>

Keywords | Subterranean termites, *Odontotermes*, Sugarcane, Botanicals, Differential-chemistry insecticides, Chlorantraniliprole, Chlorfenapyr, *Allium sativum*, *Citrus aurantium*

Introduction

With approximately 3,500 described species, termites constitute an important insect

fauna of tropical and temperate ecosystems (Engel, 2011; Majeed, 2012). Many termite species such as subterranean termites are destructive pests of agricultural crops, tree plantations and wooden

structures (Rouland-Lefèvre, 2010; Buczkowski and Bertelsmeier, 2017). *Odontotermes*, *Microtermes* and *Coptotermes* are the most important genera of subterranean termites reported from Pakistan (Ahmed *et al.*, 2005; Manzoor *et al.*, 2011; Sattar *et al.*, 2014) and *O. obesus* is the most common subterranean termite species attacking a number of agricultural crops including sugarcane, maize, sesame, cotton, wheat and gram (Sattar *et al.*, 2014; Aihetasham *et al.*, 2018).

Synthetic insecticides have been playing a major role in termite management all over the world including Pakistan. In Indo-Pak regions, a wide range of synthetic insecticides including DDT, thiodan, deltamethrin, carbosulfan and triazophos are applied against subterranean termites (Rana and Dahiya, 2001; Ahmed *et al.*, 2006; Manzoor *et al.*, 2012; Iqbal and Saeed, 2013). Most of these conventional synthetic insecticides are highly persistent with long residual effects causing the eradication of non-target organisms, including soil microbiota, predators and parasitoids, environmental contamination and insect pest resistance (Baker and Bellamy, 2006; Desneux *et al.*, 2007; Edwards, 2013).

Nevertheless, plant based insecticides and differential chemistry synthetic insecticides are emerging as relatively safer pest control tactics than conventional ones. Plants extracts and essential oils would be suitable alternate of synthetic insecticides (Ahmed *et al.*, 2007; Isman, 2008; Dubey *et al.*, 2010). Plant derived compounds are considered as a pool of chemical substances which have potential uses against a wide range of insect pests including subterranean termites (Bläske and Hertel, 2001; Peterson and Ems-Wilson, 2003; Ahmed *et al.*, 2007). For instance, essential oils extracted from eucalyptus, vetiver plant, clove bud, cedar wood, lemongrass, geranium etc. have been found exhibiting certain deterrent and poisonous effects against the subterranean termites (Isman, 2000; Zhu *et al.*, 2001; Ahmed *et al.*, 2007; Ibrahim and Demisse, 2013).

Similarly, synthetic insecticides with differential chemistry and mode of action than the conventional insecticidal groups are emerging as promising tools to mitigate the problems of pest resistance and environmental contamination. These pesticides such as chlorantraniliprole, chlorfenapyr, emamectin benzoate, indoxacarb, pymetrozine, pyriproxyfen etc.

are less persistent and more target specific and are safer to non-target fauna (Grafton-Cardwell *et al.*, 2005; Ishaaya and Degheele, 2013). Many previous studies have demonstrated the efficacy of differential chemistry insecticides against different termite species (Mao *et al.*, 2011; Rashid *et al.*, 2012; Iqbal and Saeed, 2013).

Therefore, in order to avoid the above cited detrimental effects of conventional synthetic insecticides, these should be replaced with environment friendly and biorational tactics for the control of subterranean termites. To this end, the present study was conducted to evaluate the field efficacy of some selected botanical extracts, plant essential oils and differential chemistry insecticides against subterranean termites (*O. obesus*) in a sugarcane field with an ultimate objective to find out the most effective treatments which can be recommended to the indigenous sugarcane growers for better management of termite infestations.

Materials and Methods

Experiment site

The experiment was conducted from October, 2016 to January, 2017 in a village Chak No. 101 (31°58'5.6" N; 72°45'9.4" E) located at Kandiwal Road, Sargodha (Punjab, Pakistan). The average annual rainfall and temperature of this area is 410 mm and 23.8°C, respectively (Zaka *et al.*, 2004). Wheat, rice, maize and sugarcane are main agricultural crops of the area. Soil texture is sandy loam. The sugarcane field selected for the experiment was ensured to have severe termite infestation each year and it did not receive any pesticidal application for last 4 to 5 months prior to the experiment.

Treatments

Botanical extracts and essential oils (Table 1) were prepared in the laboratory of the Department of Entomology, College of Agriculture, University of Sargodha. In brief, different plant parts as described in Table 1 were collected from the vicinity of the College of Agriculture, University of Sargodha and were washed with distilled water and were air-dried at room temperature (27°C) for few days followed by grinding of plant material to coarse powder form. Extraction of plant extracts was carried out through Soxhlet apparatus (Sigma-Aldrich, Germany) using 1:10 (w/v) methanol as extraction solvent. In brief, 50 g of plant powder was extracted using 500 mL

of methanol. Plant essential oils were extracted by the hydro distillation process using Clevenger-type apparatus. Formulations of differential chemistry insecticides (Table 2) were purchased from authorized dealers of multinational companies from the local pesticide market of the Sargodha district.

Experimental protocol

The experiment was laid out in a randomized complete block design (RCBD). Sugarcane variety HSF555 was sown on well-prepared soil ridges. Plant to plant and row to row distance was maintained at 20 and 90 cm, respectively. At the time of sowing, sugarcane setts (each having 3 bud-eyes) were dipped for two minutes in solutions of six botanicals including three essential oils (*C. citratus*, *C. aurantium* and *A. sativum* @ 2%), three crude plant extracts (*G. jasminoides*, *D. viscosa* and *N. indicum* @ 20%) and six differential-chemistry insecticides. Each treatment was independently and randomly replicated thrice in

three blocks of the experiment. A distance of 10 ft was maintained in between two replication plots as buffer zone. The total study area was 5,715 ft² and the dimension of each plot was 8 x 10 ft. All routine agronomic practices were carried out for all plots.

Data collection

Bud germination: Germination of buds on the sown sugarcane setts was recorded by tallying the germinated buds out of total setts sown in each plot. Bud germination data was taken after 30 days of planting.

Bud and shoot damage: Bud damage was assessed by exposing the buds in each plot randomly. From each plot, randomly five sugarcane setts were excavated and examined for damage of buds by subterranean termites. The bud damage data was assessed after 50 days of planting. Similarly, percentage of shoot damage was assessed after 70 days of planting.

Table 1: Different botanical extracts and essential oils evaluated under field conditions against subterranean termites (*Odontotermes obesus*) in sugarcane crop.

Plants extracted	Major bioactive constituents	Extraction type
Bulbs of garlic (<i>Allium sativum</i>)	Diallyl di- and tri-sulfides, dipropyl di-sulfides (Zhao <i>et al.</i> , 2013)	2% essential oil
Fruit peels and seeds of sour orange (<i>Citrus aurantium</i>)	Flavanone, β -pinene, α -terpineol, limonenes (Ibrahim <i>et al.</i> , 2001)	2% essential oil
Leaves of lemon grass (<i>Cymbopogon citratus</i>)	Citral, citronellol, citronella, myrcene (Dodia <i>et al.</i> , 2010)	2% essential oil
Stems of sanatha (<i>Dodonaea viscosa</i>)	Flavonoids, phenols, tannins, saponins, lupeol, and stigmasterol (Al-Snafi, 2017)	20% aqueous extract
Leaves and stems of gardenia (<i>Gardenia jasminoides</i>)	Iridoid glycosides (Li <i>et al.</i> , 2018)	20% aqueous extract
Leaves of oleander (<i>Nerium indicum</i>)	Oleandrin and oleandrogenin (Dodia <i>et al.</i> , 2010)	20% aqueous extract

Table 2: Different insecticidal formulations evaluated under field conditions against subterranean termites (*Odontotermes obesus*) in sugarcane crop.

Chemical Name (active ingredient)	Chemical family*	Mode of action	Brand Name	Company	Label dose (a.i. ha ⁻¹)
Chlorantraniliprole	28 (diamides)	Ryanodine receptor modulators	Coragen® 18.5 SC	DuPont	375 ml
Chlorfenapyr	13 (pyrroles)	Uncoupler of oxidative phosphorylation	Pirate® 360 SC	Swat Agro Chemicals	500 ml
Emamectin	6 (avermectins)	Glutamate-gated chloride channel (Glu-Cl) allosteric modulators	Proclaim® 1.9 EC	Syngenta	500 ml
Indoxacarb	22A (oxadiazines)	Voltage-dependent sodium channel blockers	Steward® 150 EC	DuPont	375 ml
Pyriproxyfen	7C (pyriproxyfens)	Juvenile hormone mimics (IGR)	Admiral® 10 EC	FMC	75 ml

*According to insecticide resistance action committee (www.irac-online.org) IRAC MoA classification version 8.3, July 2017.

Table 3: Comparison of percent mean bud germination, bud damage, shoot damage and termite count recorded in sugarcane plots treated with different botanical extracts, plant essential oils and synthetic insecticides.

Treatments	Bud germination (%)		Bud damage (%)		Shoot damage (%)		Termite count	
<i>Gardenia jasminoides</i>	83.22	±7.25 ^{ab,*}	30.09	±1.74 ^{cd}	30.27	±0.95 ^{bcd}	31.75	±6.29 ^{bc}
<i>Dodonaea viscosa</i>	70.33	±20.07 ^{abc}	34.99	±3.44 ^{cd}	24.82	±7.49 ^{cde}	33.75	±4.11 ^{bc}
<i>Nerium indicum</i>	76.03	±14.44 ^{abc}	40.85	±1.31 ^b	36.90	±2.04 ^b	37.50	±3.42 ^b
<i>Citrus aurantium</i>	90.84	±8.11 ^a	16.18	±2.35 ^f	18.04	±4.15 ^{ef}	29.00	±5.60 ^{bcd}
<i>Allium sativum</i>	85.78	±10.86 ^a	18.96	±3.34 ^{ef}	21.49	±2.47 ^{def}	28.50	±7.94 ^{bcd}
<i>Cymbopogon citratus</i>	77.70	±8.99 ^{abc}	32.18	±6.19 ^{bcd}	21.85	±2.12 ^{def}	36.50	±5.45 ^{bc}
Triflumuron	45.63	±16.64 ^{bc}	27.63	±3.65 ^{cde}	29.60	±2.23 ^{bcd}	34.75	±6.08 ^{bc}
Indoxacarb	66.88	±16.96 ^{abc}	35.23	±3.44 ^{bcd}	18.30	±5.67 ^{ef}	32.50	±4.80 ^{bc}
Pyriproxyfen	72.26	±13.25 ^{abc}	28.82	±2.21 ^{cd}	35.79	±1.53 ^b	29.50	±6.56 ^{bcd}
Chlorfenapyr	92.03	±8.88 ^a	25.97	±4.12 ^{def}	27.98	±3.03 ^{bcd}	22.50	±2.89 ^{cd}
Chlorantraniliprole	92.99	±7.47 ^a	18.67	±2.52 ^{ef}	15.56	±2.80 ^f	16.25	±4.11 ^d
Emamectin	71.49	±20.06 ^{abc}	37.41	±2.47 ^{bc}	32.04	±3.01 ^{bc}	32.00	±5.35 ^{bc}
Control	40.85	±10.18 ^c	61.92	±3.50 ^a	48.52	±2.83 ^a	68.50	±8.66 ^a

*Values are means (±SE) of five setts/plants randomly selected from three independent replications. Different alphabets indicate the statistical significance among treatment means within each column (two-way factorial ANOVA; Tukey HSD at $\alpha = 0.05$).

Termites count: The termite population was counted after 90 days of planting. For determining the termite population, sown setts were excavated out and the total individuals of subterranean termites on each plant (sett) were enumerated. Termites were identified by the termite experts as *O. obesus*.

Statistical analysis

Using Statistix® 8.1 (Analytical Software, 2005), data regarding bud germination, bud and shoot damage and termite infestation were subjected to statistical analyses by two-way ANOVA. Treatment means were compared using Tukey HSD at 0.95% level of significance.

Results and Discussion

Table 3 shows the effect of differential-chemistry insecticides, plant extracts and essential oils on sugarcane bud germination, bud and shoot damage and termite infestation. According to data analysis, there was a significant effect of all treatments (insecticides and botanicals) on the bud germination ($F_{12,38} = 4.52$; $P < 0.01$), bud damage ($F_{12,38} = 36.23$; $P < 0.001$) and shoot damage ($F_{12,38} = 18.95$; $P < 0.001$) of sugarcane setts and on the termite infestation level ($F_{12,38} = 21.41$; $P < 0.001$) (Supplementary Table 1).

Bud germination varied from maximum (92.99%) for chlorantraniliprole to minimum (45.63%) for triflumuron as compared to control plots which

exhibited 40.85% bud germination, statistically lower than that of other treatments. In case of botanicals, *C. aurantium* and *A. sativum* exhibited highest mean bud germination (i.e. 90.84 and 85.78%, respectively) and were statistically different from other botanicals (Table 3). Among insecticides, chlorantraniliprole exhibited minimum bud damage (18.67%) followed by chlorfenapyr (27.63%) and pyriproxyfen (28.82%), while maximum bud damage (~38%) was recorded in plots treated with emamectin and indoxacarb. Among botanicals, bud damage varied from maximum (40.85%) for *N. indicum* to minimum (16.18%) for *C. aurantium* as compared to control plots which showed maximum (61.92%) bud damage, statistically different from other treatments (Table 3). Similarly, shoot damage varied from maximum (36.90%) for *N. indicum* to minimum for *C. aurantium* and *A. sativum* (i.e. 18.04 and 21.27%, respectively). Among insecticides, chlorantraniliprole showed significantly less shoot damage (15.56%) than all other treatments. Maximum shoot damage (48.52%) was observed in control plots and was statistically different from all other treatments.

At 90 days of sett sowing, infestation of termites was assessed by counting termite individuals on sugarcane plants. Highest mean termite count was recorded in control plots (68.50 individuals per sett) and was statistically different from all other treatments. Among synthetic insecticides, lowest termite count was recorded in chlorantraniliprole treated plots

(16.25 individuals per sett), followed by chlorfenapyr (22.83 individuals per sett) and pyriproxyfen (30.04 individuals per sett). While among botanical treatments, essential oils of *C. aurantium* and *A. sativum* exhibited minimum termites counts (~29.42 individuals per sett), followed by botanical extract of *G. jasminoides* (31.75 individuals per sett) (Table 3).

Subterranean termites are destructive pests of agricultural crops including sugarcane, maize, sorghum, rice, wheat, cotton, gram and groundnut in Pakistan (Ahmed *et al.*, 2011). To control these pests, farmers primarily rely on extensive and irrational application of persistent synthetic insecticides and most of these chemicals are notorious for their hazards to human health and environment (Mulrooney *et al.*, 2006; Edwards, 2013). To mitigate pesticidal side-effects, searching for relatively safer and target-specific insecticidal tactics has been a core area of plant protection research. Wherefore, this study evaluated the efficacy of some promising indigenous botanicals (including plant extracts and essential oils) and differential-chemistry insecticides (which are less persistent and comparatively safer than conventional ones) against subterranean termites under field conditions.

Results of the study showed that among botanicals, essential oils of *C. aurantium* and *A. sativum* were the most effective to suppress subterranean termite infestation under field conditions. Our results are in agreement with Ahmed *et al.* (2005) and Owusu *et al.* (2008) who demonstrated *in-situ* efficacy of different indigenous plant extracts against sugarcane infesting termites (*Microtermes* spp.) suggesting that these indigenous plant materials could be utilized for effective and biorational management of subterranean termites (Ahmed *et al.*, 2007). Nevertheless, extracts of *C. aurantium* and *A. sativum* have been found effective against many insect pests (Hollingsworth, 2005; Samarasinghe *et al.*, 2007; Siskos *et al.*, 2009; Zhao *et al.*, 2013) including subterranean termites (Park and Shin, 2005; Osipitan *et al.*, 2013).

Our results demonstrated that botanical extracts and essential oils showed more or less similar results as compared to differential chemistry synthetic insecticides although the later were statistically more effective. This is in accordance with Ibrahim and Demisse (2013) who used two plant extracts (*Maesalan ceolata* and *A. indica*) and a synthetic

insecticide Diazinon 60EC against termites and found that botanicals were as effective as synthetic chemical. Our results are also in line with those of Daniel and Bekele (2006) who demonstrated the efficacy of plant extracts statistically similar to chlorpyrifos against *Macrotermes* spp.

In case of differential chemistry insecticides, chlorantraniliprole and chlorfenapyr appeared to be the most effective insecticides against *O. obesus* termite infestation in sugarcane field. These results corroborate the findings of Spomer *et al.* (2009), Mao *et al.* (2011), Neoh *et al.* (2012) and Ma and Sui (2013) who demonstrated that these differential-chemistry insecticides are highly effective against subterranean termites. Apart from chlorantraniliprole and chlorfenapyr, indoxacarb and pyriproxyfen were also effective differential chemistry insecticides causing minimum bud and shoot damage concomitantly with minimum termite infestation. These results are in line with those of Rust and Saran (2006), Rai (2014), Misbah-ul-Haq *et al.* (2016), Lewis and Forscher (2017), Akbar *et al.* (2018) and Sapkota (2018).

Conclusions and Recommendations

In brief, this study was aimed to evaluate *in-situ* efficacy of different biorational insecticides against subterranean termite infestation in sugarcane crop. Based on results of the study, essential oils of *A. sativum* and *C. aurantium* and differential chemistry insecticidal formulations of chlorantraniliprole, chlorfenapyr, indoxacarb and pyriproxyfen are recommended as biorational control options for the management of *O. obesus* and other subterranean termites. In short, the study corroborates the effectiveness of indigenous botanical extracts and differential chemistry insecticides against subterranean termite infestations in sugarcane under field conditions.

Acknowledgements

This laboratory work was financially supported by a research project (No. 6702) funded by the Higher Education Commission (HEC) of Pakistan under its National Research Program for Universities (NRPU).

Novelty Statement

This comparative *in-situ* evaluation of different

biorational pesticides against subterranean termites infesting sugarcane crop demonstrated that chlorantraniliprole, chlorfenapyr and pyriproxyfen among the differential chemistry insecticides and *A. sativum*, *C. aurantium* and *G. jasminoides* among the botanicals are effective at preventing termite infestation suggesting their future use as environmental-friendly termite management options in the field.

Authors' Contributions

MSA and MZM conceived the idea and planned the experiment. MSA and FS performed experiments and wrote first draft of the manuscript. MAR and ML performed statistical analyses. MAR and MA technically revised the manuscript. MA provided technical assistance and proofread the manuscript.

Supplementary Material

There is supplementary material associated with this article. Access the material online at: <http://dx.doi.org/10.17582/journal.sja/2021/37.1.120.127>

Conflict of interest

The authors have declared no conflict of interest.

References

- Ahmed, S., A. Hussain, M.I. Zafar, A. Hussain, M.A. Riaz and M. Shahid. 2011. Evaluation of plant extracts on mortality and tunneling activities of subterranean termites in Pakistan. In: Pesticides in the modern world-Pests control and pesticides exposure and toxicity assessment (ed. M. Stoytcheva). Intech Open Access Publisher. pp. 39-54. <https://doi.org/10.5772/18999>
- Ahmed, S., R.R. Khan and M.A. Riaz. 2007. Some studies on the field performance of plant extracts against termites (*Odontotermes guptai* and *Microtermes obesi*) in sugarcane at Faisalabad. Int. J. Agric. Biol., 9(3): 398-400.
- Ahmed, S., S. Fiaz, M.A. Riaz and A. Hussain. 2005. Comparative efficacy of *Datura alba* nees, *Calotropis procera* and Imidacloprid on termites in sugarcane at Faisalabad. Pak. Ent., 27(2): 11-14.
- Ahmed, S., T. Mustafa, M.A. Riaz and A. Hussain. 2006. Efficacy of insecticides against subterranean termites in sugarcane. Int. J. Agric. Biol., 8(4): 508-510.
- Aihetasham, A., H. Aziz and K.Z. Rasib. 2018. Efficacy of slow acting toxicants on *Heterotermes indicola* (Wasmann) (Isoptera: Rhinotermitidae). Pak. J. Zool., 50(1): 373-373. <https://doi.org/10.17582/journal.pjz/2018.50.1.sc3>
- Akbar, M.S., M.Z. Majeed and M. Afzal. 2018. Comparative toxicity of selected new-chemistry insecticides against subterranean termites *Odontotermes obesus* Ramb. (Isoptera: Termitidae). Sarhad J. Agric., 35(1): 20-26. <https://doi.org/10.17582/journal.sja/2019/35.1.20.26>
- Al-Snafi, A.E., 2017. A review on *Dodonaea viscosa*: A potential medicinal plant. IOSR J. Pharm., 7: 10-21. <https://doi.org/10.9790/3013-0702011021>
- Baker, P.B. and D.E. Bellamy. 2006. Field and laboratory evaluation of persistence and bioavailability of soil termiticides to desert subterranean termite *Heterotermes aureus* (Isoptera: Rhinotermitidae). J. Econ. Ent., 99(4): 1345-1353. <https://doi.org/10.1603/0022-0493-99.4.1345>
- Bläske, V.U. and H. Hertel. 2001. Repellent and toxic effects of plant extracts on subterranean termites (Isoptera: Rhinotermitidae). J. Econ. Ent., 94(5): 1200-1208. <https://doi.org/10.1603/0022-0493-94.5.1200>
- Buczkowski, G. and C. Bertelsmeier. 2017. Invasive termites in a changing climate: A global perspective. Ecol. Evol., 7: 974-985. <https://doi.org/10.1002/ece3.2674>
- Daniel, G. and J. Bekele. 2006. Evaluation of toxicity of crude extracts of some botanicals on different castes of *Macrotermes* termites. Pest Manage. J. Ethiopia, 8: 16-22.
- Desneux, N., A. Decourtye and J.M. Delpuech. 2007. The sublethal effects of pesticides on beneficial arthropods. Ann. Rev. Ent., 52: 81-106. <https://doi.org/10.1146/annurev.ento.52.110405.091440>
- Dodia, D.A., I.S. Patel and G.M. Patel. 2010. Botanical Pesticides for Pest Management. Sci. Pub., Jodhpur, India. pp. 354.
- Dubey, N.K., R. Shukla, A. Kumar, P. Singh and B. Prakash. 2010. Prospects of botanical pesticides in sustainable agriculture. Curr. Sci., 98: 479-480.
- Edwards, C.A., 2013. Environmental pollution

- by pesticides. Springer Science and Business Media, New York. Vol. 3.
- Engel, M.S., 2011. Family group names for termites (Isoptera). Zookeys., 148: 171-184. <https://doi.org/10.3897/zookeys.148.1682>
- Grafton-Cardwell, E., L. Godfrey, W. Chaney and W. Bentley. 2005. Various novel insecticides are less toxic to humans, more specific to key pests. Calif. Agric., 59: 29-34. <https://doi.org/10.3733/ca.v059n01p29>
- Hollingsworth, R.G., 2005. Limonene, a citrus extract, for control of mealybugs and scale insects. J. Econ. Ent., 98(3): 772-779.
- Ibrahim, A. and G. Demisse, 2013. Evaluation of some botanicals against termites' damage on hot pepper at Bako, Western Ethiopia. Int. J. Agric. Policy Res., 1(2): 48-52.
- Ibrahim, M.A., P. Kainulainen, A. Aflatuni, K. Tiilikkala and J.K. Holopainen. 2001. Insecticidal, repellent, antimicrobial activity and phytotoxicity of essential oils: with special reference to limonene and its suitability for control of insect pests. Agric. Food Sci. Finland, 10: 243-259. <https://doi.org/10.23986/afsci.5697>
- Iqbal, N. and S. Saeed. 2013. Toxicity of six new chemical insecticides against the termite, *Microtermes mycophagus* D. (Isoptera: Termitidae: Macrotermitinae). Pak. J. Zool., 45(3): 709-713.
- Ishaaya, I. and D. Degheele. 2013. Insecticides with novel modes of action: Mechanisms and application. Springer Science and Business Media.
- Isman, M.B. 2008. Botanical insecticides: for richer, for poorer. Pest Manage. Sci., 64(1): 8-11. <https://doi.org/10.1002/ps.1470>
- Isman, M.B., 2000. Plant essential oils for pest and disease management. Crop Prot., 19(8): 603-608. [https://doi.org/10.1016/S0261-2194\(00\)00079-X](https://doi.org/10.1016/S0261-2194(00)00079-X)
- Lewis, J.L. and B.T. Forschler. 2017. Transfer of five commercial termite bait formulations containing benzoylphenyl urea chitin synthesis inhibitors within groups of the subterranean termite *Reticulitermes flavipes* (Blattodea: Rhinotermitidae). Int. J. Pest Manage., 63(3): 224-233. <https://doi.org/10.1080/09670874.2016.1241911>
- Li, L., J. Zou, Q. Xia, H. Cui, S. You, Y. Liu and Q. Wang. 2018. Anti-TMV and insecticidal potential of four iridoid glycosides from *Gardenia Jasminoides* fruit. Chem. Res. Chinese Univ., 34: 697-699. <https://doi.org/10.1007/s40242-018-8197-8>
- Ma, Y.J. and X.F. Sui, 2013. Comparison on efficacy of chlorantraniliprole and bifenthrin against *Coptotermes formosanus*. Chinese J. Hyg. Insectic. Equip., 4: 015.
- Majeed, M.Z., 2012. Emissions of nitrous oxide by tropical soil macrofauna: Impact of feeding guilds and microbial communities involved. Doctoral dissertation, University of Montpellier II, France.
- Manzoor, F., A.H. Sayyed, T. Rafique and S.A. Malik. 2012. Toxicity and repellency of different insecticides against *Heterotermes indicola* (Isoptera: Rhinotermitidae). J. Anim. Plant Sci., 22(1): 65-71.
- Manzoor, F., M. Chaudhary, N. Sheikh, I.A. Khan and T. Khan. 2011. Diversity and proportion of termite species in garden trees and wheat crop in district Bhakkar, Pakistan. Pak. J. Zool., 43(3): 537-541.
- Mao, L., G.G. Henderson and C.W. Scherer. 2011. Toxicity of seven termiticides on the Formosan and eastern subterranean termites. J. Econ. Ent., 104(3): 1002-1008. <https://doi.org/10.1603/EC11005>
- Misbah-Ul-Haq, M., I.A. Khan, A. Farid, M. Ullah, D.H. Gouge and P.B. Baker. 2016. Efficacy of indoxacarb and chlorfenapyr against subterranean termite *Heterotermes indicola* (Wasmann) (Isoptera: Rhinotermitidae) in the laboratory. Turk. J. Ent., 40(3): 227-241. <https://doi.org/10.16970/ted.89871>
- Mulrooney, J.E., M.K. Davis, T.L. Wagner and R.L. Ingram. 2006. Persistence and efficacy of termiticides used in preconstruction treatments to soil in Mississippi. J. Econ. Ent., 99(2): 469-475. <https://doi.org/10.1093/jee/99.2.469>
- Neoh, K.B., J. Hu, B.H. Yeoh and C.Y. Lee. 2012. Toxicity and horizontal transfer of chlorantraniliprole against the Asian subterranean termite *Coptotermes gestroi* (Wasmann): Effects of donor: Recipient ratio, exposure duration and soil type. Pest Manage. Sci., 68(5): 749-756. <https://doi.org/10.1002/ps.2322>
- Osipitan, A.A., T.O. Jegede, D.I. Adekanmbi and I.A. Ogunbanwo. 2013. Assessment of *Datura metel*, local soap and garlic (*Allium sativum*)

- in the management of termite (Termitidae: Isoptera). *Mun. Ent. Zool.*, 8(1): 407-414.
- Owusu, E.O., K.S. Akutse and K. Afreh-Nuamah. 2008. Effect of some traditional plant components on the control of termites, *Macrotermes* spp. (Isoptera: Termitidae). *Asian J. Sci. Technol.*, 9(2): 83-91.
- Park, I.K. and S.C. Shin. 2005. Fumigant activity of plant essential oils and components from garlic (*Allium sativum*) and clove bud (*Eugenia caryophyllata*) oils against the Japanese termite (*Reticulitermes speratus* Kolbe). *J. Agric. Food Chem.*, 53(11): 4388-4392. <https://doi.org/10.1021/jf050393r>
- Peterson, C. and J. Ems-Wilson. 2003. Catnip essential oil as a barrier to subterranean termites (Isoptera: Rhinotermitidae) in the laboratory. *J. Econ. Ent.*, 96(4): 1275-1282. <https://doi.org/10.1093/jee/96.4.1275>
- Rai, V.L., 2014. Studies on diversity of major insect-pest of sugarcane along with efficacy of some novel insecticides against termites in field and household ecosystem. Doctoral dissertation. GB Pant University of Agriculture and Technology, Pantnagar-263145 (Uttarakhand).
- Rana, J.S. and K.K. Dahiya. 2001. Management of termite, *Microtermes obesi* (Holmgren) in wheat, *Triticum aestivum* through seed treatment. *Ann. Biol.*, 17(2): 207-210.
- Rashid, M., A.S. Garjan, B. Naseri and F. Saberfar. 2012. Comparative toxicity of five insecticides against subterranean termite, *Amitermes vilis* (Isoptera: Termitidae) under laboratory conditions. *Mun. Ent. Zool.*, 7(2): 1044-1050.
- Rouland-Lefèvre, C., 2010. Termites as pests of agriculture. In: *Biology of termites: A modern synthesis*. Springer, Dordrecht. pp. 499-517. https://doi.org/10.1007/978-90-481-3977-4_18
- Rust, M.K. and R.K. Saran. 2006. Toxicity, repellency, and transfer of chlorfenapyr against western subterranean termites (Isoptera: Rhinotermitidae). *J. Econ. Ent.*, 99(3): 864-872. <https://doi.org/10.1093/jee/99.3.864>
- Samarasinghe, M., B.S. Chhillar and R. Singh. 2007. Insecticidal properties of methanolic extract of *Allium sativum* L. and its fractions against *Plutella xylostella* (L.). *Pestic. Res. J.*, 19(2): 145-148.
- Sapkota, R., 2018. Residual effects of termiticides on mortality of Formosan subterranean termites, *Coptotermes formosanus* Shiraki, in substrates subjected to flooding. LSU Master Dissertation. https://digitalcommons.lsu.edu/gradschool_theses/4761.
- Sattar, A., M. Naeem and Ehsan-ul-Haq. 2014. Efficacy of plant extracts against subterranean termites i.e. *Microtermes obesi* and *Odontotermes lokanandi* (Blattodea: Termitidae). *J. Biodiv. Biopros. Develop.*, 1: 122-128.
- Siskos, E.P., M.A. Konstantopoulou and B.E. Mazomenos. 2009. Insecticidal activity of *Citrus aurantium* peel extract against *Bactrocera oleae* and *Ceratitidis capitata* adults (Diptera: Tephritidae). *J. Appl. Ent.*, 133(2): 108-116. <https://doi.org/10.1111/j.1439-0418.2008.01312.x>
- Spomer, N.A., S.T. Kamble and B.D. Siegfried. 2009. Bioavailability of chlorantraniliprole and indoxacarb to eastern subterranean termites (Isoptera: Rhinotermitidae) in various soils. *J. Econ. Ent.*, 102(5): 1922-1927. <https://doi.org/10.1603/029.102.0524>
- Zaka, M.A., N. Hussain, G. Sarwar, M.R. Malik, I. Ahmad and K.H. Gill. 2004. Fertility status of Sargodha district soils. *Pak. J. Sci. Res.*, 56(1-2): 69-75.
- Zhao, N.N., H. Zhang, X.C. Zhang, X.B. Luan, C. Zhou, Q.Z. Liu, and Z.L. Liu. 2013. Evaluation of acute toxicity of essential oil of garlic (*Allium sativum*) and its selected major constituent compounds against overwintering *Cacopsylla chinensis* (Hemiptera: Psyllidae). *J. Econ. Ent.*, 106(3): 1349-1354. <https://doi.org/10.1603/EC12191>
- Zhu, B.C., G. Henderson, F. Chen, H. Fei and R.A. Laine. 2001. Evaluation of vetiver oil and seven insect-active essential oils against the Formosan subterranean termite. *J. Chem. Ecol.*, 27(8): 1617-1625.