



Research Article

Effect of Different Sowing Dates and Cultivars on Growth and Productivity of Mungbean Crop

Maqsood Ahmad¹, Muhammad Umer Chattha¹, Imran Khan¹, Muhammad Bilal Chattha^{2*}, Faqir Hussain Anjum³, Sadia Afzal⁴, Muhammad Faran¹, Fiaz Hussain³, Muhammad Talha Aslam¹, Abdul Jabbar⁵, Muhammad Sultan Ali Bazmi⁵, Mahnoor Mehmood⁶ and Muhammad Umair Hassan¹

¹Department of Agronomy, University of Agriculture, Faisalabad, 38040, Pakistan; ²Institute of Agricultural Sciences, University of the Punjab, Lahore, Pakistan; ³Directorate of Agronomy, Ayub Agricultural Research Institute, Faisalabad, Pakistan; ⁴Department of Chemistry, University of Agriculture, Faisalabad, 38040, Pakistan; ⁵Fodder Research Institute Sargodha, Pakistan; ⁶Department of Botany, University of Agriculture, Faisalabad, Pakistan.

Abstract | Sowing time is an important agronomic factor that significantly affects plant growth, development and final production. Similarly, suitable cultivar also plays an appreciable role in final productivity. Therefore, the present study was conducted to determine the effect of variable sowing dates and cultivars on the growth and productivity of mungbean. The study was conducted in RCBD with a split arrangement having three replications at Agronomic Research Area, University of Agriculture Faisalabad. The experiment consisted of different sowing dates (SD); 25th April, 5th May and 15th May and variable mungbean cultivars Azri Mung-2006, NM-2011 and NM-2006. The maximum plant height (52.67 cm), pod length (11.39 cm), pods/plant (22.11), seed yield (1300.22 kg ha⁻¹) and biological yield (4439.56 kg ha⁻¹) was recorded in crop sown on 25th April and minimum plant height (43.67 cm), pod length (9.92 cm), pods/plant (18.33), seed yield (1975.22 kg ha⁻¹) and biological yield (3761.67 kg ha⁻¹) was recorded in crop sown on 15th May. In the case of cultivars, NM-2006 performed remarkably well and had maximum plant height (50.56 cm), pod length (11.59 cm), pods/plant (21.56), seed yield (1279.67 kg ha⁻¹), biological yield (4373.11 kg ha⁻¹) and Azri-Mung 2006 had minimum plant height (44.78 cm), pod length (9.60 cm), pods/plant (19.22), seed yield (1029.22 kg ha⁻¹) and biological yield (3794.22 kg ha⁻¹). In conclusion, mungbean should be sown on 25th April to get the maximum production in rice-wheat cropping system (RWCS). Additionally, cultivar NM-2006 can be grown in RWCS to get maximum production.

Received | April 25, 2021; **Accepted** | May 18, 2021; **Published** | June 26, 2021

***Correspondence** | Muhammad Bilal Chattha, Institute of Agricultural Sciences, University of the Punjab, Lahore, Pakistan; **Email:** bilal1409@yahoo.com

Citation | Ahmad, M., M.U. Chattha, I. Khan, M.B. Chattha, F.H. Anjum, S. Afzal, M. Faran, F. Hussain, M.T. Aslam, A. Jabbar, M.S.A. Bazmi, M. Mehmood and M.U. Hassan. 2021. Effect of different sowing dates and cultivars on growth and productivity of mungbean crop. *Journal of Innovative Sciences*, 7(1): 190-198.

DOI | <https://dx.doi.org/10.17582/journal.jis/2021/7.1.190.198>

Keywords | Cultivars, Mungbean, Rice wheat cropping system, Seed yield, Sowing times

1. Introduction

Legume grains provide food for humans and feed for livestock, along with providing these benefits legume crops also improve soil fertility (Baddeley *et al.*, 2013). Mungbean (*Vigna radiate*) being a

leguminous crop can replenish soil fertility and avert land degradation and improves crop production and livestock rearing while sustaining the ecosystem (Chattha *et al.*, 2017a). Cultivation of such dual-purpose leguminous crops enhances the overall productivity of the agricultural system and conserves

the natural resources and helps for gaining more output from the agricultural system (Kassie, 2011).

The rice-wheat cropping system (RWCS) is a vital system globally especially in South Asia where it provides the most of the grains for consumption to meet the daily dietary needs (Timsina and Connor, 2001). As for the growing population, this system suits best producing cereals in a year to fulfill the demands of the mushrooming population. As the sustainability of this system depends upon the efficient usage of the nutrients, thus it is demanded to develop and implement soil managing techniques that would enhance the soil organic matter along with increasing the activity of the microbes and retain the productivity of the soils in a sustainable way (Ali *et al.*, 2012). The inclusion of legumes as green manure crops in prevailing RWCS can enhance soil fertility in a sustainable manner (Ali *et al.*, 2012).

Among different reasons for the lower production of pulses in Pakistan, seeding time and plant population have special importance. In the changing scenario of different abiotic and biotic stress, the managerial practices must be optimized for ensuring the better crop production (Hassan *et al.*, 2020a, b). Agro-ecological conditions play a vital role in the determination of planting time. Similarly, optimum sowing time in mungbean may vary from variety to variety (Sarkar *et al.*, 2004). Planting time is of paramount importance and it has a significant effect on growth, development and yield (Asghar *et al.*, 2006; Aslam *et al.*, 2015; Hassan *et al.*, 2020c; Mohsin *et al.*, 2021). The planting time significantly varies among cultivars, therefore proper planting times should be adopted in order to get higher productivity (Aslam *et al.*, 2000).

The early sowing enhances final yield and biomass production owing to an increase in grain weight and other production traits (Barros *et al.*, 2004; Hassan *et al.*, 2020). The decrease in the growth cycle due to late sowing reduced the interception of radiations that decreases the accumulation of total dry matter and consequently leads to poor yield (Vega and Hall, 2002). Selection of the cultivar and appropriate sowing date is very essential to attain yield (Jan *et al.*, 2002). Various varieties of mungbean respond differently to sowing dates and growing season. Thus, for different varieties of mungbean there should be varied optimum sowing dates (Reddy, 2009). After identifying high

yielding cultivars, sowing at optimum dates can result in higher yields (Ali and Gupta, 2012). The variable authors reported the significant differences in the seed yield of mungbean with variable planting times. The sowing of the crop from 15th April to 15th May produced the maximum yield compared to crop sown before 15th April and after 15th May (Yan-sheng *et al.*, 2010). The late sowing after 15th May results in less dry matter production, smaller pods with lower seeds, and poor yield as compared to sowing before 15th May (Yoldas and Esiyok, 2007). We hypothesized that variable sowing dates and cultivars can have differential responses in terms of growth and seed production. Therefore, the study was conducted to determine the impact of cultivars and sowing times on the growth and productivity of mungbean crop grown in RWCS.

2. Materials and Methods

2.1 Experimental site

The current study was executed at Agronomic Research Area, University of Agriculture Faisalabad in 2018 to determine the potential of mungbean as a catch crop in RWCS. The experimental site has hot and humid semi-arid conditions (Chattha *et al.*, 2019; Zamir *et al.*, 2020) further prevailed weather conditions during the crop period are given below in Table 1. The composite samples were collected with the help of a soil auger from a depth of 0-30 cm. The soil was tested according to the procedure of Homer and Pratt (1961). The soil was sandy loam with pH 7.84, organic matter 0.84%, available nitrogen 0.041%, available phosphorus 6.63 mg kg⁻¹, and available potassium 179 mg kg⁻¹.

Table 1: The climatic conditions during the study period.

Months	Monthly average temperature (°C)	Rainfall (mm)	Relative humidity (%)
April	28.5	7.9	47.3
May	32.0	21.6	29.8
June	33.9	92.0	56.5
July	33.0	195.8	70.2

2.2 Experimental detail and crop husbandry

The experiment was conducted using a net plot size of 15 m² under randomized complete block design (RCBD) with a split plot arrangement having three replications. The study was comprised of mungbean cultivars; AZRI Mung-2006, NM-2011 and NM-

2016 and different sowing dates; 25th April, 5th May and 15th May. The soil was pulverized with cultivation followed by planking to prepare the fine seedbed. The mungbean crop was sown in 30 cm apart rows using the hand drill. Urea (46%N), di-ammonium phosphate (46%P, 18%N) and sulphate of potash (50% K) were used to apply recommended dose of NPK @ 20:40:40 per hectare. The full dose of P and K and half of N was applied as basal dose and the remaining N was applied at first irrigation. All other agronomic practices were kept the same for getting optimum yield production.

2.3 Observations

In each plot, one square meter area was marked and both broad and narrow weeds were uprooted and weighed to determine their biomass 30 and 60 days after sowing (DAS) of mungbean crop. For the determination of leaf area one meter long row in each plot was harvested and a sub-sample (10 g) of leaves was taken and leaf area was measured by leaf area meter and leaf area index was measured by the method of [Watson \(1947\)](#). Moreover, leaf samples taken for determining leaf area were oven-dried and crop growth rate was determined by methods of [Hunt \(1978\)](#). Ten plants were marked in each plot, plant height was measured and pods were counted and averaged. Likewise, ten pods were taken and their length was measured and grains were counted from each pod and the average was worked out. Complete plots of each treatment were harvested and bundles were made and weighed to determine biological yield and threshed to determine seed yield and later converted into t ha⁻¹.

2.4 Statistical analysis

The observations on growth, yield and weed traits were analyzed by Fisher's analysis of variance technique ([Steel et al., 1997](#)). Furthermore, the differences among treatment means were compared by LSD test at a 5 % level of probability.

3. Results and Discussion

3.1 Weeds biomass

The results indicated that sowing dates (SD) significantly affected broad and narrow leaves weed biomass however mungbean cultivars had a non-significant impact on weeds biomass ([Table 2](#)). The maximum broad and narrow leaves weeds biomass was noted in the crop sown on 15th May, followed by

the crop sown on the 5th May and minimum broad and narrow weeds biomass was noted in crop sown on 25th April ([Table 2](#)). The higher rainfalls in mid-May and the start of June reduced the seedling emergence of mungbean and provided the opportunity for weeds to emerge which resulted in higher weeds biomass in this sowing date. These results are the same with outcomes of [Knezev et al. \(2002\)](#) they also noticed more weed biomass in the later sown crops.

3.2 Growth attributes

The variables SD and cultivars had a significant impact on the leaf area index (LAI) and crop growth rate ([Figures 1, 2](#)). LAI and CGR increased over time and reached to maximum values at 60DAS, afterwards the LAI and CGR started declining. The maximum LAI and CGR were noticed in the crop sown on the 25th April, and the lowest LAI and CGR were recorded in the crop sown on 15th May. Amongst cultivars, NM-2006 had maximum LAI and CGR and lowest LAI and CGR were noticed for Azri-mung, 2006. After 60DAS both LAI and CGR started decreasing and minimum reduction in LAI and CGR was recorded from crop sown on 25th April and in cultivar NM-2006 ([Figures 1, 2](#)). The early sown crop faced better climatic conditions during the life cycle, and it produced longer leaves with maximum width, therefore, had more leaf area. Likewise, in early sowing higher LAI resulted in maximum light-harvesting which favored higher dry matter production and consequently led to higher CGR in crop sown on 25th April. Variations amid the cultivars for the LAI can be due the differences in the leaf length and leaf width. Cultivar NM-2006 had longer leaves therefore, had more LAI, likewise, higher LAI was responsible for the higher CGR in this cultivar compared to others. The earlier plating provides a longer growth period for the crops which ensures better light-harvesting and produced more assimilates therefore, had better LAI and CGR ([Rao et al., 2012](#); [Darany, 2016](#)). Moreover, these outcomes are the same with the finding of [Wiedenfeld and Matocha \(2010\)](#), [Hassan et al. \(2018a, b\)](#) they also found differences amid cultivars for LAI and CGR.

3.3 Yield and yield traits

The results indicated that different SD and cultivars significantly affected plant height, pod length and pods/plant ([Table 3](#)). The maximum plant height (52.67 cm), pod length (11.39 cm), pods/plant (22.11) and grains/pod (13.56) were noticed in crop sown on

Table 2: Effect of different cultivars and sowing times on broad and narrow leaf weeds grown in mungbean crop.

Cultivars	Broad leaf weeds bio-mass (g) 30 DAS	Broad leaf weeds bio-mass (g) 60 DAS	Narrow leaf weeds biomass (g) 30 DAS	Narrow leaf weeds biomass (g) 60 DAS
Azri-Mung 2006	427.00	561.22	56.89	64.00
NM-2011	432.22	569.67	59.78	67.22
NM-2006	431.11	577.00	59.11	63.67
LSD $\leq$ 0.05P	NS	NS	NS	NS
Sowing dates (SD)				
25 th April	392.11C	469.11C	47.22C	57.22B
5 th May	415.11B	580.22B	59.22B	68.00A
15 th May	483.11A	658.56A	69.33A	69.67A
LSD $\leq$ 0.05P	4.77	25.17	6.65	6.28
Cultivars $\times$ SD				
Azri-Mung 2006 $\times$ 25 th April	391.67	456.33	45.00	55.00
NM-2011 $\times$ 25 th April	397.33	464.00	48.67	58.67
NM-2006 $\times$ 25 th April	387.33	487.00	48.00	58.00
Azri-Mung 2006 $\times$ 5 th May	413.67	576.00	57.67	67.67
NM-2011 $\times$ 5 th May	416.00	586.33	60.33	70.67
NM-2006 $\times$ 5 th May	415.67	578.33	59.67	65.67
Azri-Mung 2006 $\times$ 15 th May	475.67	651.33	68.00	69.33
NM-2011 $\times$ 15 th May	483.33	658.67	70.33	72.33
NM-2006 $\times$ 15 th May	490.33	665.67	69.67	67.33
LSD $\leq$ 0.05P	NS	NS	NS	NS

Means with different letters differed at 0.05 P level. DAS: days after sowing.

Table 3: Effect of different cultivars and sowing times on yield traits of mungbean crop.

Cultivars	Plant height (cm)	Pod length (cm)	Pods per plant	Grains per pod
Azri-Mung 2006	44.78C	9.60C	19.22B	10.56C
NM-2011	48.22B	10.71B	20.11B	11.44B
NM-2006	50.56A	11.59A	21.56A	12.67A
LSD $\leq$ 0.05P	2.03	0.68	1.12	0.85
Sowing dates (SD)				
25 th April	52.67A	11.39A	22.11A	13.56A
5 th May	47.22B	10.59B	20.44B	11.11B
15 th May	43.67C	9.92C	18.33C	10.00B
LSD $\leq$ 0.05P	2.12	0.23	1.56	1.15
Cultivars $\times$ SD				
Azri-Mung 2006 $\times$ 25 th April	49.33	10.27de	21.00bc	12.33
NM-2011 $\times$ 25 th April	53.00	11.60ab	21.33b	13.67
NM-2006 $\times$ 25 th April	55.67	12.30a	24.00a	14.67
Azri-Mung 2006 $\times$ 5 th May	44.00	9.60ef	19.00cd	9.67
NM-2011 $\times$ 5 th May	47.33	10.60cde	20.67bc	11.00
NM-2006 $\times$ 5 th May	50.33	11.57abc	21.67b	12.67
Azri-Mung 2006 $\times$ 15 th May	41.00	8.93f	17.67d	9.67
NM-2011 $\times$ 15 th May	44.33	9.93def	18.33 d	9.67
NM-2006 $\times$ 15 th May	45.67	10.90bcd	19.00cd	10.67
LSD $\leq$ 0.05P	NS	1.19	1.93	NS

Means with different letters differed at 0.05 P level.




Figure 1: Effect of different sowing dates (A) and cultivars (B) on leaf area index of mungbean crop

25th April, after that 5th May and minimum plant height (43.67 cm), pod length (9.92 cm), pods/plant (18.33) and grains/pod (10) were noticed in crop sown on 15th May (Table 3). In case of cultivars maximum plant height (50.56 cm), pod length (11.59 cm), pods/plant (21.56) and grains/pod (12.67) were recorded in NM-2006, and minimum plant height (44.78 cm), pod length (9.60 cm), pods/plant (19.22) and grains/pod (10.56) were recorded in Azri-mung 2006 (Table 3). The earlier sown crop produced the taller plants due to optimum temperature and moisture conditions faced by the crop (Soomro, 2003). The cultivars also had significant impact on the plant height; cultivar NM-2006 produced the taller plant owing to vigorous growth as compared to other cultivars (Siddique *et al.*, 2006). The earlier sown crop got the longer time period for the crop growth and development and produced more assimilates thus it resulted in production of longer pods with more grains. Moreover, crop sown on 15th May faced higher

temperature and high rains during the flowering stages which led to smaller pods with less grains. The current outcomes are similar with outcomes of Mondal (2004) they also noticed that early sown crop produced longer pods. The cultivars also behaved differently for the pod length, pods/plant and grains/pod; the difference among the cultivars for the pod length, pods/plant and grains/pod can be described to differences in the genetic makeup (Chattha *et al.*, 2017b; Hassan *et al.*, 2019a, b).




Figure 2: Effect of different sowing dates (A) and cultivars (B) on crop growth rate ($\text{g m}^{-2} \text{ day}^{-1}$) of mungbean crop.

The variable SD and cultivars also significantly affected seed and biomass yield and harvest index. The maximum seed yield, biological yield and harvest index were noted in crop sown on 25th April after that 5th May and minimum seed and biological yield and

harvest were recorded from crop sown on 15th May (Table 4). Likewise, in the case of cultivars maximum seed and biological yield and harvest index were noticed in NM-2006 after that NM-2011 and minimum seed yield, biological yield and harvest index were recorded in Azri-mung 2006 (Table 4). The seed yield is the interplay of yield components. The maximum seed yield in the early sown crop can be due to maximum emergence count, longer pods with more seeds and seed weight as compared to the other SD (Khattak *et al.*, 2006; Ahmad *et al.*, 2008). Likewise, the maximum biological yield was noticed in crop sown on 25th April that can be due to maximum emergence count, longer pods with more seeds and seed weight as compared to the other SD. The cultivars also behaved differently for the seed and biological yield; NM-2006 had more seed and biological yield as compared to other cultivars owing to higher emergence count, longer pods with more seeds. Earlier Seijoon *et al.* (2000) and Singh *et al.* (2006) also noted the significant difference among cultivars for seed and biological yield owing to differences in the yield characters.

Table 4: Effect of different cultivars and sowing times on seed, biological yield and harvest index of mungbean crop.

Cultivars	Seed yield (kg ha ⁻¹)	Biological yield (kg ha ⁻¹)	Harvest index (%)
Azri-Mung 2006	1029.22C	3794.22C	26.99B
NM-2011	1130.11B	4176.33B	27.04B
NM-2006	1279.67A	4373.11A	29.25A
LSD≤0.05P	55.62	155.57	1.92
Sowing dates (SD)			
25 th April	1300.22A	4439.56A	29.33A
5 th May	1163.56B	4142.44B	28.09AB
15 th May	975.22C	3761.67C	25.87B
LSD≤0.05P	82.49	166.02	2.67
Cultivars × SD			
Azri-Mung 2006×25 th April	1195.00bc	4046.67	29.54
NM-2011×25 th April	1282.00b	4418.00	29.09
NM-2006×25 th April	1423.67a	4854.00	29.36
Azri-Mung 2006×5 th May	1076.67de	3887.67	27.74
NM-2011×5 th May	1123.00cd	4191.00	26.84
NM-2006×5 th May	1291.00b	4348.67	29.68
Azri-Mung 2006×15 th May	816.00f	3448.33	23.69
NM-2011×15 th May	985.33e	3920.00	25.19
NM-2006×15 th May	1124.33cd	3916.67	28.72
LSD≤0.05P	96.34	NS	NS

Means with different letters differed at 0.05 P level.

Conclusions and Recommendations

The results depicted the significant impact of sowing dates and cultivars on the growth and productivity of mungbean crop. The sowing on 25th April resulted in better growth, and seed yield compared to other sowing dates, however, sowing at 15th caused a significant reduction in growth and yield of mungbean. Therefore, mungbean crop can be grown from 25th April to 5th May in rice-wheat crop system to obtain maximum yield from the mungbean.

Novelty Statement

The yield potential of different mungbean cultivars under different sowing dates was not fully explored. Thus this study was performed to determine the yield potential of different mungbean cultivars under different sowing dates.

Author's Contribution

Muhammad Umer Chattha and Imran Khan: Conceived and designed the experiment.

Maqsood Ahmad: Performed the experiment.

Maqsood Ahmad, Sadia Afzal and Muhammad Faran: Data collection.

Muhammad Umer Chattha, Imran Khan, Muhammad Umair Hassan and Muhammad Talha Aslam: Writing original draft.

Muhammad Bilal Chattha, Faqir Hussain Anjum, Fiaz Hussain, Abdul Jabbar, Muhammad Sultan Ali Bazmi and Mahnoor Mehmood: Reviewing and editing.

Conflict of interest

The authors have declared no conflict of interest.

References

- Ahmad, M.S.A., Hossain, M., Ijaz, S. and Alvi, A.K., 2008. Photosynthetic performance of two mungbean (*Vigna radiata*) cultivars under lead and copper stress. *International Journal of Agriculture and Biology*, 10: 167–172.
- Almodares, A. and Darany, S.M.M., 2006. Effects of planting date and time of nitrogen application on yield and sugar content of sweet sorghum. *Journal of Environmental Biology*, 27(3): 601–605.
- Ali, R.I., Awan, T.H.M., Ahmad, A., Saleem, M.U.

- and Akhtar, M., 2012. Diversification of rice-based cropping systems to improve soil fertility, sustainable productivity and economics. *Journal of Animal and Plant Science*, 22: 108-112.
- Ali, M. and Gupta, S., 2012. Carrying capacity of Indian agriculture: Pulse crops. *Current Science*, 102: 874-881.
- Asghar, A., Chaudhry, A.M., Tanveer, A., 2006. Response of mung bean (*Vigna radiata* L. genotypes to rhizobia culture. *Pakistan Journal of Agriculture Science*, 37: 1-2.
- Aslam, M., Mahmood, I.A., Sultan, T., Ahmad, S., 2000. Inoculation approach to legume crops and their production assessment in Pakistan-A Review. *Pakistan Journal of Biological Science*, 2: 193-195. <https://doi.org/10.3923/pjbs.2000.193.195>
- Aslam, M.M., Zeeshan, M., Irum, A., Hassan, M.U., Ali, S., Hussain, R., Ramzani, P.M.A. and Rashid, M.F., 2015. Influence of seedling age and nitrogen rates on productivity of rice (*Oryza sativa* L.): A Review. *American Journal of Plant Sciences*, 6: 1361-1369. <https://doi.org/10.4236/ajps.2015.69135>
- Baddeley, J.A., Jones, S., Topp, C.F.E., Watson, C.A., Helming, J., and Stoddard, F.L., 2013. Biological nitrogen fixation (BNF) in Europe. *Legume Future Report*, 1: 5-11.
- Barros, J.F., Carvalho, M., and Basch, G., 2004. Response of sunflower (*Helianthus annuus* L.) to sowing date and plant density under Mediterranean conditions. *European Journal of Agronomy*, 21: 347-356. <https://doi.org/10.1016/j.eja.2003.10.005>
- Chattha, M.U., Hassan, M.U., Khan, I., Chattha, M.B., Ashraf, I., Ishque, W., Farooq, M.U., Usman, M. and Kharal, M., 2017a. Effect of different nitrogen and phosphorus fertilizer levels in combination with nitrogen and phosphorus solubilizing inoculants on the growth and yield of mung bean. *Pakistan Journal of Life and Social Sciences*, 15(1): 31-36.
- Chattha, M.U., Khan, I., Hassan, M.U., Nawaz, M., Chattha, M.B., Ahamd, I., Khan, N.H., Usman, M., Kharal, M. and Khan, A.U., 2017b. Agronomic appraisal of amaranth accessions under semiarid conditions of Pakistan. *Pakistan Journal of Life and Social Sciences*, 15(3): 144-149.
- Chattha, M.U., Hassan, M.U., Barbanti, L., Chattha, M.B., Khan, I., Usman, M., Ali, A. and Nawaz, M., 2019. Composted sugarcane by-product press mud cake supports wheat growth and improves soil properties. *International Journal of Plant Production*, 13(3): 241-249. <https://doi.org/10.1007/s42106-019-00051-x>
- Chattha, M.U., Hassan, M.U., Khan, I., Chattha, M.B., Aamer, M., Nawaz, M., Anjum, S.A., Ashraf, U. and Kharal, M., 2020. Impact of planting methods on biomass production, chemical composition and methane yield of sorghum cultivars. *Pakistan Journal of Agricultural Sciences*, 57(1): 43-51.
- Hassan, M.U., Chattha, M.U., Mahmood, A. and Sahi, S.T., 2018a. Performance of sorghum cultivars for biomass quality and biomethane yield grown in semi-arid area of Pakistan. *Environmental Science and Pollution Research*, 25(13): 12800-12807. <https://doi.org/10.1007/s11356-018-1575-4>
- Hassan, M.U., Chattha, M.U., Chattha, M.B., Mahmood, A. and Sahi, S.T., 2018b. Bio-methane production from sorghum elite lines under the climatic conditions of Pakistan. *Maydica*, 63(1): 1-8.
- Hassan, M.U., Chattha, M.U., Barbanti, L., Chattha, M.B., Mahmood, A., Khan, I. and Nawaz, M., 2019a. Combined cultivar and harvest time to enhance biomass and methane yield in sorghum under warm dry conditions in Pakistan. *Industrial Crops and Products*, 132: 84-91. <https://doi.org/10.1016/j.indcrop.2019.02.019>
- Hassan, M.U., Chattha, M.U., Chattha, M.B., Mahmood, A. and Sahi, S.T., 2019b. Chemical composition and methane yield of sorghum as influenced by planting methods and cultivars. *The Journal of Animal and Plant Sciences*, 29(1): 251-259.
- Hassan, M.U., Chattha, M.U., Khan, I., Chattha, M.B., Barbanti, L., Aamer, M., Iqbal, M.M., Nawaz, M., Mahmood, A., Ali, A. and Aslam, M.T., 2020a. Heat stress in cultivated plants: Nature, impact, mechanisms, and mitigation strategies. A review. *Plant Biosystems*, 155: 211-234. <https://doi.org/10.1080/11263504.2020.1727987>
- Hassan, M.U., Aamer, M., Chattha, M.U., Haiying, T., Shahzad, B., Barbanti, L., Nawaz, M., Rasheed, A., Afzal, A., Liu, Y. and Guoqin, H., 2020b. The critical role of zinc in plants facing the drought stress. *Agriculture*, 10(9): 396. <https://doi.org/10.3390/agriculture10090396>
- Hassan, M.U., Chattha, M.U., Barbanti, L.,

- Mahmood, A., Chattha, M.B., Khan, I., Mirza, S., Aziz, S.A., Nawaz, M. and Aamer, M., 2020c. Cultivar and seeding time role in sorghum to optimize biomass and methane yield under warm dry climate. *Industrial Crops and Products*, 145: 111983. <https://doi.org/10.1016/j.indcrop.2019.111983>
- Homer, D.C. and Pratt, P.F., 1961. Methods of analysis for soils, plants and waters. Davis: University of California, Davis.
- Hunt, R., 1978. *Plant growth analysis. The institute Biology's studies in Biology*. Edward Arnold (Pub) Ltd, London, 96: 8–38.
- Jan, A., F.H. Taj., Khan, I.A. and Khan, N., 2002. Effect of sowing dates on yield and yield components of Mashbean varieties. *Asian Journal of Plant Science*, 10: 622–624. <https://doi.org/10.3923/ajps.2002.622.624>
- Kassie, M., 2011. *Economic and environmental benefits of forage legume-cereal intercropping in the mixed farming system: A case study in west Gojam, Ethiopia*. Addis Ababa, Ethiopia: EDRI.
- Khattak, G.S.S., Ashraf, M., Saeed, I. and Alam, A., 2006. A new high yielding mungbean (*Vigna radiata* L.) variety "Ramzan" for the agro climatic conditions of NWFP. *Pakistan Journal of Botany*, 38: 301–309.
- Knezevic, S.Z., Evans, S.P., Blankenship, E.E., VanAcker, R.C. and Lindquist, J.L., 2002. Critical period for weed control: The concept and data analysis. *Weed Science*, 50: 773–786. [https://doi.org/10.1614/0043-1745\(2002\)050\[0773:CPFWCT\]2.0.CO;2](https://doi.org/10.1614/0043-1745(2002)050[0773:CPFWCT]2.0.CO;2)
- Mondal, M.M.A., 2004. Performance of four summer mungbean varieties at Rangpur Zone of Bangladesh. *Journal of Nuclear Agriculture and Biology*, 19: 145–149.
- Muhsin, M., Nawaz, M., Khan, I., Chattha, M.B., Khan, S., Aslam, M.T., Iqbal, M.M., Amin, M.Z., Ayub, M.A., Anwar, U. and Hassan, M.U., 2021. Efficacy of seed size to improve field performance of wheat under late sowing conditions. *Pakistan Journal of Agricultural Research*, 34(1): 247–253. <https://doi.org/10.17582/journal.pjar/2021/34.1.247.253>
- Reddy, A.A., 2009. Pulses production technology: Status and way forward. *Economic and Political Weekly*, 44: 73–80. <https://doi.org/10.2139/ssrn.1537540>
- Rao, S.S., Patil, J., Prasad, P.V.V., Reddy, D.C.S., Mishra, J.S., Umakanth, A.V., Reddy, B.V.S., and Kumar, A.A., 2012. Sweet sorghum planting effects on stalk yield and sugar quality in semi-arid tropical environment. *Agronomy Journal*, 105: 1458–1465. <https://doi.org/10.2134/agronj2013.0156>
- Sarkar, M.A.R., Kabir, M.H., Begum, M. and Salam, M.A., 2004. Yield performance of mungbean as affected by planting date, variety and plant density. *Journal of Agronomy*, 3(1): 18–24. <https://doi.org/10.3923/ja.2004.18.24>
- Seijoon, P., Wookhan, K. and Rakehun, S., 2000. Influence of different planting times on harvest index and yield determination factors in soybean. *Korean Journal of Crop Science*, 45: 97–102.
- Siddique, M., Malik, M.F.A. and Awan, S.I., 2006. Genetic divergence, association and performance evaluation of different genotypes of mungbean (*Vigna radiata*). *International Journal of Agriculture and Biology*, 8: 793–795.
- Singh, J., Mathur, N., Bohra, S., Bohra, A. and Vyas, A., 2006. Comparative performance of Mungbean (*Vigna radiata* L.) varieties under rainfed condition in Indian Thar desert. *American Journal of Agriculture and Environmental Science*, 1: 48–50.
- Soomro, N.A., 2003. Response of mungbean genotypes to different dates of sowing in kharif season under rainfed condition. *Asian Journal of Plant Science*, 2: 377–379. <https://doi.org/10.3923/ajps.2003.377.379>
- Steel, R.G.D., Torrie, J.H. and Dickey, D.A., 1997. Principles and procedures of statistics: A biometric approach, 3rd edn. McGraw Hill Book Co. Inc., New York.
- Timsina, J. and Connor, D.J., 2001. Productivity and management of rice–wheat cropping systems: issues and challenges. *Field Crops Research*, 69: 93–132. [https://doi.org/10.1016/S0378-4290\(00\)00143-X](https://doi.org/10.1016/S0378-4290(00)00143-X)
- Vega, A.J. and Hall, A.J., 2002. Effects of planting date, genotype, and their interactions on sunflower yield. *Crop Science*, 42: 1191–1201. <https://doi.org/10.2135/cropsci2002.1191>
- Watson, D.J., 1947. Comparative physiological studies in the growth of field crops. I: variation in net assimilation rate and leaf area between species and varieties, and within and between years. *Annals of Botany*, 11: 41–76. <https://doi.org/10.1093/oxfordjournals.aob.a083148>
- Wiedenfeld, B. and Matocha, J., 2010. Planting

- date, row configuration and plant population effects on growth and yield of dryland sorghum in subtropical South Texas. *Archives of Agronomy and Soil Science*, 56(1): 39-47. <https://doi.org/10.1080/03650340902922696>
- Yan-sheng, L., Qiu-ying, Z., Qing, S.G., Herbert, J. and Hashemi, A., 2010. Influence of sowing date on phonological stages, seed growth and marketable yield of four vegetable soybean cultivars in North-eastern USA. *African Journal of Agriculture Research*, 5: 2556-2562.
- Yoldas, F. and Esiyok, D., 2007. Effects of sowing dates and cultural treatments on growth, quality and yield of processing beans. *Pakistan Journal of Biological Science*, 10: 2470-2474. <https://doi.org/10.3923/pjbs.2007.2470.2474>
- Zamir, S.I., Haq, I.U., Chattha, M.U., Hassan, M.U., Khan, I. Chattha, M.B., Saeed, N., Iqbal, M.M., Ayub, M.A. and Rehman, A., 2020. Harvesting at milking stage along with urea and molasses addition improved the quality and fermentation characteristics of corn silage. *International Journal of Agriculture and Biology*, 23:253-258.