



Research Article

Management of Bitter Gourd Whitefly *Bemisia tabaci* (Genn) (Hemiptera: Aleyrodidae) Under Field Condition

Riaz Hussain¹, Adnan Ihsan^{2*}, Jawad Sarwar², Nauman Ahmad², Noor Ullah², Arbab Luqman², Muhammad Junaid² and Hizer Ali²

¹Department of Entomology, The University of Agriculture Swat, Pakistan, 19120 Pakistan; ²Department of Entomology, The University of Agriculture Peshawar, Pakistan, 25100 Pakistan; ³Department of Agricultural Chemistry and Biochemistry, The University of Agriculture Peshawar, Pakistan, 25100 Pakistan.

Abstract | Bitter gourd (*Momordica charantia* L.) is one of the most important vegetable crop of warm season. Various number of insect pests attack on this crop but whitefly (*Bemisia tabaci* Genn.) is one of them. An investigation was carried out at farmer's field in Malam jaba Swat during Kharif crop 2018 to evaluate the effectiveness of synthetic and botanical insecticides against insect pests of bitter gourd i.e Whitefly Significant differences were observed among all the treatments as compared to untreated plot. Results revealed that Flonicamid 50WG (0.3g/l) was significantly better in minimizing the whitefly population, followed by Diafenthiuron 50WP (0.6g/l), Diptrex (Trichlorofom) 80-SP at 1250 gm/ha, Thiamethoxam 25 WG (0.2g/l), Chinaberry (200g/l), Garlic (20g/l), and Neem oil (2% cont). Similarly, highest yield (1556.38 kg/ha) and cost benefit ratio (1:9) was observed in Flonicamid 50WG (0.3g/l) treated plots. From the above study, it is concluded that all the insecticides reduce the population whitefly and increases net production of the crop. But chemical insecticide Flonicamid 50WG (0.3g/l) is recommended against various sucking insect pests.

Received | May 17, 2023; **Accepted** | December 06, 2024; **Published** | December 26, 2023

***Correspondence** | Adnan Ihsan, Department of Entomology, The University of Agriculture Peshawar-Pakistan, 25100 Pakistan; **Email:** adnanihsan@aup.edu.pk

Citation | Hussain, R., A. Ihsan, J. Sarwar, N. Ahmad, N. Ullah, A. Luqman, M. Junaid and H. Ali. 2023. Management of bitter gourd whitefly *Bemisia tabaci* (Genn) (Hemiptera: Aleyrodidae) under field condition. *Pakistan Journal of Weed Science Research*, 29(4): 238-242.

DOI | <https://dx.doi.org/10.17582/journal.PJWSR/2023/29.4.238.242>

Keywords | Chemical insecticides, Botanical extracts, Bittergourd, Whitefly, sucking insect, Swat



Copyright: 2023 by the authors. Licensee ResearchersLinks Ltd, England, UK.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Introduction

Bitter gourd (*Momordica charantia* L.) is the most important vegetable crops used in pickles, fresh and dry food. It contains 32% oil in its seed and is rich sources of protein, vitamins and minerals (Tindall, 1978). It is mostly effective in controlling diabetes (Yuwai *et al.*, 1991). Many insect pest attacks on bittergourd i.e melon fruit fly (*Bactrocera cucurbitae*),

widely spread in South Asian countries and causes highly economic losses. Whitefly *Bemisia tabaci* Genn. (Hemiptera: Aleyrodidae) is another major insect pest of bitter gourd which suck the cell sap from the leaves, secrete honey dew and serves as medium for sooty mold fungus (Mustafa, 1995). It slows down the processes of photosynthesis (Berlinger *et al.*, 2002) and hence it effects the market value of the crop. It is the most destructive pest of vegetables, fruits, fibers,

ornamental crops and plantation crops in tropical and subtropical regions of the world (Oliveira *et al.*, 2001).

Whitefly can cause several plant viral diseases as the result of severe yield losses. Severe infestation may lead to complete destruction of the crop (Berlinger, 2002). Excessive use of pesticides showed resistance in insect pests (Mahrotra and Phokela, 1992) which became a serious threat to the agriculture. In current situation it is crucial to develop an eco-friendly and effective IPM strategy to manage the whitefly infestation (Zhu *et al.*, 2016). Botanical extracts are eco-friendly as compared to chemical insecticides (Hussain *et al.*, 2022a). Plant extracts proved as a better alternative to synthetic insecticides (Sithisut *et al.*, 2011). The comparative study of botanical and chemical insecticides is very important to minimize the indiscriminate use of chemical insecticides. Thus, the current study was carried out to evaluate the effectiveness of botanical and chemical insecticides against whitefly on bitter melon crop under field condition.

Materials and Methods

The experiment was conducted in the farmer's field at Malam Jaba Swat during kharif crop 2018 to test the efficacy synthetic and botanical insecticides against insect pest of bitter melon.

The experiment was laid out in a Randomized Complete Block design (RCBD) having eight treatments including control and replicated three times. Insecticides viz Diptrex (trichlorfon) 80-SP (1250 gm/ha), Diafenthiuron 50WP (0.6g/l), Flonicamid 50WG (0.3g/l), Thiamethoxam 25 WG (0.2g/l) and Neem oil were purchased from local market. Extract of Garlic bulbs were obtained from field by crushing 20g garlic bulbs. The crushed garlic bulbs were then added to 1 liter of water with minute quantity of detergent and kept for 24 hours to get 2% solution (Khan *et al.*, 2022; Ali *et al.*, 2022). The chinaberry extracts were obtained by crushing 200gm of dry fruit and mixed with 1 liter of water at room temperature for 48 hrs. The solution was then filtered for field application (Hussain *et al.*, 2022a, b, d).

Field application

The insecticides were applied with 10 days intervals by using knapsack hand sprayer. The data were recorded from randomly selected 5 plants. In each plant, 6 leaves (top, mid and bottom) were randomly selected.

The data were recorded before and after 24, 48, 72 and 1 week interval. Mean pest populations data were calculated by the following formula.

$$\text{Mean insect population} = \text{sum of } X/n$$

Where X is sum of value, n is number of values.

Yield (kg/ha)

The Healthy fruits weight in each picking was noted individually and the yield was calculated which was then converted into per hectare with the following formula.

$$\text{yield kg per ha} = \frac{\text{Fruit weight (kg)}}{\text{Area harvested (m}^2\text{)}} \times 10000$$

Cost benefit ratio (CBR)

The cost benefit ratio for finding best treatment was obtained by the method used by Hussain *et al.* (2022a).

$$C.B.R = \frac{\text{estimated net benefit}}{\text{total expenditure}}$$

Statistical analysis

All the data were subjected to statistical analysis of variance after appropriate transformation given by Gomez and Gomez (1976).

Results and Discussion

Mean density of whitefly leaf⁻¹

The current study was carried out in bitter melon to evaluate the effectiveness of different botanical and chemical insecticides against whitefly at District Swat, Khyber Pakhtunkhwa. Before spray application the population of whitefly was nonsignificant. After spray application the chemical Ulala 50WG (flonicamid) was most effective in all the applied insecticides. The current results revealed that Ulala (flonicamid) remains active for 3 weeks. Morita *et al.* (2014) also observed the long-lasting efficiency of flonicamid in field. The chemical insecticide flonicamid gave better result in all spray applications against whitefly. This finding is similar to the finding of Hussain *et al.* (2022b).

After 1st spray application (Table 1) minimum mean density was observed in plot treated with Flonicamid and Diafenthiuron i.e., 1.92 Whitefly/leaf and 2.07 Whitefly/leaf, while highest (3.89 Whitefly/leaf)

mean density was observed in control plot.

Table 1: Means density of whitefly population on Bittergourd after each spray applications.

Treatments	1 st Appli- cation	2 nd Appli- cation	3 rd Appli- cation	Over all means
Neem oil	2.83b	2.33b	1.31b	2.16b
Garlic	2.69bc	2.25b	0.62cd	1.85c
Flonicamid	1.92d	1.22d	0.39d	1.17e
Chinaberry	2.73b	1.93bc	0.73c	1.80c
Thiamethoxam	2.28bcd	1.86bc	0.73c	1.62cd
Diafenthion	2.07cd	1.57cd	0.76c	1.47de
Diptrex	2.35bcd	1.65cd	0.67c	1.56cd
Control	3.89a	4.16a	3.11a	3.72a
LSD	0.16	0.23	0.44	0.20

Means followed by same letters within a column are nonsignificant at 5% level of significance (LSD test).

After 2nd spray application the minimum mean density was observed in plot treated with Flonicamid and Diafenthion i.e., 1.22 Whitefly/leaf and 1.57 Whitefly/leaf, while the highest mean density was observed in control.

After 3rd spray application, minimum mean density was observed in plot treated with Flonicamid and Garlic i.e., 0.39 Whitefly/leaf and 0.62 Whitefly/leaf, while the highest (3.11 Whitefly/leaf) mean density was recorded in control plot.

Minimum mean density of whitefly in all three applications was observed in plot treated with Flonicamid (1.17) followed by Diafenthion (1.47) while highest mean density of whitefly was recorded in control plot (3.72). In current study, the comparative efficacy of synthetic and botanical insecticides clearly indicated that Flonicamid gave better result as

compared to other insecticides. These agreements are close to [Zia et al. \(2022\)](#), [Ihsan et al. \(2022\)](#), [Hussain et al. \(2022b\)](#) and [Hussain et al. \(2022c\)](#) findings for controlling whiteflies.

Yield and cost benefit ratios (CBR)

[Table 2](#) showed that, Significant difference was recorded in all the treatments as compare to control plot. Flonicamid treated plot showed maximum yield (1556.38 kg/ha), followed by Garlic (1388 kg/ha), Neem oil (1385 kg/ha), Chinaberry (1353.05 kg/ha), Thiamethoxam (1330.83 kg/ha), Diafenthion (1320.27 kg/ha), Diptrex (1309.72 kg/ha) and the lowest yield was recorded in untreated plot (908 kg/ha) respectively. The study supported that maximum population of whitefly gave minimum production and vice versa. Current results are like the results of [Mehra et al. \(2018\)](#) and [Zia et al. \(2022\)](#). Highest (CBR) was recorded in plot treated with Flonicamid (1:9) followed by Chinaberry (1:8.04), Garlic (1:7.32), Diptrex (1:7.04), Diafenthion (1:6.79), Thiamethoxam (1:6.03) while the lowest CBR was noted in Neem oil (1:5.15). CBR and yield were recorded higher in synthetic insecticides as compare to botanicals. The current findings cannot be associated with the results of former investigators, as the commodity cost of control is fluctuating from region to region.

Conclusions and Recommendations

It was concluded from the study that the botanical extracts are less effective as compared to synthetic insecticides in controlling whitefly infestation. Therefore, it is recommended that chemicals insecticides are better to control against sucking insect pests. However, synthetic insecticides are hazardous to environment. Therefore, botanical insecticides are recommended for future study with different concentration.

Table 2: Cost benefit ratio of different synthetic and botanical insecticides against whitefly of bitter gourd.

Treatments	Yield (kg/ha)	Gross income	Cost of control	Return over control	Estimated benefit E= (D-C)	C:B (F=E/C)
Neem oil	1385	210500	8200.12	50400	42199.88	5.15
Garlic	1388	235500	9060.23	75400	66339.77	7.32
Flonicamid	1556.38	245800	8170.12	85700	77529.88	9.49
Chinaberry	1353.05	236000	8400.45	75900	67499.55	8.04
Thiamethoxam	1330.83	223540	9030.01	63440	54409.99	6.03
Diafenthion	1320.27	227510	8653.34	67410	58756.66	6.79
Diptrex	1309.72	227200	8345.34	67100	58754.66	7.04
Control	908	160100	--	--	--	--

Acknowledgements

The authors are thankful to Dr. Toheed Iqbal (Assistant Professor, AMK Campus, Mardan, The University of Agriculture, Peshawar, Pakistan) for providing technical support in conducting experiment and writing manuscript.

Novelty Statement

This study is novel in describing the use of different concentration of plant extracts as a substitute for conventional insecticides.

Author's Contribution

All authors listed, have made substantial, direct, and intellectual contribution to the work, and approved it for publication.

Conflict of interest

The authors have declared no conflict of interest.

References

- Ali, A., R. Hussain, H. Ullah, J. Anwar, S. A. Khan, K. Ullah, F. Ali, S. Khan, M. Ali, O. Riaz, W. Amin, F. Khan and S. Ahmad. 2022. Management of major insect pest of bitter melon (*Momordica charantia*) through different pesticides. J. Xi'an Shiyu Univ. Natl. Sci. Ed., 18(11): 403-411. <https://doi.org/10.1371/journal.pone.0286823>
- Banchio, E., G. Valladares, M. Defag, S. Palacios and C. Carpinella. 2003. Effects of *Melia azedarach* (Meliaceae) fruit extracts on the leaf miner *Liriomyza huidobrensis* (Diptera, Agromyzidae): Assessment in laboratory and field experiments. Ann. Appl. Biol., 143: 187-193. <https://doi.org/10.1111/j.1744-7348.2003.tb00285.x>
- Berlinger, M.J., R.A.J. Taylor, L. Mordechi, S. Shalhevet and I. Spharim. 2002. Efficacy of exclusion screens for preventing whitefly transmission of tomato yellow leaf curl virus of tomatoes in Israel. Bull. Entomol. Res., 92(5): 367-373. <https://doi.org/10.1079/BER2002180>
- Ghelani, M.K., B.B. Kabaria and S.K. Chhodavadia. 2014. Field efficacy of various insecticides against major sucking pests of BT cotton. J. Biopest., 7: 27-32.
- Gomez, K.A. and A.A. Gomez. 1976. Statistical

procedure for agricultural research (2nd 149 Ed). A Willey inter science Publication. New York. pp. 680.

- Hammad, A.F., N.M. Nemer, Z.K. Haw and L.T. Hanna. 2000. Responses of the sweet potato whitefly, *Bemisia tabaci*, to the chinaberry tree (*Melia azedarach* L.) and its extracts. Ann. Appl. Biol., 137: 79-88. <https://doi.org/10.1111/j.1744-7348.2000.tb00039.x>
- Hussain, R., A. Ihsan, A.A. Shah, N. Ullah, H. Iftikhar and R. Jalal. 2022a. Ecofriendly management of green pea (*Pisum sativum* L.) insect pests through plant extracts. Sarhad J. Agric., 38(4): 1405-1411. <https://doi.org/10.17582/journal.sja/2022/38.4.1405.1411>
- Hussain, R., A. Ihsan, N. Ullah, A.A. Shah, S.F. Shah, M. Usman and M.A. Khan. 2022b. Appraisal of plant extracts and chemical control against whitefly *Bemisia Tabaci*, (Genn) and its effect on associated natural enemies in round chili. Ann. Roman. Soc. Cell Biol., 26(1): 1121-1132. <https://annalsofrscb.ro/index.php/journal/article/10959>.
- Hussain, R., A. Ullah, M.A. Awan, M. K. Shah, A. Ali, S. Ali, M. A. Khan, K. Ullah, M. Salman, S. Ahmad and M. Shakur. 2022c. Bio-efficacy of botanical and chemical insecticides against whitefly (*Bemisia tabaci*, (Genn) and its effect on yield of the okra. Nat. Volatiles Essent. Oils. 9(2): 852-861.
- Hussain, R., H. Ullah., J. Anwar., F. Ali., A. Ali., I.U. Haq., S. Ahmad., M.A. Awan., A. Shakir, H. Iftikhar, K. Sana and H. Ali. 2022d. Management of Cabbage butterfly, *Pieris brassicae* through Herbal pesticides and its effect on yield. J. Xi'an Shiyu Univ. Natl. Sci. Ed., 18(11): 188-197.
- Ihsan, A., R. Hussain, J. Sarwar, N. Ullah, T. Iqbal, A. Mujeeb, H. Ahad, U. Qamar and N.U. Haq. 2022. Manipulation of synthetic and aqueous herbal extracts for the management of onion thrips, *Thrips tabaci*, LIND (Thysanoptera: Thripidae). J. Xi'an Shiyu Univ. Natl. Sci. Ed., 18(9): 696-710.
- Jazzar, C. and A.F. Hammad. 2003. The efficacy of enhanced aqueous extracts of *Melia azedarach* leaves and fruits integrated with the *Camptotylus reuteri* releases against the sweet potato whitefly nymphs. Bull. Insectol., 56(2): 269-275.
- Khan, A.A., R. Hussain, A. Ihsan, N. Ullah, A. Ali, I. Ullah, M. Amin, A.A. Khan, H. Iftikhar,

- K. Sana, R. Jalal and A.S. Zia. 2022. Relative efficacy of synthetic and botanical insecticides against pea leaf miner (*Phytomyza horticola*) on Pea crop. Nat. Volatiles Essent. Oils, 9(2): 302-309.
- Mahrotra, K.N. and A. Phokela. 1992. Pyrethroid resistance in *Heliothis armigera* Hub. V. Response of population in Punjab cotton. Pest. Res. J., 4: 59-61.
- Meghana, H., S.B. Jagginavar and N.D. Sunitha. 2018. Efficacy of Insecticides and Bio Pesticides against Sucking Insect Pests on BT Cotton. Int. J. Curr. Microbiol. App. Sci., 7(6): 2872-2883. <https://doi.org/10.20546/ijcmas.2018.706.338>
- Mehra, S., K. Rolania and M. Rathee. 2018. Management of cotton whitefly *Bemisia tabaci* (gennadius) (Hemiptera: Aleyrodidae). Ind. J. Entomol., 80(3): 1-4. <https://doi.org/10.5958/0974-8172.2018.00166.9>
- Morita, M., T. Yoneda and N. Akiyoshi. 2014. Research and development of a novel insecticide, flonicamid. J. Pest. Sci., 39(3): 179-180.
- Mustafa, G., 1995. Transmission of leaf curl virus of sucking insects. Proceedings of the National Workshop on Cotton Leaf Curl Virus, (CLCV'95), CCRI, Multan.
- Oliveira, M.R.V., T.J. Henneberry and P.K. Anderson. 2001. History, current status and collaborative research projects for *Bemisia tabaci*. Crop Prot. 20: 709-723. [https://doi.org/10.1016/S0261-2194\(01\)00108-9](https://doi.org/10.1016/S0261-2194(01)00108-9)
- Peshin, R. and A.K. Dhawan. 2009. Integrated pest management: Innovation, development Process. Springer. 1: 1-687. https://doi.org/10.1007/978-1-4020-8992-3_22
- Razaq, M., S. Anjum, M. Aslam, A. Jalal, M.A. Saleem and M.H.A. Khan. 2005. Evaluation of neonicotinoids and conventional insecticides against cotton jassid, *Amrasca devastans* and cotton whitefly, *Bemisia tabaci* on cotton. Pak. Entomol., 27(1): 75-78.
- Sarwar, M., A. Ali, N. Ahmad, M. Sattar and M. Tofique. 2005. Expediency of different botanical products intended for managing population of rice stem borers. Proc. Pak. Congr. Zool., 25: 15-23.
- Shaheen, N. 2008. Is organic farming suitable solution for Pakistan. SDPI Res. News Bull., 15(2): 78-81.
- Shannag, H.K., J.M. Al-Qudah, I.M. Makhadmeh and N.M. Freihat. 2007. Differences in growth and yield responses to *Aphis gossypii* G. between different okra (*Abelmoschus esculentus* L. Moench) varieties. Pl. Prot. Sci., 43: 109-116. <https://doi.org/10.17221/2250-PPS>
- Sithisut, D., P.G. Fields and A. Chandrapathya. 2011. Contact toxicity, feeding reduction and repellency of essential oils from three plants from the ginger family (Zingiberaceae) and their major components against *Sitophilus zeamais* and *Tribolium castaneum*. J. Stored Prod., 104: 1445-1454. <https://doi.org/10.1603/EC11050>
- Tindall, H.D., 1978. Commercial vegetable growing. Oxford University Press, London. pp. 150-151.
- Ullah, F., M.U. Mulk, A. Farid, M.Q. Saeed and S. Sattar. 2010. Population dynamics and chemical control of Onion Thrips (*Thrips tabaci*, Lindemann). Pak. J. Zool. 42(4): 401- 194 406.
- Yuwei, K.E., K.S. Rao, C. Kaluwin, G.P. Jones and D.E. Rivett. 1991. Chemical composition of *Momordica charantia* L. fruits. J. Agric. Food Chem., 39: 1762-1763. <https://doi.org/10.1021/jf00010a013>
- Zhu, F., L. Lavine, S. Neal, M. Lavine, C. Foss and D. Walsh. 2016. Insecticide Resistance and Management Strategies in Urban Ecosystems. SAAS Bull. Biochem. Biotechnol., 7(1): 1-26. <https://doi.org/10.3390/insects7010002>
- Zia, S., A. A. Usman, R. Hussain, A. Ihsan, N. Ullah, A.A. Shah, A. And M. Usman. 2022. Assessment of different modules for the control of whitefly *Bemisia tabaci* (Genn) and its effect on associated natural enemies in okra. Annals Roman. Soc. Cell Biol., 26(1): 1227-1239.