

Comparative Study of Growth, Hematology and Proximate Composition of Nile Tilapia (*Oreochromis niloticus*) in Biofloc Technology and Conventional Pond Culture System

Yousaf Jamal¹, Muhammad Zubair Anjum^{1*}, Zahir Muhammad¹, Saira Amin¹, Muhammad Irfan¹ and Muhammad Qayash Khan²

¹Department of Zoology, Wildlife and Fisheries, Faculty of Sciences, Pir Mehr Ali Shah Arid Agriculture University Rawalpindi, Pakistan.

²Department of Zoology, Abdul Wali Khan University Mardan, Khyber Pakhtunkhwa, Pakistan.

ABSTRACT

The current study was designed to evaluate the growth, hematology and proximate composition of Nile Tilapia (*Oreochromis niloticus*) cultured in super intensive Biofloc technology (BFT) and conventional pond culture systems (CPCS). Fish samples (n=5) of the same age (120 days) were collected and analyzed. Results showed significantly ($p > 0.05$) higher growth performance in BFT as compared to CPCS in terms of daily weight gain (DWG), total weight gain (TWG), specific growth rate (SGR), productivity and feed conversion ratio (FCR). Various hematological parameters i.e., white blood cells (WBC), red blood cells (RBC), hemoglobin (Hb), hematocrit (HCT), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), platelets (PLT) were found significantly ($p \leq 0.05$) higher while the results of proximate analysis i.e., crude protein (CP), crude fat (CF) and total ash (TA) were found significantly ($p \leq 0.05$) lower in BFT as compared to CPCS. Moreover, the moisture content was observed significantly higher in BFT than CPCS. The survival rate was approximately 100% in both culture systems. Overall the current study showed improved growth performance and hematology in BFT while better proximate composition of Nile tilapia reared in CPCS.

Article Information

Received 14 April 2022

Revised 20 April 2023

Accepted 06 May 2023

Available online 28 August 2023
(early access)

Published 10 January 2025

Authors' Contribution

YJ collected samples, analysed data and wrote the manuscript. MZA conceived idea, supervised the research, and revised the manuscript. ZM collected samples and revised the manuscript. SA revised the manuscript. MI did statistical analysis and guided manuscript writing.

Key words

Biofloc technology, Conventional pond, Nile tilapia, Growth, Hematology, Proximate composition

INTRODUCTION

Aquaculture is one of the fastest-growing food industries with an annual average growth rate of 5.3% during the period of 2001-2018. For the last five years, aquaculture has contributed 46% to world fish production and is estimated to grow from 82.1 million tonnes in 2018 to 109 million tonnes in 2050 (FAO, 2020). Due to the rapid growth in the world population which is expected to reach 9.5 billion in 2050 and the world food demand will increase about 70% (de Marsily and Abarca-del-Rio, 2016). Pakistan's human population increased with an

annual growth rate of 2.4% from 132 million in 1998 to 208 million in 2017 and is estimated to rise to 250.2 million by 2050. Although this growth rate is lower than 2.7% of the previous intercensal period (1981-1998) but much higher than the anticipated growth rate of 1.9% (Goujon *et al.*, 2020). Aquaculture and fisheries contribute a major role in the economy of a nation. Fisheries is a major source of nutrition, food, and occupation (Welcomme *et al.*, 2010) supporting livelihoods of about 60 million people and supplying about 17% of global protein (FAO, 2018). Fish being an important source of various proteins, deal with the "hidden hunger" i.e. health-related issues due to micronutrients deficiency (Kennedy, 2002). To fulfill the increased food demand the aquaculture sector being an important source of nutrition, employment, and food needed to be well developed (Sharifinia *et al.*, 2019). Intensification of the farming system is needed to meet the increasing food demand (Garlock *et al.*, 2020).

Pakistan fish production mainly comes from capture fisheries and aquaculture practices. Initially, the aquaculture contribution was very low (3517 million tons/ year from 1950 to 1990) but with time it increases

* Corresponding author: zubair.anjum@uaar.edu.pk
0030-9923/2025/0001-0267 \$ 9.00/00



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

significantly (5460 million tons/ year from 1991 to 2017). The highest aquaculture production contributing to total capture fisheries production was recorded as 1,53,230 million tons in the year 2017 (Shah *et al.*, 2018). Growth in Pakistan's fish production in the past two decades has come largely from aquaculture (FAO, 2018). The inland aquaculture in Pakistan is mainly relying on intensive, semi-intensive, and extensive culture systems. Although the growth rate of aquaculture production is increasing in Pakistan, but it is still far behind from the production level of other countries (Suplicy *et al.*, 2017). To increase the production level within the available resources many new technologies are being developed worldwide e.g., Biofloc, recycling aquaculture system (RAS), Aquaponics, etc. Fish farmers in Pakistan are also adopting these technologies according to their environment, financial budget, and available resources.

Challenges associated with the expansion and intensification of the aquaculture sector are aquatic animal disease (Stentiford *et al.*, 2017), degradation of the ecosystem (Reverter *et al.*, 2020), and water pollution due to high load of nitrogenous wastes (Crab *et al.*, 2007). Biofloc technology is the best solution to the problems associated with the expansion of the aquaculture industry such as its impact on the environment and the limited water and land resources (Khanjani and Sharifinia, 2020).

Biofloc technology (BFT) is a cost-effective, sustainable, and environmental friendly (due to zero or minimum water exchange) way of aquaculture to improves water quality as well as in situ production of microbial protein that reduces the feed conversion ratio (FCR) (Sgnaulin *et al.*, 2018). It is sustainable alternative for intensive aquaculture of Nile tilapia (*Oreochromis niloticus*), Pacific white shrimp (*Litopenaeus vannamei*) and polyculture of silver carp (*Hypophthalmichthys molitrix*), common carp (*Cyprinus carpio* L) and bighead carp (*Aristichthys nobilis*) (Zhao *et al.*, 2014). BFT might be the best solution for sustainable aquaculture production in culturing both detritivorous and filter-feeding species (Luo *et al.*, 2014).

Tilapia (*Oreochromis niloticus*) is the most important aquaculture fish species (FAO, 2007). After carps, tilapia is the second widely cultured fish species (Ridha, 2006) boosting world aquaculture production and is the important source of animal proteins and income (Sosa *et al.*, 2016). Tilapia is the best candidate for aquaculture due to its stress tolerance, widely accepted taste, fast growth, and easy breeding in captivity (Guerrero, 2019). Now a day's tilapia consumption has increased worldwide and is known as aquatic chicken (Fitzsimmons, 2005).

MATERIALS AND METHODS

Fish stocking

Monosex tilapia (*Oreochromis niloticus*) seed was obtained from Tawakul Tilapia Fish Hatchery Muradabad Muzaffargarh, Punjab. The fish was transported in Polythene bags with sufficient oxygen provided, to Smart Biofloc Fish Farm Peshawar and to Faisal Tilapia Fish Farm, Chakri Rawalpindi for rearing under Biofloc technology and conventional pond culture system respectively. A total of 800 fish with an initial average weight of 25 (g) was stocked in BFT circular tank of 4×1.5-meter (Height X Diameter) with 10,000-liter water capacity. The CPCS pond was stocked at the rate of 3400 fingerling/acre.

Pond preparation and feeding

The conventional pond was prepared and fertilized pre-stocking by adding cow dung 250 kg/acre, Urea 2 kg/acre, and DAP 4 kg/acre while Biofloc tank was added with 100 g probiotics 15×10^9 CFU/gram (Napro company imported for China), 10 m³ water, 2kg molasses, 100 g CaCO₃, and 10 kg raw salt. Fish was fed with commercially available 30% CP soya feed twice a day at the rate of 5% wet body weight in both BFT and CPCS.

Water management

Water parameters of the Biofloc tanks were maintained by adding fermented carbon organic (FCO). FCO was prepared by mixing probiotics (100 g) and molasses (5 kg) /200liter water and was provided with aeration for 24 h. FCO was added regularly 40% of the total feed used to maintain a C/N ratio of 10:1. The conventional pond were maintained by using AC-Pro aquaculture treatment chemical.

Growth performance

Growth performance of Nile Tilapia from both culture systems was obtained viz., specific growth rate (SGR) and daily weight gain (DWG) using standard formulas (Hopkins, 1992).

Specific growth rate (SGR) = $(\ln(W_f) - \ln(W_i)) / t$ (days)

Total weight gain (TWG) (g) = Final weight (g) - Initial weight (g)

Daily weight gain (DWG) (g/day) = $(W_f - W_i) / \text{days}$

Feed conversion ratio (FCR) and Survival rate percentage was determined by using formula given by Khanjani *et al.* (2017).

Feed conversion ratio (FCR) = total amount of dry feed consumed (g) / wet weight gain of fish (g)

Survival rate (%) = $(\text{final number of fish} / \text{initial number of fish}) \times 100$

Hematology

Neubauer haemocytometer was used to count the white blood cells (WBC), red blood cells (RBC) and platelets (PLT) in the blood of fish sampled randomly from both culture systems. Mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC) were calculated (Wintrobe, 1933). The hemoglobin (Hb) and hematocrit (HCT) was measured by using Sahli's haemometer and micro haematocrit reader (NUVE, NF-048 Turkey), respectively.

Proximate analysis

For proximate composition the fish were dissected and viscera was removed along with body scales. Fish meat was oven dried at 60 °C till constant weight and grinded. Moisture content, crude protein, crude fat and total ash were determined by following the standard procedure of AOAC (2005).

Statistical analysis

The data is expressed in terms of Mean±SD. All hematological and meat data were incorporated into SPSS software version 21 for desktop. T-test (two-tailed) was used to calculate the difference between the proximate composition of fish samples from the BFT and CPCS. Significance was set at ($p < 0.05$).

RESULTS

Growth performance

Growth performance, survival and productivity of Nile Tilapia (*O. niloticus*) in BFT and CPCS at the end of 120 days is presented in Figure 1. Survival rate was 100% in both culture systems while SGR, DWG, TWG, and productivity were significantly higher ($p \leq 0.05$) in BFT as compared to CPCS. FCR in CPCS was significantly higher ($p \leq 0.05$) than BFT.

Hematology

Hematological parameters of Nile Tilapia (*O. niloticus*) reared in biofloc technology and conventional pond culture systems (CPCS) are shown in Table I. The various hematological parameters were found significantly higher ($p \leq 0.05$) in BFT.

Proximate analysis

The proximate analysis of Nile Tilapia (*O. niloticus*) cultured in a conventional pond was significantly higher ($p \leq 0.05$) in terms of crude protein (CP), crude fat (CF), and total ash (TA) than fish cultured in a biofloc system. The moisture content was recorded higher ($p > 0.05$) in biofloc

cultured fish (Fig. 2).




Fig. 1. Growth performance of Nile tilapia (*O. niloticus*) reared in biofloc technology and conventional pond culture system. DWG (g/day) = daily weight gain; TWG(g) = Total weight gain; FCR = feed conversion ratio; SGR%, specific growth rate; ns, not different significantly; *, significantly different.




Fig. 2. Comparative proximate composition of Nile tilapia (*O. niloticus*) reared in biofloc technology and conventional pond culture system. CP, crude protein; CF, crude fat; TA, total ash.

DISCUSSION

Growth performance

Results of the present study revealed better growth performance of Nile Tilapia reared in biofloc technology as compared to conventional pond culture system.

Table I. Hematology of Nile Tilapia (*Oreochromis niloticus*) reared in biofloc technology (BFT) and conventional pond culture system (CPCS).

Parameters	Culture system	
	BFT	CPCS
WBC ($10^3/\mu\text{L}$)	185.00 ± 5.2^b	151.60 ± 6.10^a
RBC ($10^6/\mu\text{L}$)	2.44 ± 0.08^b	2.32 ± 0.14^a
Hb (g/dL)	13.80 ± 0.83^b	11.94 ± 0.48^a
HCT (%)	52.44 ± 1.18^b	41.75 ± 2.27^a
MCV (fL)	189.60 ± 7.70^b	174.20 ± 3.96^a
MCH (pg)	49.56 ± 1.69^b	46.52 ± 1.47^a
MCHC (g/dL)	31.82 ± 1.12^b	28.78 ± 0.95^a
PLT ($10^3/\mu\text{L}$)	287.20 ± 7.19^b	272.80 ± 4.08^a

Means $\pm$ SD with different alphabets are significantly different ($p \leq 0.05$). WBC ($10^3/\mu\text{L}$), white blood cells, RBC ($10^6/\mu\text{L}$), red blood cells; HGB (g/dL), hemoglobin; HCT(%), hematocrit; MCV(fL), mean corpuscular volume; MCH (pg), mean corpuscular hemoglobin, MCHC (g/dL), mean corpuscular hemoglobin concentration, PLT ($10^3/\mu\text{L}$), platelets.

These results are in accordance with various studies that reported increased growth performance of cultured animals by the presence of biofloc in culture systems (Panigrahi *et al.*, 2019; Adineh *et al.*, 2019). In support to our study Azim and Little (2008) also observed higher individual fish weight in biofloc treatment as compared to the control using a recirculating system. BFT treatments in the experiment also contributed 44-46% greater individual weight gain and net fish production than those in controls. Food conversion ratio (FCR) value was also significantly higher in the control as compared to the BFT treatment tanks. In biofloc probiotics, essential nutrients and several organic compounds are present (Najdegerami *et al.*, 2016). The presence of biofloc and the supplementary feed provides a complete diet to the cultured aquatic animals (Khanjani and Sharifinia, 2020). The increased growth performance in the BFT treatment might be in part due to the maintenance of optimum quality tank water and to the constant availability of nutritious floc (the richest source of many essential fatty acids) (Ekasari *et al.*, 2010; Toledo *et al.*, 2016). Many bioactive compounds were evaluated in biofloc such as carotenoids, chlorophylls, and phytosteroids, which are expected to contribute to better growth of cultured organisms in the biofloc system (Ju *et al.*, 2008). In contrast to our study Little *et al.* (2008) reported better growth performance in ordinary recirculating culture system as compared to that in indoor and outdoor BFT systems. Such results might be due to the biological turbidity provoked by biofloc and/or the chronic stress arising from deteriorated and fluctuated water quality.

Hematology

Hematological parameters are important to monitor health of cultured animals and for detecting abnormalities and disease etc. (Harikrishnan *et al.*, 2012). In the present study hematological parameters including WBC, RBC, Hb, HCT, MCV, MCH, MCHC and PLT were significantly higher ($p \leq 0.05$) in BFT than in the conventional pond. These results were in accordance with Zafar *et al.* (2021) who also reported increased hematological parameters of stinging catfish (*Heteropneustes fossilis*) in BFT as compared to control. Higher values of hematological parameters in *O. niloticus* reported by Gayed *et al.* (2021) with probiotics supplemented diet support the results of present study. The improved hematological parameters in BFT as compared to conventional pond may be due to the application of probiotics (Salam *et al.*, 2021; Saravanan *et al.*, 2021; Iwashita *et al.*, 2015). To the best of our knowledge data regarding the effect of BFT on hematological parameters is insufficient and contradictory. Azim and Little (2008) observed non significant difference in HCT% between control and BFT groups in Nile Tilapia. In contrast to our study Hisano *et al.* (2021) found non significant difference in hematological parameters of *O. niloticus* cultured in BFT and RAS and in another study conducted by Long *et al.* (2015) also non significant hematological parameters were found between indoor BFT with no water exchange and control with water exchange.

Proximate analysis

Proximate composition of fish meat i.e., Crude protein, crude fat, ash and moisture are the important factors to evaluate the muscle quality of edible organisms (Lie, 2001; Yang *et al.*, 2018a). Different environmental conditions lead to alteration in the proximate components (Yang *et al.*, 2018). Data regarding comparative analysis of the proximate composition of *O. niloticus* is insufficient in BFT. Numerous studies have been conducted to evaluate the proximate composition of tilapia and various other species in BFT, RAS, Floating cages. Comparative studies of the proximate composition of wild and pond-reared fish have also been conducted (Hisano *et al.*, 2021; Long *et al.*, 2015). In our study, significantly higher values of crude protein, crude fat, and total ash of *O. niloticus* were found in the conventional pond culture system as compared to fish reared in biofloc technology while moisture content was significantly low in the conventional pond. Significantly higher moisture content in BFT (72.82%) was recorded as compared to the conventional pond (67.26%). Azim and Little (2008) also found higher moisture content (71.25%) for *O. niloticus* fed with 35% crude protein along with biofloc and that of 67.83% in fish fed with 24% crude

protein feed with addition of biofloc. Zafar *et al.* (2021) reported the moisture content of 75.74% and 75.62% in BFT and control for Asian stinging catfish (*Heteropneustes fossilis*). Another study conducted by Desta *et al.* (2019) on the proximate composition of *O. niloticus* reported higher moisture content (73.62%) in semi-intensive ponds as compared to wild fish (66.77%) of Lake Hawassa.

The crude protein content in the present study was (56.00 ± 1.58) and (61.26 ± 1.58) for BFT and conventional pond, respectively. These results were in accordance to Azim and Little (2008) who reported the crude protein values of 53.41%, 55.28%, and 49.55% for *O. niloticus* cultured in indoor tanks and fed with different experimental diets i.e. 35% CP with biofloc, 35% CP without biofloc and 24% CP with biofloc respectively. Khanjani *et al.* (2021) also reported similar values of crude protein 59.23% for *O. niloticus* in Biofloc systems provided with Barley flour as a carbon source. The improved proximate parameters in pond cultured fish may be due to the foraging movement which leads to the breakdown of lipids and accumulation of protein content in the fish muscles (Yang *et al.*, 2018; Xu *et al.*, 2006, 2015). Chowdhury *et al.* (2000) and Uddin *et al.* (2001) also reported higher protein content in paddy culture systems as compared to conventional pond due to the availability of more diverse microbes and other supplementary nutrients.

Crude fat content in the current study calculated as 31.60 ± 1.95 and 36.58 ± 1.22 for BFT and conventional pond, respectively was in accordance with Khanjani *et al.* (2021) and Azim and Little (2008) who reported 25.69% and 27.83% respectively for *O. niloticus* in biofloc cultured systems.

Total ash content found in present study (7.46%) for the conventional pond was significantly higher as compared to BFT i.e., 4.52%. These results were in contrast to Desta *et al.* (2019) who reported low ash content in pond reared (1.51%) as compared to wild *O. niloticus* (1.89%).

Overall the present study showed that BFT is one of the best option to adopt for intensive fish culture to fulfill the increasing world food demand in a sustainable way. Better Growth and hematological parameters of Nile Tilapia indicating healthy environment in BFT as compared to CPCS. However better proximate composition of fish meal was recorded in conventional pond culture system but it requires more land as well as water. Further improvements in BFT is needed to produce quality fish with better proximate composition. Hybrid culture system of BFT and conventional pond may also fill the gap to attain better quality fish meat.

ACKNOWLEDGEMENT

The provision of research facility at Smart Biofloc

Fish Farm in Peshawar, KP by Mr. Salim Nawaz and the conventional fish pond at Rawalpindi, Punjab by Mr. Muhammad Faisal is greatly acknowledged.

Funding

It was a self funded research work, no funds were acquired from any organization for this study.

IRB approval

Not applicable.

Ethics statement

All the rules and regulations approved by ethical committee of Pir Mehr Ali Shah Arid Agriculture University, Rawalpindi were followed.

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Adineh, H., Naderi, M., Hamidi, M.K. and Harsij, M., 2019. Biofloc technology improves growth, innate immune responses, oxidative status, and resistance to acute stress in common carp (*Cyprinus carpio*) under high stocking density. *Fish Shellf. Immunol.*, **95**: 440-448. <https://doi.org/10.1016/j.fsi.2019.10.057>
- AOAC, 2005. *Official methods of analysis*, 18th ed. Association of Official Analytical Chemist, Washington, DC, US.
- Azim, M.E. and Little, D.C., 2008. The biofloc technology (BFT) in indoor tanks: Water quality, biofloc composition, and growth and welfare of Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, **283**: 29–35. <https://doi.org/10.1016/j.aquaculture.2008.06.036>
- Chowdhury, M.T.H., Dewan, S., Wahab, M.A., Uddin, M.J. and Thilsted, S.H., 2000. Water quality parameters of the rice fields used for rice cum fish culture. *Bangladesh J. Fish*, **23**: 25-29.
- Crab, R., Avnimelech, Y., Defoirdt, T., Bossier, P. and Verstraete, W., 2007. Nitrogen removal techniques in aquaculture for a sustainable production. *Aquaculture*, **270**: 1–14. <https://doi.org/10.1016/j.aquaculture.2007.05.006>
- de Marsily, G. and Abarca-del-Rio, R., 2016. Water and food in the twenty-first century. *Surv. Geophys.*, **37**: 503–527. <https://doi.org/10.1007/s10712-015-9335-1>
- Desta, D., Zello, G., Alemayehu, F., Estfanos, T., Zatti,

- K. and Drew, M., 2019. Proximate analysis of Nile Tilapia (*Oreochromis niloticus*), Fish fillet harvested from farmers pond and Lake Hawassa, Southern Ethiopia. *Int. J. Res. Dev. Technol.*, **11**: 94-99.
- Ekasari, J., Crab, R. and Verstraete, W., 2010. Primary nutritional content of bio-flocs cultured with different organic carbon sources and salinity. *HAYATI J. Biosci.*, **17**: 125-130. <https://doi.org/10.4308/hjb.17.3.125>
- FAO, 2007. *Nations Fish State and Production*, pp. 1970-2002. Rome.
- FAO, 2018. *The state of world fisheries and aquaculture 2018*. Meeting the sustainable development goals. Rome.
- FAO, 2020. *The state of world fisheries and aquaculture 2020*. Sustainability in action. FAO.
- Fitzsimmons, K., 2005. Tilapia culture. *Am. Fisher. Soc. Symp.*, **2005**: 563-590.
- Garlock, T., Asche, F., Anderson, J., Bjørndal, T., Kumar, G., Lorenzen, K., Ropicki, A., Smith, M.D. and Tveterås, R., 2020. A global blue revolution: Aquaculture growth across regions, species, and countries. *Rev. Fish. Sci. Aquacult.*, **28**: 107-116. <https://doi.org/10.1080/23308249.2019.1678111>
- Gayed, M.A., Elabd, H., Tageldin, M. and Abbass, A., 2021. Probiotic Zado (*Ruminococcus Flavofaciens*) boosts hematology, immune, serum proteins, and growth profiles in Nile tilapia (*Oreochromis niloticus*). *Fish Shellf. Immunol.*, Reports, **2**, 100021.
- Goujon, A., Wazir, A., Gailey, N. and Population, I., 2020. Pakistan: A population giant falling behind in its demographic transition. *Popul. Soc.*, **576**: 1-4. <https://doi.org/10.3917/popsoc.576.0001>
- Guerrero, R.D., 2019. Farmed tilapia production in the Philippines is declining: What has happened and what can be done. *Philippine J. Sci.*, **148**: xi-xv.
- Harikrishnan, R., Kim, J.S., Kim, M.C., Balasundaram, C. and Heo, M.S., 2012. Pomegranate enriched diet enhances the hematology, innate immune response, and disease resistance in olive flounder against *Philasterides dicentrarchi*. *Vet. Parasitol.*, **187**: 147-156. <https://doi.org/10.1016/j.vetpar.2011.12.006>
- Hisano, H., Barbosa, P.T.L., de Arruda Hayd, L. and Mattioli, C.C., 2021. Comparative study of growth, feed efficiency, and hematological profile of Nile tilapia fingerlings in biofloc technology and recirculating aquaculture system. *Trop. Anim. Hlth. Prod.*, **53**: 1-7. <https://doi.org/10.1007/s11250-020-02523-z>
- Hopkins, K.D., 1992. Reporting fish growth: A review of the basics. *J. World Aquacult. Soc.*, **23**: 173-179. <https://doi.org/10.1111/j.1749-7345.1992.tb00766.x>
- Iwashita, M.K.P., Nakandakare, I.B., Terhune, J.S., Wood, T. and Ranzani-Paiva, M.J.T., 2015. Dietary supplementation with *Bacillus subtilis*, *Saccharomyces cerevisiae* and *Aspergillus oryzae* enhance immunity and disease resistance against *Aeromonas hydrophila* and *Streptococcus iniae* infection in juvenile tilapia *Oreochromis niloticus*. *Fish Shellfish Immunol.*, **43**: 60-66. <https://doi.org/10.1016/j.fsi.2014.12.008>
- Ju, Z.Y., Forster, I., Conquest, L. and Dominy, W., 2008. Enhanced growth effects on shrimp (*Litopenaeus vannamei*) from inclusion of whole shrimp feces or feces fractions to a formulated diet. *Aquacult. Nutr.*, **14**: 533-543. <https://doi.org/10.1111/j.1365-2095.2007.00559.x>
- Kennedy, G., 2002. The scourge of hidden hunger; global dimensions of micronutrient deficiencies. *Fd. Nutr. Agric.*, **32**.
- Khanjani, M.H., Alizadeh, M., Mohammadi, M. and Sarsangi, A.H., 2021. Biofloc system applied to Nile tilapia (*Oreochromis niloticus*) farming using different carbon sources: Growth performance, carcass analysis, digestive and hepatic enzyme activity. *Iran. J. Fish. Sci.*, **20**: 490-513.
- Khanjani, M.H., Sajjadi, M.M., Alizadeh, M. and Sourinejad, I., 2017. Nursery performance of Pacific white shrimp (*Litopenaeus vannamei* Boone, 1931) cultivated in a biofloc system: the effect of adding different carbon sources. *Aquacult. Res.*, **48**: 1491-1501. <https://doi.org/10.1111/are.12985>
- Khanjani, M.H. and Sharifinia, M., 2020. *Biofloc technology as a promising tool to improve aquaculture production*. pp. 1836-1850. <https://doi.org/10.1111/raq.12412>
- Lie, 2001. Flesh quality. The role of nutrition. *Aquacult. Res.*, **32**(Suppl. 1): 341-348. <https://doi.org/10.1046/j.1355-557x.2001.00026.x>
- Little, D.C., Murray, F.J., Azim, E., Leschen, W., Boyd, K., Watterson, A. and Young, J.A., 2008. Options for producing a warm-water fish in the UK: Limits to green growth? *Trends Fd. Sci. Technol.*, **19**: 255-264. <https://doi.org/10.1016/j.tifs.2007.12.003>
- Long, L., Yang, J., Li, Y., Guan, C. and Wu, F., 2015. Effect of biofloc technology on growth, digestive enzyme activity, hematology, and immune response of genetically improved farmed tilapia (*Oreochromis niloticus*). *Aquaculture*, **448**: 135-141. <https://doi.org/10.1016/j.aquaculture.2015.05.017>

- Luo, G., Gao, Q., Wang, C., Liu, W., Sun, D., Li, L. and Tan, H., 2014. Growth, digestive activity, welfare, and partial cost-effectiveness of genetically improved farmed tilapia (*Oreochromis niloticus*) cultured in a recirculating aquaculture system and an indoor biofloc system. *Aquaculture*, **422–423**: 1–7. <https://doi.org/10.1016/j.aquaculture.2013.11.023>
- Najdegerami, E.H., Bakhshi, F. and Lakani, F.B., 2016. Effects of biofloc on growth performance, digestive enzyme activities and liver histology of common carp (*Cyprinus carpio* L.) fingerlings in zero-water exchange system. *Fish Physiol. Biochem.*, **42**: 457–465. <https://doi.org/10.1007/s10695-015-0151-9>
- Official Methods of Analysis of AOAC International. 2005. *Official Method 2005.08 RE*. 18th Ed., AOAC International, Gaithersburg, MD, USA,
- Panigrahi, A., Sundram, M., Jebha, J., Otta, S.K., Bhuvaneshwari, T., Saraswathy, R. and Ravichandran, P., 2017. Biofloc based nutrient dense culture system for nursery and grow-out farming of pacific white shrimp *Penaeus vannamei*. Boone, **1931**. <https://doi.org/10.21077/ijf.2017.64.special-issue.76189-04>
- Reverter, M., Sarter, S., Caruso, D., Avarre, J.C., Combe, M., Pepey, E., Pouyaud, L., Vega-Heredía, S., de Verdál, H. and Gozlan, R.E., 2020. Aquaculture at the crossroads of global warming and antimicrobial resistance. *Nat. Commun.*, **11**: 1–8. <https://doi.org/10.1038/s41467-020-15735-6>
- Ridha, M.T., 2006. Comparative study of growth performance of three strains of Nile tilapia, *Oreochromis niloticus*, L. at two stocking densities. *Aquacult. Res.*, **37**: 172–179. <https://doi.org/10.1111/j.1365-2109.2005.01415.x>
- Salam, M.A., Islam, M.A., Paul, S.I., Rahman, M.M., Rahman, M.L., Islam, F., Rahman, A., Shaha, D.C., Alam, M.S. and Islam, T., 2021. Gut probiotic bacteria of *Barbonymus gonionotus* improve growth, hematological parameters and reproductive performances of the host. *Sci. Rep.*, **11**: <https://doi.org/10.1038/s41598-021-90158-x>
- Saravanan, K., Sivaramakrishnan, T., Praveenraj, J., Kiruba-Sankar, R., Haridas, H., Kumar, S. and Varghese, B., 2021. Effects of single and multi-strain probiotics on the growth, hemato-immunological, enzymatic activity, gut morphology and disease resistance in Rohu, *Labeo rohita*. *Aquaculture*, **540**: 736749. <https://doi.org/10.1016/j.aquaculture.2021.736749>
- Sgnaulin, T., de Mello, G.L., Thomas, M.C., Garcia, J.R.E., de Oca, G.A.R.M. and Emerenciano, M.G.C., 2018. Biofloc technology (BFT): An alternative aquaculture system for piracanjuba *Brycon orbignyanus*? *Aquaculture*, **485**: 119–123. <https://doi.org/10.1016/j.aquaculture.2017.11.043>
- Shah, S.B.H., Mu, Y., Mohsin, M., Pavase, T.R., Kalhor, M.T., Memon, A.M., Jattak, Z., Hydar, S.S. and Soomro, M.A., 2018. An economic analysis of the fisheries sector in Pakistan (1950–2013). *Indian J. Geo-Mar. Sci.*, **47**: 903–909.
- Sharifinia, M., Afshari, B.Z., Smith, W.O., Yap, C.K. and Keshavarzifard, M., 2019. Prevention is better than cure: Persian Gulf biodiversity vulnerability to the impacts of desalination plants. *Glob. Change Biol.*, **25**: 4022–4033. <https://doi.org/10.1111/gcb.14808>
- Sosa-Villalobos, C., del Refugio, C-C.M., Amaro-Espejo, I.A., Galaviz-Villa, I. and Lango-Reynoso, F., 2016. Diagnosis of the current state of aquaculture production systems with regard to the environment in Mexico. *Latin Am. J. aquat. Res.*, **44**: 193–201. <https://doi.org/10.3856/vol44-issue2-fulltext-1>
- Stentiford, G.D., Sritunyaluksana, K., Flegel, T.W., Williams, B.A.P., Withyachumnarnkul, B., Itsathitphaisarn, O. and Bass, D., 2017. New paradigms to help solve the global aquaculture disease crisis. *PLoS Pathog.*, **13**: 1–6. <https://doi.org/10.1371/journal.ppat.1006160>
- Suplicy, F.M., Vianna, L.F.D.N., Rupp, G.S., Novaes, A.L.T., Garbossa, L.H.P., de Souza, R.V., Guzenski, J., da Costa, S.W., Silva, F.M. and dos Santos, A.A., 2017. Planning and management for sustainable coastal aquaculture development in Santa Catarina State, south Brazil. *Rev. Aquacult.*, **9**: 107–124. <https://doi.org/10.1111/raq.12107>
- Toledo, T.M., Silva, B.C., Vieira, F.D.N., Mourinho, J.L.P. and Seiffert, W.Q., 2016. Effects of different dietary lipid levels and fatty acids profile in the culture of white shrimp *Litopenaeus vannamei* (Boone) in biofloc technology: Water quality, biofloc composition, growth and health. *Aquacult. Res.*, **47**: 1841–1851. <https://doi.org/10.1111/are.12642>
- Uddin, M.J., Dewan, S., Ashrafuzzaman, M. and Haque, M.M., 2001. Growth and yield of GIFT (*Oreochromis niloticus*) and Thai silver barb (*Barbodes gonionotus* Bleeker) in rice fields and their effects on the yield of rice. *Bangladesh J. Fish. Res.*, **5**: 29–35.
- Welcomme, R.L., Cowx, I.G., Coates, D., Béné, C., Funge-Smith, S., Halls, A. and Lorenzen, K., 2010. Inland capture fisheries. *Philos. Trans. R. Soc. B Biol. Sci.*, **365**: 2881–2896. <https://doi.org/10.1098/>

- [rstb.2010.0168](#)
- Wintrobe, M.M., 1933. Variations in size and haemoglobin concentration of erythrocytes in the blood of various vertebrates. *Folia Haematol.*, **51**: 32-49.
- Xu, G.F., Wang, Y.Y., Han, Y., Liu, Y., Yang, Y.H., Yu, S.L. and Mou, Z.B., 2015. Growth, feed utilization and body composition of juvenile Manchurian trout, *Brachymystax lenok* (Pallas) fed different dietary protein and lipid levels. *Aquacult. Nutr.*, **21**: 332-340. <https://doi.org/10.1111/anu.12165>
- Xu, J., Liu, Y., Cui, S. and Miao, X., 2006. Behavioral responses of tilapia (*Oreochromis niloticus*) to acute fluctuations in dissolved oxygen levels as monitored by computer vision. *Aquacult. Eng.*, **35**: 207-217. <https://doi.org/10.1016/j.aquaeng.2006.02.004>
- Yang, H., He, J., Lv, M., Huang, G., Feng, P., Wen, L. and Hu, T., 2018. Preliminary effects of dietary protein levels on muscle quality and digestive enzyme activities in GIFT *Oreochromis niloticus*. *Aquacult. Stud.*, **18**: 1-10.
- Zafar, M.A., Talha, M.A. and Rana, M.M., 2021. Effect of biofloc technology on growth performance, digestive enzyme activity, proximate composition, and hematological parameters of Asian stinging catfish (*Heteropneustes fossilis*). *J. appl. Aquacult.*, pp. 1-19. <https://doi.org/10.1080/10454438.2021.1957053>
- Zhao, Z., Xu, Q., Luo, L., Li, J. and Wang, L., 2014. Effect of feed C/N ratio promoted bioflocs on water quality and production performance of bottom and filter feeder carp in minimum-water exchanged pond polyculture system. *Aquaculture*, **434**: 442-448. <https://doi.org/10.1016/j.aquaculture.2014.09.006>