



Research Article

Assessment of Soil Physico-Chemical Characteristics in Response to Biochar and Inorganic Fertilizers in Maize Field

Mujeebur Rahman¹, Muhammad Ilyas^{2*}, Arshad Ullah³, Rooh Ullah⁴ and Sanam Zarif²

¹Department of Soil and Water Conservation, Khyber Pakhtunkhwa, Pakistan; ²Pakistan Forest Institute, Peshawar, Khyber Pakhtunkhwa, Pakistan; ³Institute of Soil Science Chinese Academy of Sciences Nanjing, China; ⁴The University of Agriculture Peshawar, Khyber Pakhtunkhwa, Pakistan.

Abstract | A field experiment was conducted during the Kharif season, 2017 at The University of Agriculture, Peshawar, to examine the impact of biochar and inorganic fertilizers on soil physico-chemical properties in a maize field. The experiment followed a two-factor randomized complete block design (RCBD) with three replications. Factor A comprised three biochar levels: control, 5 t ha⁻¹, and 10 t ha⁻¹. Factor B included two application rates of inorganic fertilizers i.e., Nitrogen, Phosphorus and Potassium (NPK): 100% and 50% of the recommended doses. Statistical analysis revealed the application of 10 t ha⁻¹ biochar resulted in the highest values for soil organic matter, pH, electrical conductivity, total soil nitrogen, extractable phosphorus, and potassium, followed by 5 t ha⁻¹ biochar, compared to control plots. The bulk density of soil was significantly reduced with 10 t ha⁻¹ biochar. For inorganic fertilizer doses, the application of full-dose NPK, significantly enhanced total soil nitrogen, extractable phosphorus, and potassium compared to half-dose treatments. However, soil organic matter, bulk density, and pH did not show differences at the 5% significance level for NPK doses. The interaction between biochar and inorganic doses was insignificant for all soil parameters. The lack of significant interaction between biochar and NPK suggests that biochar can independently enhance soil quality, making it a promising soil amendment for sustainable agricultural practices.

Received | October 17, 2024; **Accepted** | December 14, 2025; **Published** | December 27, 2024

***Correspondence** | Muhammad Ilyas, Pakistan Forest Institute, Peshawar, Khyber Pakhtunkhwa, Pakistan; **Email:** muhammad.ilyas@pfi.edu.pk

Citation | Rahman, M., M. Ilyas, A. Ullah, R. Ullah and S. Zarif. 2024. Assessment of soil physico-chemical characteristics in response to biochar and inorganic fertilizers in maize field. *Pakistan Journal of Forestry*, 74(2): 92-100.

DOI | <https://dx.doi.org/10.17582/journal.PJF/2024/74.2.92.100>

Keywords | Soil properties, Biochar, Inorganic fertilizers, Maize crop



Copyright: 2024 by the authors. Licensee ResearchersLinks Ltd, England, UK.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Introduction

Soil quality refers to the dynamic interactions between soil and plants in growth processes. To optimize agronomic crop production, soil fertility assessment is typically carried out. Within agricultural systems, soil quality includes factors such as soil pH, electrical conductivity (EC), the

availability of essential minerals, moisture levels, soil air composition, and biotic elements. Both organic and inorganic amendments can significantly affect each of these aspects, either directly or indirectly (Mays *et al.*, 1973; Pagliai *et al.*, 1981; Tester, 1990).

Organic farming minimizes the use of artificial fertilizers. To sustain soil fertility, organic fertilizers

play a continuous role, mitigating the adverse environmental impacts resulting from extensive chemical fertilizer use (Ghafoor *et al.*, 2015). Biochar, an organic substance rich in carbon, is made by heating biomass without oxygen. It is being applied increasingly as a soil amendment to enhance physical, chemical, and biological soil attributes, decrease greenhouse gas emissions, and contribute to soil properties in beneficial ways. Beyond being a carbon source, biochar has shown potential for modifying soil characteristics, improving plant growth when used as an organic amendment. Numerous studies, including field experiments, have highlighted biochar's ability to decrease soil bulk density, as reported by Laird *et al.* (2010), with Obia *et al.* (2016) further confirming this effect in sandy loam soils. Furthermore, applying biochar at rates between 0–20 g·kg⁻¹ was found to significantly enhance the specific surface area, increasing it from 130 to 150 m²/g (Laird *et al.*, 2010).

Amending soil with biochar notably influences soil hydrological characteristics, including moisture levels, water-holding and retention capacities, and infiltration rate, which are closely related to bulk density, surface area, porosity, and aggregate stability (Laghari *et al.*, 2015). Research has shown that biochar can enhance soil aeration, increase soil pH, improve nutrient and water accessibility for crops, and shift soil microbial communities, thereby promoting plant growth and yield (Woo, 2013; Lee *et al.*, 2018; Wu *et al.*, 2020). The incorporation of biochar into soil can also change chemical properties, raising nutrient availability for plants. Biochar has been shown to enhance various soil properties, including higher levels of carbon (C), nitrogen (N), and available phosphorus (P), as well as improvements in pH, cation exchange capacity (CEC), and the availability of exchangeable cations such as calcium (Ca), magnesium (Mg), sodium (Na), and potassium (K) (Chan *et al.*, 2008). Studies have also found that biochar, derived from green waste, can raise pH and organic carbon, and increase levels of exchangeable Na, K, and Ca, while reducing exchangeable aluminum (Al) in soil (Chan *et al.*, 2008). The extent of these changes correlates with biochar application rates. Soil organic carbon (SOC) is a core indicator of soil quality, potentially enhancing degraded soils that are nutrient-poor. Recent studies have validated biochar's efficacy as an amendment to raise soil organic carbon and total nitrogen (Han *et al.*, 2016; Wang *et al.*, 2015). Biochar can either directly supply nutrients or improve nutrient accessibility

indirectly. The ash content of biochar is rich in essential nutrients, including nitrogen (N), calcium (Ca), phosphorus (P), sulfur (S), magnesium (Mg), manganese (Mn), potassium (K), iron (Fe), and zinc (Zn), which are crucial for supporting plant growth. Research by Smider and Singh (2014) demonstrated that adding biochar, with a high ash content derived from tomato waste, to soil increased corn shoot dry matter, attributed to nutrient release and liming effects.

Inorganic fertilizers are pivotal in modern agriculture, providing essential nutrients such as nitrogen, phosphorus, and potassium to enhance crop productivity. However, excessive use can lead to adverse effects, including soil acidification and increased electrical conductivity, which degrade soil quality and reduce fertility. These fertilizers are vital for sustaining crop yields but must be managed judiciously to avoid environmental consequences such as nutrient runoff and greenhouse gas emissions (Kang *et al.*, 2022). In Pakistani soils, nitrogen is 100% deficient, with phosphorus showing a 90% deficiency, and potassium tends to be deficient due to fixation (Laghari *et al.*, 2015). Relying solely on inorganic fertilizers disrupts soil surfaces by damaging structure, increase soil EC and reduce soil pH and (Kang *et al.*, 2022), worsening plant growth, soil quality, fertility, and speeding up acidification (Wu *et al.*, 2020; Wang *et al.*, 2020). Organic fertilizers provide ongoing nutrient supply, enhancing soil structure, pH, and EC (Chauhan and Bhatnagar, 2014). Despite their benefits, organic fertilizers are not preferred alone because labor-intensive and lack of quick results (Yang *et al.*, 2020).

The co-application of biochar and inorganic fertilizers synergistically improves soil physical and chemical properties. Biochar enhances soil structure and water retention, while inorganic fertilizers supply essential nutrients. This combination improves nutrient use efficiency, reduces nutrient leaching, and mitigates the adverse effects of excessive fertilizer use. Additionally, biochar's ability to increase cation exchange capacity and stabilize nutrients complements inorganic fertilizers, fostering sustainable soil fertility management (Zhang *et al.*, 2021; Kang *et al.*, 2022). Combining organic and inorganic fertilizers has proven effective for soil fertility improvement and erosion reduction (Negassa *et al.*, 2007), making it an ideal solution for enhanced soil health and crop productivity. Therefore, this study was proposed

to evaluate the soil quality parameters in response to biochar and inorganic fertilizers alone and in combination under maize crop at the research farm, The University of Agriculture, Peshawar.

Materials and Methods

Site description

During the kharif season of 2017, a field experiment was conducted to assess soil physicochemical parameters in response to biochar and inorganic fertilizers on a maize field at The University of Agriculture, Peshawar's research farm. The Azam maize variety was sown at a row spacing of 70 cm, with plants spaced 15 cm apart. Prior to seeding, the soil was amended with biochar and NPK fertilizers. All suggested cultural practices were followed throughout the growth period.

Treatments and experimental design

The experimental design consisted of a randomized complete block design (RCBD) with two treatment factors, each replicated three times. Factor A included biochar application rates (control, 5, and 10 t ha⁻¹), whereas Factor B included two levels of NPK fertilizer: 50% of the recommended dose (75, 45, 30 kg ha⁻¹) and 100% of the recommended dose (150, 90, 60 kg ha⁻¹). Nitrogen was supplied through urea, phosphorus through Di-ammonium phosphate (DAP), and potassium through sulphate of potash (SOP).

The treatment combination were following:

Treatments	Factor (A): Biochar (ton ha ⁻¹)	Factor (B): N. P. K. (kg/ha)
1	0	75-45-30, respectively
2	0	150-90-60, respectively
3	5	75-45-30, respectively
4	5	150-90-60, respectively
5	10	75-45-30, respectively
6	10	150-90-60, respectively

Soil sample collection

Pre-planting and post-harvest soil samples were collected from each plot to a depth of 0–15 cm. The samples were then packaged in labeled bags and transported to the soil science laboratory for further analysis. Twigs or stone were taken out of the samples and grinded after they had air-dried at room temperature. Following a 2 mm mesh sieve, the samples were stored for subsequent physico-chemical examination.

Soil parameter recorded

The pH of the soil was assessed using a pH meter, with a 1:5 soil-to-water suspension prepared according to McLean's (1982) method. Electrical conductivity (EC) was also measured in the same suspension using an EC meter, following the procedure described by Rhoades (1982). Soil organic matter (SOM) was assessed by oxidizing soil samples with K₂Cr₂O₇ and H₂SO₄, followed by titration with FeSO₄, according to Nelson and Sommers (1996). Bulk density was measured by drying soil samples taken from core samplers and calculating the mass per unit volume (Blake and Hartge, 1984). Total nitrogen content was measured using the Kjeldahl method (Bremner, 1996), which involved acid digestion, distillation, and titration to determine nitrogen concentration. Phosphorus (P) and potassium (K) were extracted with AB-DTPA solution; P was measured spectrophotometrically at 880 nm, while K was analysed by flame photometer, following the method of Soltanpour and Schwab (1977).

Statistical analysis

The collected data were subjected to analysis of variance (ANOVA) using MS Excel and the Statistix 8.1 software package. Treatment means were compared using the LSD test at a 0.05 probability level, as described by Jan *et al.* (2009).

Results and Discussion

A field experiment was undertaken at the research farm of The University of Agriculture, Peshawar, to investigate the impact of biochar and inorganic fertilizers on soil physico-chemical properties in a maize cropping system. The findings are detailed and critically analyzed in the subsequent sections.

Table 1: Soil physico-chemical characteristics prior to biochar and NPK application.

Properties	Unit	Concentration
Clay	%	11
Silt	%	63.4
Sand	%	25.6
Textural class	--	Silt-loam
Soil pH _(1:5)	--	7.5
Soil EC _(1:5)	dS/m	0.126
Soil organic matter	%	0.34
Bulk density	gcm ⁻³	1.45
Extra. P	mg kg ⁻¹	2.1
Extra. K	mg kg ⁻¹	48.5

Soil pH

Soil pH responses to the combined application of biochar and inorganic fertilizers are summarized in Table 2 and Figure 1. The statistical analysis indicated that biochar and NPK fertilizers had a profound effect on soil pH. The highest soil pH (7.61) was recorded at 10 t ha⁻¹ application of biochar followed by pH of 7.56 at 5 t ha⁻¹. Inorganic fertilizers application had no significant effect on soil pH and the interaction was also found non-significant.




Figure 1: Graphical representation of soil pH, influenced by biochar and NPK fertilizers under maize crop.

Soil EC (dSm⁻¹)

Soil EC responses to the combined application of biochar and inorganic fertilizers are summarized in Table 2 and Figure 2. The statistical analysis indicated that biochar and NPK fertilizers had a profound effect on soil EC. Soil EC showed a statistically significant increase with escalating biochar application rates. The maximum EC value of 0.140 dS/m was recorded in plots treated with 10 t ha⁻¹ biochar, with the 5 t ha⁻¹ treatment ranking second. Among the NPK doses the soil EC was also significantly influenced. Application of full recommended dose resulted higher EC (0.137 dS/m). The interactive effect of both factors was found non-significant for soil EC.




Figure 2: Graphical representation of soil electrical conductivity (dS/m), influenced by biochar and NPK fertilizers under maize crop.

Bulk density of soil

Soil bulk density (g cm⁻³) responses to the combined application of biochar and inorganic fertilizers are summarized in Table 2 and Figure 3. The statistical analysis indicated that biochar and NPK fertilizers had a profound effect on soil bulk density. Application of 10t ha⁻¹ biochar decreased soil bulk density (1.38 gcm⁻³), followed by 5 t ha⁻¹ over no treatment plots. Notably, the application of NPK fertilizer, with or without biochar, failed to induce significant changes in soil bulk density.




Figure 3: Graphical representation of bulk density of soil (gcm⁻³), influenced by biochar and NPK fertilizers under maize crop.

Table 2: Effect of biochar with NPK fertilizers on soil pH, EC (dS m⁻¹) and bulk density (g cm⁻³).

Treatments	pH	EC (dS m ⁻¹)	Bulk density (g cm ⁻³)
Biochar (t ha⁻¹)			
0	7.43 b	0.128 c	1.45 a
5	7.56 a	0.134 b	1.43 a
10	7.61 a	0.140 a	1.38 b
LSD (0.05)	0.005	0.0042	0.039
NPK (kg ha⁻¹)			
Half NPK (75, 45, 30)	7.52	0.132 b	1.42
Full NPK (150, 90, 60)	7.55	0.137 a	1.42
LSD (0.05)	ns	0.0034	ns
Interaction			
Biochar × NPK	ns	ns	ns

Soil organic matter (%)

Soil organic matter (%) responses to the combined application of biochar and inorganic fertilizers are summarized in Table 3 and Figure 4. The statistical analysis indicated that biochar and NPK fertilizers had a profound effect on soil organic matter. Highest organic matter (0.60%) was found in the plot, where 10t ha⁻¹ biochar were applied, and then 5t ha⁻¹ (0.56%). Inorganic fertilizer levels had no significant effect on

soil organic matter however an increasing trend were recorded from half to full NPK doses. The interaction between biochar and NPK application was found insignificant.




Figure 4: Graphical representation of soil organic matter (%), influenced by biochar and NPK fertilizers under maize crop.

Table 3: Effects of biochar and NPK fertilizer application on soil organic matter and total nitrogen contents.

Treatment	SOM (%)	Total Nitrogen (%)
Biochar (t ha⁻¹)		
0	0.54 b	0.123 c
5	0.56 ab	0.126 b
10	0.60 a	0.136 a
LSD (0.05)	0.04	0.0027
NPK (kg ha⁻¹)		
Half NPK (75, 45, 30)	0.56	0.124 b
Full NPK (150, 90, 60)	0.57	0.132 a
LSD (0.05)	ns	0.0025
Interaction		
Biochar × NPK	ns	ns

Soil total nitrogen (%)

Soil total nitrogen (%) responses to the combined application of biochar and inorganic fertilizers are summarized in Table 4 and Figure 5. The statistical analysis indicated that biochar and NPK fertilizers had a profound effect on soil total nitrogen. With increasing the biochar level total nitrogen content were increased over control plots. Highest soil nitrogen (0.136 %) was found on 10 t ha⁻¹ followed by 5 t ha⁻¹ (0.126 %). Among the inorganic fertilizer levels, maximum total soil nitrogen (0.132 %) was recorded by full recommended NPK dose (150, 90, 60 kg ha⁻¹ NPK, respectively). The interaction between biochar and NPK doses were found non-significant.




Figure 5: Graphical representation of soil total nitrogen (%), influenced by biochar and NPK fertilizers under maize crop.

Extractable phosphorus (mg kg⁻¹)

Soil extractable phosphorus (mg kg⁻¹) responses to the combined application of biochar and inorganic fertilizers are summarized in Table 4 and Figure 6. The statistical analysis indicated that biochar and NPK fertilizers had a profound effect on soil extractable phosphorus. With increasing the biochar level phosphorus content were increased over control plots. Highest AB-DTPA extractable phosphorus (2.70 mg kg⁻¹) was found on 10 t ha⁻¹ followed by 5 t ha⁻¹ (2.64 mg kg⁻¹). Among the inorganic fertilizer levels, maximum soil phosphorus (2.68 mg kg⁻¹) was recorded by full recommended NPK dose. The interactive effect between both factors were found insignificant for soil phosphorus.




Figure 6: Graphical representation of Extr. soil phosphorus (mg kg⁻¹), influenced by biochar and NPK fertilizers under maize crop.

Extractable soil potassium (mg kg⁻¹)

Soil extractable potassium (mg kg⁻¹) responses to the combined application of biochar and inorganic fertilizers are summarized in Table 4 and Figure 7. The statistical analysis indicated that biochar and NPK fertilizers had a profound effect on soil extractable potassium. With increasing the biochar

level potassium content were increased over control plots. Highest AB-DTPA extractable potassium (73.65 mg kg^{-1}) was found on 10 t ha^{-1} followed by 5 t ha^{-1} (72.45 mg kg^{-1}). Among the inorganic fertilizer levels, maximum extractable soil potassium (72.76 mg kg^{-1}) was recorded by full recommended NPK dose. The interaction between biochar and NPK doses were found non-significant.

Table 4: Effects of biochar with NPK fertilizer on extractable soil phosphorous and potassium (mg kg^{-1}).

Treatment	Phosphorous (mg kg^{-1})	Potassium (mg kg^{-1})
Biochar (t ha^{-1})		
0	2.59 b	70.51 c
5	2.64 ab	72.45 b
10	2.70 a	73.65 a
LSD (0.05)	0.09	1.07
NPK (kg ha^{-1})		
Half NPK (75, 45, 30)	2.60 b	71.64 b
Full NPK (150, 90, 60)	2.68 a	72.76 a
LSD (0.05)	0.07	0.88
Interaction		
Biochar $\times$ NPK	ns	ns




Figure 7: Graphical representation of Extr. soil potassium (mg kg^{-1}), influenced by biochar and NPK fertilizers under maize crop.

The beneficial influence of biochar on soil's physical and chemical attributes has been well-documented. The current findings correspond with [Laird et al. \(2010\)](#), who noted reductions in soil bulk density with biochar incorporation, and were later substantiated by [Obia et al. \(2016\)](#) for sandy loam soils. Similarly, [Mukherjee and Lal \(2013\)](#) concluded from their literature review that a biochar application rate of approximately 2% (w/w) is sufficient to achieve a measurable soil bulk density decrease. Similar findings were also reported by [Pandian et al. \(2016\)](#)

and [Yuan et al. \(2011\)](#), highlighted that biochar not only lowers soil bulk density but also improves moisture retention, increases soil pH, and boosts soil organic carbon content. [Lehmann and Joseph \(2009\)](#) also reported similar results and suggested that soil nutrient retention and its availability might be due to biochar application. [Chan et al. \(2008\)](#) similarly reported increases in available phosphorus, soil organic carbon, nitrogen, pH, and exchangeable potassium, along with a reduction in exchangeable aluminum, by the application of biochar from green waste. Organic carbon (SOC) is a key indicator of soil quality, particularly in degraded or nutrient-poor soils. Studies by [Han et al. \(2016\)](#) and [Wang et al. \(2015\)](#) found parallel results and reported that biochar application increases SOC and total nitrogen. Several other studies also confirmed biochar's capacity to enhance total nitrogen levels in the soil ([DeLuca et al., 2006](#); [Smider and Singh, 2014](#); [Zhao et al., 2014](#)). These findings emphasize the role of biochar in improving soil fertility, particularly through its ash content, which supplies essential plant nutrients, including nitrogen, phosphorus, potassium, sulfur, calcium, magnesium, manganese, iron, and zinc. Inorganic fertilizer also has great impact in nutrient availability alone as well with Biochar. [Fagbenro et al. \(2015\)](#) found similar results that increasing biochar and NPK levels improve nutrient availability specifically N, P and K, enhancing tree growth, stem length, and dry matter in moringa, though biochar-fertilizer interaction was non-significant. Overall, the literature highlights biochar's potential to improve various soil properties, including bulk density, moisture retention, pH, CEC, nutrient availability, and SOC. These improvements contribute to enhanced crop performance and soil health, making biochar a valuable tool for sustainable soil management and agricultural productivity.

Conclusions and Recommendations

The study concludes that integrated application of biochar and NPK fertilizers significantly improves soil physico-chemical properties. The highest biochar level (10 t ha^{-1}) showed the most pronounced effects, enhancing soil organic matter, nitrogen, phosphorus, and potassium, while also reducing bulk density. The full recommended dose of NPK fertilizers improved soil nitrogen, phosphorus, and potassium levels, though its effects on organic matter, bulk density, and pH were not statistically significant. Enhanced levels of ABDTPA phosphorus and potassium were

observed when 10 t ha⁻¹ biochar was combined with the full NPK fertilizer dose (150, 90, 60 kg ha⁻¹), indicating a positive interaction between the two treatments. Biochar at 10 t ha⁻¹ or higher can be effectively used to amend soil characteristics and improve soil health and quality.

Acknowledgement

I express my sincere gratitude to Dr. Hamida Bibi, Assistant Professor, Department of Soil and Environmental Sciences, The University of Agriculture, Peshawar, for her invaluable guidance, constructive feedback, and consistent support throughout the course of this study. I am also thankful to Prof. Dr. Jamal Khan Khattak, Chairman, Department of Soil and Environmental Sciences, The University of Agriculture, Peshawar, for his encouragement and support. Additionally, I acknowledge the support and assistance provided by Aqib Nouman Khattak, Qasim Afridi, and Arshad Ullah during the course of this research. Their encouragement and collaboration were greatly appreciated.

Novelty Statement

This study examines the independent and combined effects of biochar and inorganic fertilizers on soil physico-chemical properties in a maize cropping system under the agro-ecological conditions of Peshawar. It demonstrates that biochar has the potential to improve soil parameters both independently and in combination with inorganic fertilizers, enhancing key soil quality attributes such as organic matter content, bulk density, and nutrient availability. These findings highlight biochar as a sustainable soil amendment for improving soil health in intensive agricultural systems.

Author's Contribution

Mujeebur Rahman: Wrote original draft, resources, Methodology, Investigation, Formal analysis.

Muhammad Ilyas: Writing – review & editing, software, Resources, Investigation, Data curation, Conceptualization.

Arshad Ullah: Contributed in composition of review manuscripts and paraphrasing the literature data.

Rooh Ullah: Contributed in collecting and tabulating the data and proof reading of review manuscript.

Sanam Zarif: Contributed to the critical review and

editing of the manuscript.

Conflict of interest

The authors have declared no conflict of interest.

References

- Atkinson, C.J., Fitzgerald, J.D. and Hipps, N.A., 2010. Potential mechanisms for achieving agricultural benefits from biochar application to temperate soils: A review. *Plant Soil*, 337(1–18). <https://doi.org/10.1007/s11104-010-0464-5>
- Blake, G.R. and Hartge, K.H., 1986. Bulk density. In: A. Klute (Ed.), *Methods of Soil Analysis, Part 1—Physical and Mineralogical Methods* (2nd ed., pp. 363–382). Madison, WI: American Society of Agronomy–Soil Science Society of America.
- Bremner, J.M., 1996. Nitrogen-total. In: D.L. Sparks (Ed.), *Methods of Soil Analysis, Part 3—Chemical Methods* (pp. 1085–1122). Madison, WI: SSSA, Inc., ASA, Inc. <https://doi.org/10.2136/sssabookser5.3.c37>
- Chan, K., Van Zwieten, L., Meszaros, I., Downie, A. and Joseph, S., 2008. Agronomic values of greenwaste biochar as a soil amendment. *Soil Res.*, 45: 629. <https://doi.org/10.1071/SR07109>
- Chauhan, S.S. and Bhatnagar, R.K., 2014. Influence of long-term use of organic and inorganic manures on soil fertility and sustainable productivity of wheat in Vertisols of Madhya Pradesh. *Asian J. Soil Sci.*, 9(1): 113–116.
- Chintala, R., Mollinedo, J., Schumacher, T.E., Malo, D.D. and Julson, J.L., 2014. Effect of biochar on chemical properties of acidic soil. *Arch. Agron. Soil Sci.*, 60: 393. <https://doi.org/10.1080/03650340.2013.789870>
- DeLuca, T., MacKenzie, M., Gundale, M. and Holben, W., 2006. Wildfire-produced charcoal directly influences nitrogen cycling in ponderosa pine forests. *Soil Sci. Soc. Am. J.*, 70: 448. <https://doi.org/10.2136/sssaj2005.0096>
- Fagbenro, J.A., Oshunsanya, O.S. and Oyeleye, A.B., 2015. Effects of Gliricidia biochar and inorganic fertilizer on Moringa plant grown in an Oxisol. *Commun. Soil Sci. Plant Anal.*, 46: 619–626. <https://doi.org/10.1080/00103624.2015.1005222>
- Ghafoor, A.M.R., Ahmed, S.T. and Ahmed, M.Q., 2015. Influence of different organic fertilizers on growth and yield of wheat. *Am. Eurasian J.*

- Agric. Environ. Sci., 15(6): 1123–1126.
- Han, F., Ren, L. and Zhang, X.C., 2016. Effect of biochar on the soil nutrients about different grasslands in the Loess Plateau. *Catena*, 137: 554. <https://doi.org/10.1016/j.catena.2015.11.002>
- Jan, M.T., Shah, P., Hoointon, P.A., Khan, M.J. and Sohail, Q., 2009. Agriculture research: Design and analysis. Department of Agronomy, KPK Agriculture University, Peshawar, Pakistan.
- Fagbenro, J.A., Oshunsanya, S.O., and Oyeleye, B. A. (2015). Effect of Gliricidia biochar and inorganic fertilizer on Moringa plant grown in an Oxisol. *Communications in Soil Science and Plant Analysis*, 46(5), 620–636. <https://doi.org/10.1080/00103624.2015.1005222>
- Kang, Y.G., Chun, J.H., Lee, J.H., Park, S.Y., Luyima, D., Oh, T.K. and Yun, Y.U., 2022. Effects of varying rates of nitrogen and biochar pH on NH₃ emissions and agronomic performance of Chinese cabbage (*Brassica rapa* ssp. *pekinensis*). *Agronomy*, 12(1): 61. <https://doi.org/10.3390/agronomy12010061>
- Kang, Y.G., Lee, J.H., Chun, J.H., Yun, Y.U., Hatamleh, A.A., Dosary, M.A., Alwasel, Y.A., Lee, K.S., and Oh, T.K. 2022. Influence of individual and co-application of organic and inorganic fertilizers on NH₃ volatilization and soil quality. *J. King Saud Univ. Sci.*, 34: 102068. <https://doi.org/10.1016/j.jksus.2022.102068>
- Laghari, M., Mirjat, M.S., Hu, Z., Fazal, S., Xiao, B., Hu, M., Chen, Z. and Guo, D., 2015. Effects of biochar application rate on sandy desert soil properties and sorghum growth. *Catena*, 135: 313. <https://doi.org/10.1016/j.catena.2015.08.013>
- Laird, D., Fleming, P., Wang, B., Horton, R. and Karlen, D., 2010. Biochar impact on nutrient leaching from a Midwestern agricultural soil. *Geoderma*, 158: 436. <https://doi.org/10.1016/j.geoderma.2010.05.012>
- Lee, J.H., Seong, C.J., Kang, S.S., Lee, H.C., Kim, S.H., Lim, J.S., Kim, J.H., Yoo, J.H., Park, J.H. and Oh, T.K., 2018. Effect of different types of biochar on the growth of Chinese cabbage (*Brassica chinensis*). *Korean J. Environ. Agric.*, 45(2): 197–203.
- Laghari, S.H., Wahocho, N.A., Laghari, G.M., Laghar, A.H., Bhabhan, G.M., Talpur, K.H., Bhutto, T.A., Wahocho, S.A. and Lashari, A.A., 2016. Role of nitrogen for plant growth and development: A review. *Adv. Environ. Biol.*, 10(9): 209–218.
- Lehmann, J. and Joseph, S., 2009. Biochar for environmental management. Earthscan Publisher Ltd.
- Masto, R.E., Kumar, S., Rout, T., Sarkar, P., George, J. and Ram, L., 2013. Biochar from water hyacinth (*Eichornia crassipes*) and its impact on soil biological activity. *Catena*, 111: 64. <https://doi.org/10.1016/j.catena.2013.06.025>
- Mays, D.A., Terman, G.L. and Duggan, J.C., 1973. Municipal compost: Effects on crop yields and soil properties. *J. Environ. Qual.*, 2: 89–92. <https://doi.org/10.2134/jeq1973.00472425000200010011x>
- McLean, E.O., 1982. Soil pH and lime requirement. In: A.L. Page, R.H. Miller and D.R. Keeney (Eds.), *Methods of Soil Analysis, Part 2* (2nd ed., pp. 209–223). Agronomy.
- Mukherjee, A. and Lal, R., 2013. Biochar impacts on soil physical properties and greenhouse gas emissions. *Agron. J.*, 3: 313. <https://doi.org/10.3390/agronomy3020313>
- Negassa, W., Getaneh, F., Deressa, A. and Dinsa, B., 2007. Integrated use of organic and inorganic fertilizers for maize production. In: *Utilization of Diversity in Land Use Systems: Sustainable and Organic Approaches to Meet Human Needs*. Tropentag 2007, October 9–11, 2007, Witzenhausen.
- Nelson, D.W. and Sommer, L.E., 1996. Total carbon, organic carbon, and organic matter. In: A.L. Page, R.H. Miller and D.R. Keeney (Eds.), *Methods of Soil Analysis, Part 2* (2nd ed., pp. 574–577). Agronomy.
- Obia, A., Mulder, J., Martinsen, V., Cornelissen, G. and Børresen, T., 2016. *In situ* effects of biochar on aggregation, water retention, and porosity in light-textured tropical soils. *Soil Tillage Res.*, 155: 35. <https://doi.org/10.1016/j.still.2015.08.002>
- Pagliai, M., Guidi, G., La Marca, M., Gichetti, M. and Lucamante, G., 1981. Effect of sewage sludges and composts on soil porosity and aggregation. *J. Environ. Qual.*, 10: 556–561. <https://doi.org/10.2134/jeq1981.00472425001000040028x>
- Pandian, K., Subramaniam, P., Gnasekaran, P. and Chitraputhirapillai, S., 2016. Effect of biochar amendment on soil physical, chemical, and biological properties and groundnut yield in rainfed Alfisol of semiarid tropics. *Arch. Agron.*

- Soil Sci., 62(9): 1293–1310. <https://doi.org/10.1080/03650340.2016.1139086>
- Rhoades, J.D., 1982. Testing of soils for salinity and sodicity. In: R.L. Westerman (Ed.), soil testing and plant analysis (3rd ed., pp. 200–336).
- Smider, B. and Singh, B., 2014. Agronomic performance of a high ash biochar in two contrasting soils. *Agric. Ecosyst. Environ.*, 191: 99. <https://doi.org/10.1016/j.agee.2014.01.024>
- Soltanpour, P.N. and Schwab, A.P., 1977. A new soil test for simultaneous extraction of macro and micro nutrients in alkaline soils. *Soil Sci. Plant Anal.*, 8: 195–207. <https://doi.org/10.1080/00103627709366714>
- Tester, C.F., 1990. Organic amendment effects on physical and chemical properties of a sandy soil. *Proc. Soil Sci. Soc. Am.*, 54: 827–831. <https://doi.org/10.2136/sssaj1990.03615995005400030035x>
- Wang, J., Li, R., Zhang, H., Wei, G. and Li, Z., 2020. Beneficial bacteria activate nutrients and promote wheat growth under conditions of reduced fertilizer application. *BMC Microbiol.*, 20(1): 1–12. <https://doi.org/10.1186/s12866-020-1708-z>
- Wang, R., Kang, Y. and Wan, S., 2015. Effects of different drip irrigation regimes on saline–sodic soil nutrients and cotton yield in an arid region of Northwest China. *Agric. Water Manage.*, 153: 1. <https://doi.org/10.1016/j.agwat.2015.01.025>
- Woo, S.H., 2013. Biochar for soil carbon sequestration. *Clean Technol.*, 19(3): 201–211. <https://doi.org/10.7464/ksct.2013.19.3.201>
- Wu, L., Jiang, Y., Zhao, F., He, X., Liu, H. and Yu, K., 2020. Increased organic fertilizer application and reduced chemical fertilizer application affect the soil properties and bacterial communities of grape rhizosphere soil. *Sci. Rep.*, 10(1): 1–10. <https://doi.org/10.1038/s41598-020-66648-9>
- Yang, Q., Liu, P., Shuting, D., Zhang, J. and Zhao, B., 2020. Combined application of organic and inorganic fertilizers mitigates ammonia and nitrous oxide emissions in a maize field. *Nutr. Cycl. Agroecosyst.*, 117(1): 13–27. <https://doi.org/10.1007/s10705-020-10060-2>
- Yuan, J.H., Xu, R.K., Qian, W. and Wang, R.H., 2011. Comparison of the ameliorating effects on an acidic ultisol between four crop straws and their biochars. *J. Soils Sediments*, 11: 741. <https://doi.org/10.1007/s11368-011-0365-0>
- Zhang, Y., Wang, J. and Feng, Y., 2021. The effects of biochar addition on soil physicochemical properties: A review. *Catena*, 202: 105284. <https://doi.org/10.1016/j.catena.2021.105284>
- Zhao, X., Wang, J., Wang, S. and Xing, G., 2014. Successive straw biochar application as a strategy to sequester carbon and improve fertility: A pot experiment with two rice/wheat rotations in paddy soil. *Plant Soil*, 378: 279. <https://doi.org/10.1007/s11104-014-2025-9>