

Distributional Patterns of Carabid Beetles under Humid Subtropical Climate in Croplands of Sialkot

Somia Liaqat, Mubashar Hussain*, Kiran Aftab, Maryam Khalid and Suleman Hussain Shah

Department of Zoology, University of Gujrat, Gujrat, Punjab, Pakistan

ABSTRACT

Pakistan with its diverse climatic and topographical features, is expected to harbor a great diversity of insect species; however, insect fauna of global significance lacks taxonomic and ecological exploration in Pakistan. We explored species diversity and community composition of carabid beetles from croplands. Sampling was conducted fortnightly from August 2020 to July 2021 from district Sialkot i.e., Sialkot, Sambrial, and Pasrur. Specimens were collected by handpicking during the day while pitfall traps were used for nocturnal sampling. Diversity indices and similarities percentage (SIMPER) were calculated for all study sites. We reported 15 species belonging to four subfamilies, six tribes, and eight genera. *Calosoma inquisitor* (15.18%), *Pheropsophus verticalis* (12.37%), and *Pheropsophus darwini* (11.54%) were the most abundant species in district Sialkot. The values of diversity indices calculated for different sites show high species diversity and evenness in Sambrial ($D = 0.09$, $e^H/S = 0.94$) followed by Pasrur ($D = 0.08$, $e^H/S = 0.93$) and Sialkot ($D = 0.11$, $e^H/S = 0.84$). Species richness was highest in Pasrur ($H = 2.49$) followed by Sambrial ($H = 2.42$) and Sialkot ($H = 2.22$). *Galerita bicolor* and *Galerita lecontei veracrucis* were reported as new to Pakistan. The study emphasized that ground beetles are one of the key components of agroecosystems and need to be explored extensively concerning accelerated habitat modification and habitat types.

Article Information

Received 13 June 2023

Revised 05 September 2023

Accepted 18 September 2023

Available online 28 November 2023
(early access)

Published 02 April 2025

Authors' Contribution

MH developed the original hypothesis, designed the experiment, and finalized the manuscript. SL collected the data for this study, conducted the statistical analysis and prepared the draft. MK helped in interpreting the results. KA and SHS proofread the manuscript.

Key words

Ground beetles, Carabidae, *Calosoma*, Croplands, Carabids

INTRODUCTION

One of the key global issues is biodiversity loss in response to climate change, land-use practices, pollution, overexploitation, and agricultural intensification resulting in habitat modification, fragmentation, microclimatic variation, and habitat loss (Cardoso *et al.*, 2020; Rounsevell *et al.*, 2018; van der Meer *et al.*, 2020). Climate change and extensive agriculture are key drivers of the global decrease in insect biodiversity (Raven and Wagner, 2021). The rate of species extinction is 1000 times faster compared to the historical trends, mainly due to climate change and associated anthropogenic activities (JM *et al.*, 2015). To mitigate biodiversity decline, the conservation, and re-establishment of semi-natural habitats

and less intensively managed habitats in agricultural landscapes have been encouraged (Wang *et al.*, 2021). Several studies have identified agricultural practices and extension as principal contributing factors to species extinction (Wepprich *et al.*, 2019; Forister *et al.*, 2016; Raven and Wagner, 2021).

Several studies have been conducted to explore major insect fauna from Pakistan (Hussain and Umar, 2021; Hussain *et al.*, 2023; Ramzan *et al.*, 2021; Asghar *et al.*, 2022; Maryam *et al.*, 2020; Ane and Hussain, 2016). Major insect taxa assessed from different habitats and climatic zones during last two decades include butterflies (Roberts, 2001; Khan *et al.*, 2014, 2015, 2021; Abbas *et al.*, 2002; Bibi *et al.*, 2022; Yu-Feng *et al.*, 2020; Mangi *et al.*, 2023), dung beetles (Siddiqui *et al.*, 2014; Noreen *et al.*, 2015b; Ghazanfar *et al.*, 2017; Nasir *et al.*, 2016; Hussain *et al.*, 2020, 2022a; Abbas *et al.*, 2015; Ali *et al.*, 2015; Noreen *et al.*, 2015a, 2021), coccinellid beetles (Hussain *et al.*, 2018, 2022b; Ahmed *et al.*, 2017), grasshoppers (Hussain *et al.*, 2017; Ali and Panhwar, 2017), bees and wasps (Bashir *et al.*, 2019; Hussain *et al.*, 2010, 2012; Bashir *et al.*, 2023), and ground beetles (Rafi *et al.*, 2017; Rahim *et al.*, 2013b; Azadbakhsh and Rafi, 2017; Bibi and Rehman, 2020; Hussain *et al.*, 2022c). Coleoptera play an important role in agroecosystems due to their high diversity, abundance,

* Corresponding author: dr.mubashar@uog.edu.pk
0030-9923/2025/0002-0839 \$ 9.00/00



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

wide distribution, and role as biocontrol agents of pests (Manzoor *et al.*, 2020; Perveen, 2015; Bashir *et al.*, 2023).

Ground beetles (Carabidae) are abundant and species-rich in cultivable areas across the globe with almost 40,000 species (Lövei and Sunderland, 1996). Carabid beetles exhibit various morphological forms and diverse ecological roles i.e., predation, herbivory, and scavenging (Lövei and Sunderland, 1996). Carabid species are sensitive to anthropogenic changes in soil habitats. Thus, they act as bioindicators and are used to characterize habitat disturbance by responding to changes in vegetation height, and supplementary habitats (corridors) for generalist and mesophilous species (Lövei and Sunderland, 1996; Rainio and Niemelä, 2003; Rebrina *et al.*, 2022). Moreover, they are important pest-control agents due to their predatory polyphagous diet (Symondson *et al.*, 2002; Vichitbandha and Wise, 2002). Carabid communities may serve as an important bio-indicators as well biocontrol agents in agricultural landscapes (Avgin and Luff, 2010; Skłodowski, 2017; Schwerk *et al.*, 2020; Soomro *et al.*, 2021).

The checklist of ground beetles (Carabidae) presented earlier included 238 species from Pakistan (Kazi *et al.*, 2016). Studies on carabid beetles in Pakistan reported five species from Haripur, KPK, Pakistan (Bibi and Rehman, 2020), 38 species from National Insect Museum of Pakistan (Azadbakhsh and Rafi, 2017) 11 species from Mekran division, Baluchistan (Khatri *et al.*, 2016), 12 species from Mansehra Division, KPK (Mishkatullah, 2018), five species from Sindh Province (Soomro *et al.*, 2021), and six species of ground beetles from Upper Sindh Plains.

Keeping in view the global importance of carabid beetles, we explored the diversity of carabid beetles from Sialkot. The objective of the study is to document the carabid beetles from farmland and seminatural habitats from Sialkot, Punjab, Pakistan. The study would help to assess diversity and the pattern of ground beetles' distribution in croplands of Sialkot.

MATERIALS AND METHODS

Study area

Sialkot (32.3811° N, 74.4995° E) is one of the main districts of the province of Punjab, Pakistan. It is located in the northeast part of the province and is surrounded by Jammu Kashmir on the north, Narowal on the south, Gujranwala on the west, and Gujrat on the northwest side (Noreen *et al.*, 2020). It covers an area of 1200 square miles and is 256m above sea level. Besides the Chenab River on the north side of Sialkot, three seasonal streams also flow through the city i.e., Palkhu, Aik, and Beer. The weather of Sialkot is extremely cold during winters while hot and humid during summer. Sialkot is the sub-humid,

warm, and sub-tropical monsoon region of Pakistan (Khan, 2019). Winter temperature might fall to -2°C and the average rainfall of district Sialkot is almost 1000 mm the highest rainfall is observed from July to September (District Sialkot, 2023).

It has great agricultural potential and plays an important role in the agronomic development of Pakistan (Malik *et al.*, 2010). Sialkot district is very fertile land rich with flora. There is a variety of natural vegetation growing in different seasons throughout the year. Major plant varieties of this area include *Catharanthus roseus* (Sada bahar), and *Conyza canadensis* (Horseweed), *Cannabis sativa* (Bhang), *Ipomea carnea* (Bush morning glory), *Cucumis melo* (Wild watermelon), *Ricinus communis* (Castor oil), *Cassia fistula* (Golden shower), *Solanum nigrum* (Nightshade), *Datura inoxia* (Thorn apple), *Withania somnifera* (Winter cherry) and *Tribulus terrestris* (Puncture vine) (Arshad *et al.*, 2011).




Fig. 1. Map showing study sites in Sialkot, Pasrur, and Sambrial tehsils of district Sialkot.

Sampling procedure

Sampling was carried out fortnightly from randomly selected 15 sites within three tehsils i.e., Pasrur, Sambrial and Sialkot (Fig. 1). Sites were selected based on the vegetation type i.e., croplands and semi-natural habitats. Specimens were collected by handpicking from agricultural fields and seminatural habitats. Pitfall traps were also placed for nocturnal sampling in both type of habitats (Ullah *et al.*, 2017). We used disposable plastic cups to prepare the traps with dimensions: diameter (52mm), top diameter (81mm), and height (110 mm). One-third bottom of the trap was filled with vinegar to kill and preserve the specimen. Traps were placed in a 1km transects. Transects were separated by ~ 100 m. A total of nine traps were placed at each sampling site which were separated by

~150m. The open mouthed traps were surveyed after 24 h and the captured specimens were collected and transferred in labelled plastic containers. For the reuse of already installed traps, the killing agent was drained and the traps were refilled (Mishkatullah, 2018).

Killing, preservation, and identification

The specimens were identified by using a microscope (CZM6), labelled and preserved in 70% ethanol (Schauff, 2001; King and Porter, 2004; Post *et al.*, 1993). For identification of the specimens, taxonomic keys, field guides and literature were consulted (Abdullah and Azmir, 2021; Choate, 1999, 2001; Ghannem *et al.*, 2016; Ribera *et al.*, 1999).

Statistical analysis

The simpson and shannon-weiner diversity indices were applied using PAST3 statistical software to calculate species abundance, richness, and evenness across the study area. The similarities percentage (SIMPER) notes the species contributing to dissimilarity among different sites (Simpson, 1949; Shannon, 1949; Clarke and Gorley, 2015; Clarke, 1993; Clarke and Warwick, 1994). Seasonal relative abundance was calculated by arranging data seasonally, i.e. summer (April–October) and winter (November–March) (Hussain *et al.*, 2022a).

RESULTS

Overall diversity and abundance

The results indicated that a total of 4558 ground

beetle specimens were collected from Sialkot, Pasrur and Sambrial tehsils of district Sialkot from August 2020 to July 2021. The specimens belonged to four subfamilies, six tribes, eight genera, and fifteen species. The maximum number of species was recorded from genus *Pheropsophus* (7 species) followed by genus *Galerita* (2 species, Table I).

Overall relative abundance of species at district Sialkot indicated that *Pheropsophus hilaris sobrinus* (15.18%) was the most abundant species followed by *Pheropsophus hilaris hilaris* (12.37%), and *Pheropsophus lissoderus* (11.54%) (Fig. 2). We also indentified that *Galerita bicolor* and *Galerita lecontei veracrucis* were new to Pakistan and were being reported for the first time from this region.




Fig. 2. Overall relative abundance of carabid beetles in District Sialkot.

Table I. Relative abundance (%) of carabid beetles (family Carabidae) in different tehsils of district Sialkot.

Subfamily	Tribe	Genus	Species	Guild	Relative abundance (%)		
					Pasrur	Sialkot	Sambrial
Carabinae	Carabini	<i>Calosoma</i>	<i>Calosoma inquisitor</i>	Predator	5.81	--	5.23
Harpalinae	Chlaeniini	<i>Chlaenius</i>	<i>Chlaenius pictus</i>	Predator	8.06	--	7.52
	Lebiini	<i>Cymindis</i>	<i>Cymindis platicollis</i>	Predator	4.15	0.58	--
	Harpalini	<i>Dioryche</i>	<i>Dioryche subrecta</i>	Predator	3.08	3.19	--
	Galeritini	<i>Galerita</i>	<i>Galerita bicolor</i>	Predator	6.75	9.57	10.07
			<i>G. lecontei veracrucis</i>	Predator	5.69	6.96	5.91
Brachininae	Pterostichini	<i>Poecilus</i>	<i>Poecilus versicolor</i>	Predator	--	--	4.97
	Brachiniini	<i>Pheropsophus</i>	<i>Pheropsophus africanus</i>	Predator	9.95	11.01	7.25
			<i>P. catoirei</i>	Predator	8.77	10.43	8.72
			<i>P. darwini</i>	Predator	--	4.35	6.58
			<i>P. hilaris hilaris</i>	Predator	13.15	14.49	9.53
			<i>P. hilaris sobrinus</i>	Predator	11.97	17.83	16.38
			<i>P. lissoderus</i>	Predator	9.83	13.77	11.41
Scaritinae	Scaritini	<i>Scarites</i>	<i>P. verticalis</i>	Predator	4.98	--	6.44
			<i>Scarites subterraneus</i>	Predator	7.82	7.83	--

Table II. Diversity indices calculated for different sites from August 2020 to July 2021.

	Sialkot	Sambrial	Pasrur	Overall
Taxa_S	11	12	13	15
Individuals	1380	1490	1688	4558
Dominance_D	0.1178	0.09474	0.0878	0.09137
Simpson_1-D	0.8822	0.9053	0.9122	0.9086
Shannon_H	2.224	2.423	2.493	2.521
Evenness_e^H/S	0.8408	0.94	0.9306	0.8295
Brillouin	2.202	2.399	2.47	2.51
Menhinick	0.2961	0.3109	0.3164	0.2222
Margalef	1.383	1.506	1.615	1.662
Equitability_J	0.928	0.975	0.972	0.931
Fisher_alpha	1.632	1.783	1.917	1.931
Berger-Parker	0.1783	0.1638	0.1315	0.1518

Pheropsophus hilaris sobrinus (17.83%) showed the highest relative abundance at tehsil Sialkot, followed by *Pheropsophus hilaris hilaris* (14.49%) and *Pheropsophus lissoderus* (13.77%). The lowest relative abundance was recorded for *Cymindis platicollis* (0.58%). At tehsil Sambrial, the highest relative abundance was recorded for *Pheropsophus hilaris sobrinus* (16.38%) followed by *Pheropsophus lissoderus* (11.41%) and *Galerita bicolor* (10.07%). *Poecilus versicolor* (4.97%) showed the lowest relative abundance from tehsil Sambrial. At tehsil Pasrur,

Pheropsophus hilaris hilaris (13.15%) showed the highest relative abundance followed by *Pheropsophus hilaris sobrinus* (11.97%) and *Pheropsophus africanus* (9.95%). The lowest relative abundance was recorded for *Cymindis platicollis* (4.15%) and *Dioryche subrecta* (3.08%) (Table I).

Diversity indices

The values of species dominance (1-D) showed variations in the diversity of species at a given site (value closer to 0 indicates less diversity while 1 indicates high diversity (Table II). The Simpson's index (1-D) shows that species are more diverse in Pasrur (1-D = 0.91) followed by Sambrial (1-D = 0.90) and Sialkot (1-D = 0.88). Similarly, Shannon index (H) indicate high species richness in Pasrur (H = 2.49) followed by Sambrial (H = 2.42) and Sialkot (H = 2.2). The evenness value (e^H/S) indicated closer to 1 showed high evenness while a 0 value shows low evenness i.e., Sambrial (e^H/S = 0.94) followed by Pasrur (e^H/S = 0.93) and Sialkot (e^H/S = 0.84). The equitability value (J) indicated that the species are almost equally distributed in Sambrial (J = 0.975) and Pasrur (J = 0.972) whereas in Sialkot (J = 0.927), species are less equally distributed as compared to Sambrial and Pasrur (Table II). Margalef index explains the impact of sample size on the species richness i.e., the higher the index the greater the richness. Pasrur (1.615) exhibited higher richness as compared to sambrial (1.506) and Sialkot (1.383).

Table III. Relative abundance (%) of carabid beetles from study sites during summer and winter. Each value (%) calculated from total abundance at each site during seasons.

Species	Sialkot		Sambrial		Pasrur	
	Winter	Summer	Winter	Summer	Winter	Summer
<i>Pheropsophus africanus</i>	21.62	17.15	14.41	16.72	13.85	12.45
<i>Galerita lecontei veracruzis</i>	18.92	9.29	4.50	9.46	10.77	9.37
<i>Cymindis platicollis</i>	17.57	13.94	8.11	9.78	16.15	13.57
<i>Dioryche subrecta</i>	14.86	10.42	4.50	7.73	13.85	10.21
<i>Pheropsophus lissoderus</i>	14.86	13.46	8.11	11.99	9.23	10.49
<i>Galerita bicolor</i>	5.41	--	--	--	3.08	4.34
<i>Pheropsophus verticalis</i>	2.70	8.01	1.80	6.62	3.08	6.57
<i>Pheropsophus catoirei</i>	2.70	8.33	--	--	4.62	8.67
<i>Poecilus versicolor</i>	1.35	4.01	--	--	--	3.64
<i>Calosoma inquisitor</i>	--	--	33.33	0.32	--	0.42
<i>Chlaenius pictus</i>	--	--	9.01	7.26	12.31	7.97
<i>Pheropsophus darwini</i>	--	10.58	5.41	10.88	6.92	7.55
<i>Scarites subterraneus</i>	--	4.81	0.90	7.57	--	--
<i>Pheropsophus hilaris hilaris</i>	--	--	5.41	6.62	6.15	4.76
<i>Pheropsophus hilaris sobrinus</i>	--	--	4.50	5.05	--	--

Table IV. SIMPER results including species contributing to dissimilarities among sites.

Species	Overall dissimilarity	Mean dissimilarity	Contribution %	Cumulative %	Mean abundance (No.)	
					Sialkot	Sambrial
<i>Chlaenius pictus</i>	25.16 %	4.46	18.33	18.33	--	10.60
<i>Scarites subterraneus</i>		4.34	18.00	36.33	10.4	--
<i>Pheropsophus verticalis</i>		4.18	16.97	53.3	--	09.80
<i>Poecilus versicolor</i>		3.59	14.90	68.2	--	08.60
<i>Dioryche subrecta</i>		2.77	11.49	79.69	6.63	--
<i>Cymindis platicollis</i>		1.18	04.89	84.59	2.83	--
<i>Pheropsophus hilaris hilaris</i>		0.93	03.87	88.44	14.1	11.90
<i>Pheropsophus darwini</i>		0.90	03.73	92.17	7.75	09.90
<i>Pheropsophus africanus</i>		0.81	03.36	95.53	12.3	10.40
<i>Galerita bicolor</i>		0.32	01.32	96.84	11.5	12.20
<i>Pheropsophus lissoderus</i>		0.32	01.30	98.13	13.8	13.00
<i>Pheropsophus catoirei</i>		0.25	01.04	99.17	12.0	11.40
<i>Pheropsophus lecontei veracrucis</i>		0.18	0.72	99.89	9.80	09.38
<i>Pheropsophus hilaris sobrinus</i>		0.03	0.11	100	15.70	15.60
<i>Chlaenius pictus</i>	16.06 %	04.62	28.78	28.78	--	11.7
<i>Pheropsophus verticalis</i>		03.63	22.62	51.39	--	9.16
<i>Pheropsophus darwini</i>		03.07	19.11	70.51	7.75	--
<i>Cymindis platicollis</i>		02.19	13.67	84.17	2.83	8.37
<i>Pheropsophus hilaris sobrinus</i>		0.583	03.63	87.80	15.7	14.2
<i>Scarites subterraneus</i>		0.434	02.71	90.51	10.4	11.5
<i>Pheropsophus lissoderus</i>		0.356	02.22	92.73	13.8	12.9
<i>Galerita bicolor</i>		0.321	02.01	94.73	11.5	10.7
<i>Pheropsophus hilaris hilaris</i>		0.301	01.87	96.6	14.1	14.9
<i>Pheropsophus africanus</i>		0.251	1.56	98.16	12.3	13.0
<i>Dioryche subrecta</i>		0.230	1.43	99.59	6.63	7.21
<i>Pheropsophus catoirei</i>		0.06579	0.4096	100	12	12.2
<i>Poecilus versicolor</i>		0	0	100	0	0
<i>Galerita lecontei veracrucis</i>		0	0	100	9.8	9.8
					Sambrial	Pasrur
<i>Scarites subterraneus</i>	22.12 %	4.447	20.10	20.10	--	11.5
<i>Pheropsophus darwini</i>		3.831	17.32	37.43	9.90	--
<i>Poecilus versicolor</i>		3.329	15.05	52.48	8.60	--
<i>Cymindis platicollis</i>		3.238	14.64	67.12	--	8.37
<i>Dioryche subrecta</i>		2.791	12.62	79.74	--	7.21
<i>Pheropsophus hilaris hilaris</i>		1.155	5.222	84.96	11.9	14.9
<i>Pheropsophus africanus</i>		0.9943	4.496	89.46	10.4	13.0
<i>Galerita bicolor</i>		0.6077	2.747	92.21	12.2	10.7
<i>Pheropsophus hilaris sobrinus</i>		0.5446	2.462	94.67	15.6	14.2
<i>Chlaenius pictus</i>		0.4176	1.888	96.56	10.6	11.7
<i>Pheropsophus catoirei</i>		0.2957	1.337	97.89	11.4	12.2
<i>Pheropsophus verticalis</i>		0.245	1.108	99.00	09.8	9.16
<i>Galerita lecontei veracrucis</i>		0.1614	0.7297	99.73	9.38	9.80
<i>Pheropsophus lissoderus</i>		0.0596	0.2695	100	13.0	12.9

Species contributing to dissimilarity among different sites

SIMPER results for the species contributing to dissimilarity among different sites in district Sialkot. Overall dissimilarity among Sialkot and Sambrial is 25.16%. Top species contributing to dissimilarity include *Chlaenius pictus* (18.33%), *Scarites subterraneus* (18%), and *Pheropsophus verticalis* (16.97%) whereas *Pheropsophus hilaris sobrinus* (0.11%) has the least contribution to dissimilarity among the species. Sialkot and Pasrur showed 16.06% dissimilarity in species distribution. *Chlaenius pictus* (28.78%) and *Pheropsophus verticalis* (22.62%) have the highest contribution to dissimilarity whereas, *Poecilus versicolor* and *Galerita lecontei veracrucis* have no contribution to dissimilarity (Table IV).

Similarly, Sambrial and Pasrur have 22.12% overall dissimilarity among the species. The top species contributing to dissimilarity include *Scarites subterraneus* (20.1%) and *Pheropsophus darwini* (17.32%) whereas, *Galerita lecontei veracrucis* (0.72%) and *Pheropsophus lissoderus* (0.26%) have the least contribution to dissimilarity among the species at Sambrial and Pasrur.

Rank abundance curve

Rank abundance curve reflects evenness in the slope of the line. Whereas a steep gradient represents low evenness, while a shallow gradient is indicative of higher species evenness. The rank abundance graph of species in three study sites showed a shallow gradient of Pasrur and Sambrial which indicated evenness in the distribution of species whereas the steep curve of Sialkot indicates low evenness in species abundance (Fig. 3). Pasrur and Sambrial were equally dominant sites as compared to Sialkot. Dominant species included *Pheropsophus africanus*, *Galerita lecontei veracrucis*, *Cymindis platicollis*, *Dioryche subrecta* and *Pheropsophus lissoderus* (Table III).




Fig. 3. Rank abundance graph of species in three tehsils (Sialkot, Sambrial, and Pasrur).

DISCUSSION

Ground beetles (family Carabidae) belong to a large family of predatory ground-dwelling beetles. The studies on carabid beetles report their dependence on biotic and abiotic features of their local habitats (Pizzolotto *et al.*, 2018; Heino *et al.*, 2019). Carabid community rely on vegetation characteristics mainly through local effects (Schaffers *et al.*, 2008; Worthen and Merriman, 2013). Microclimatic conditions like food sources, temperature, moisture and sunlight influence carabid assemblages (Luff, 1980; Ernsting and Isaaks, 2000). In our study, we recorded 15 species of carabid beetles that belonged to the carnivorous guild from agricultural landscapes. Our study sites represent similarities in microclimatic conditions (temperature, light, and biotic communities), however local differences in vegetation and other features in three tehsils (Sialkot, Pasrur and Sambrial) were observed which have influenced species richness and abundance.

Ground beetle assemblages have been reported amongst important insect communities from the agricultural landscapes across the globe (Szyszko-Podgórska *et al.*, 2021; Rainio and Niemelä, 2003). Interestingly, most species of carabids were observed with reasonably higher magnitudes from all three study sites except *Scarites subterraneus* (from Sambrial), *Pheropsophus darwini* (from Pasrur), *Cymindis platicollis* (from Sambrial) and *Dioryche subrecta* (from Sambrial). *Poecilus versicolor* was not recorded from Sialkot and Pasrur. *Calosoma inquisitor*, *Chlaenius pictus*, *Pheropsophus verticalis* and *Poecilus versicolor* (also from Pasrur) were not recorded from Sialkot. Additionally, these results of ground beetle assemblages recorded from district Sialkot indicated slight variations in the species abundance, richness, and distribution across the sites i.e., Sialkot, Sambrial, and Pasrur.

Vegetation types and higher altitude and lower temperature ranges influence species diversity. Such results of lower diversity (only five species: *Carabus caschmirensis*, *Chlaenius quadricolor*, *Pheropsophus sobrinus*, *Chlaenius laticollis* and *Chlaenius hamifer*) were reported earlier from District Poonch, Azad Kashmir (Rahim *et al.*, 2013a). Surprisingly lower diversity of species and abundance of ground beetles (Coleoptera: Carabidae) from Upper Sindh Plains. Only six species were reported: *Calosoma auropunctatum*, *Calosoma sycophantum*, *Harpalus erythropus*, *Harpalus suensoni*, *Mecyclothorax cordicollis* and *Nesambyops oreobius*. Fourteen species of Brachinini tribe have been reported from northern Pakistan (Ullah *et al.*, 2017). Variations in the topography, altitudes, climatic conditions, and vegetation type determine carabid community composition.

SIMPER results highlighted that Sialkot-Sambrial comparison indicated 25.16 % contribution in the dissimilarity of species abundance. In Sialkot-Pasrur, 16.06% average dissimilarity was observed. Main species contributing *Chlaenius pictus* (18.33% in Sialkot-Sambrial, 28.78% in Sialkot-Pasrur) and *Scarites subterraneus* (20.10 % in Sambrial-Pasrur). The distribution patterns of *Chlaenius pictus* in agricultural landscapes indicated variations in the abundance in response to vegetation, temperature, and microenvironment (Satpathi, 2021; Knapp et al., 2019).

Twelve species were reported from coastal areas of Sindh with most abundant species include *Calosoma aurapunctatum*, *Calosoma scyophantum* and *Anthia sexumatta*. The species belonging to the genus *Pheropsophus* were found to be the most abundant in Sialkot. Species were found to be evenly distributed in Sambrial and Pasrur (less anthropogenic activities) as compared to Sialkot (more anthropogenic activities). Similar results have been reported for changes in the composition and abundance of carabid species due to the disturbance caused by anthropogenic activities (Rainio and Niemelä, 2003). Habitat fragmentation, vegetation, and soil water content are the major factors affecting the distribution and abundance of carabid beetles (Wei et al., 2020; Kazi et al., 2016). The values of diversity indices show high species diversity and evenness in Sambrial followed by Pasrur and Sialkot. Species richness was highest in Pasrur followed by Sambrial and Sialkot. The seasonal diversity of carabid beetles has also been reported similar to our findings from Sindh, Pakistan (Sahito et al., 2020).

CONCLUSION

Carabid beetle assemblages indicated dominance of predatory species in croplands of Sialkot. Among 15 species recorded, *Galerita bicolor* and *Galerita lecontei veracrucis* were reported as new to Pakistan. Our study described *Pheropsophus* as the most abundant genus with seven species out of total fifteen species belong to this genus. Variations in the number of species from three tehsils indicated local differences in the microclimatic conditions in the sites.

ACKNOWLEDGEMENT

Authors are grateful to the Head of Laboratory of Systematics and Pest Management, Department of Zoology-UOG, for providing all the resources and feasible environment for conducting the research work.

Funding

The study received no external funding.

IRB approval

The study was approved by the Ethical Committee of the University of Gujrat, Gujrat.

Ethical statement

All efforts were made to minimize pain and discomfort to the specimens during research.

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Abbas, M., Bai, M. and Yang, X., 2015. Study on dung beetles (Coleoptera: Carabaeidae: Scarabaeinae) of northern Pakistan with a new record from Pakistan. *Entomotax*, **37**: 257-267.
- Abbas, M., Rafi, M.A., Inayatullah, M., Khan, M.R. and Pavulaan, H., 2002. *Taxonomy and distribution of butterflies (Papilionoidea) of the Skardu region, Pakistan*. The Taxonomic Report of the International Lepidoptera Survey. University of Nebraska - Lincoln.
- Abdullah, N.H. and Azmir, I.A., 2021. A pictorial key for the identification of beetle (Order: Coleoptera) and diversity study in selected area within Pelangai Forest Reserve, Negeri Sembilan, Malaysia. *Biodiv. J. Biol. Divers.*, **22**: 947-955. <https://doi.org/10.13057/biodiv/d220250>
- Ahmed, K.S., Majeed, M.Z., Rafi, M.A., Sellami, F. and Afzal, M., 2017. Biodiversity and species distribution of coccinellids (Coccinellidae: Coleoptera) in District Sargodha (Punjab), Pakistan. *Pakistan J. Zool.*, **49**: 1749-1759. <https://doi.org/10.17582/journal.pjz/2017.49.5.1749.1759>
- Ali, S. and Panhwar, W.A., 2017. A checklist of Acrididae (Orthoptera) of Hazara Division Khyber Pakhtunkhwa Pakistan. *J. Ent. Zool. Stud.*, **5**: 96-100.
- Ali, S.M.S., Naeem, M., Baig, F., Shazad, A. and Zia, A. 2015. New records, distributional notes and species diversity of dung beetles (Coleoptera: Scarabaeidae: Scarabaeinae) from Pothohar Plateau of Punjab. *Pakistan. J. Ent. Zool. Stud.*, **3**: 1-6.
- Ane, N.U. and Hussain, M., 2016. Diversity of insect pests in major rice growing areas of the world. *J. Ent. Zool. Stud.*, **4**: 36-41.
- Arshad, M., Nisar, M.F., Majeed, A., Ismail, S. and Ahmad, M., 2011. Ethnomedicinal flora in district

- sialkot, Punjab, Pakistan. *Middle East J. Sci. Res.*, **9**: 209-214.
- Asghar, A., Qadeer, O., Mushtaq, S., Maalik, S., Majeed, W., Bano, N. and Nargis, S., 2022. Assessment of insects diversity with the influence of industrial pollutants in agricultural zones of District Sialkot, Pakistan. *Biodiv. J. Biol. Divers.*, **23**: 2047-2053. <https://doi.org/10.13057/biodiv/d230440>
- Avgin, S.S. and Luff, M.L., 2010. Ground beetles (Coleoptera: Carabidae) as bioindicators of human impact. *Munis Ent. Zool.*, **5**: 209-215.
- Azadbakhsh, S. and Rafi, M.A., 2017. Checklist of ground beetles (Coleoptera, Carabidae) deposited in National Insect Museum of Pakistan. *Orient. Insects*, **51**: 305-312. <https://doi.org/10.1080/00305316.2017.1283256>
- Bashir, M.A., Saeed, S., Sajjad, A., Khan, K.A., Ghramh, H.A., Shehzad, M.A., Mubarak, H., Mirza, N., Mahpara, S. and Rehmani, M.I.A., 2019. Insect pollinator diversity in four forested ecosystems of southern Punjab, Pakistan. *Saudi J. biol. Sci.*, **26**: 1835-1842. <https://doi.org/10.1016/j.sjbs.2018.02.007>
- Bashir, S., Malik, M.F. and Hussain, M., 2023. Spatiotemporal occurrence of beehives of genus *Apis* in Northern Punjab and Azad Jammu and Kashmir, Pakistan. *Kuwait J. Sci.*, **50**: 40-46. <https://doi.org/10.1016/j.kjs.2023.02.007>
- Bibi, M., Bibi, S., Akhtar, N., Ullah, Z., Khan, M.F. and Qureshi, I.Z., 2022. Butterfly (Order: Lepidoptera) species Richness, diversity and distribution in different localities of Battagram, Pakistan. *Saudi J. biol. Sci.*, **29**: 1853-1857. <https://doi.org/10.1016/j.sjbs.2021.10.039>
- Bibi, S. and Rehman, A., 2020. Systematic and diversity of ground beetles (Carabidae: Coleoptera) from district Haripur KPK, Pakistan. *Acta Ent. Zool.*, **1**: 5-7. <https://doi.org/10.33545/27080013.2020.v1.i1a.2>
- Cardoso, P., Barton, P.S., Birkhofer, K., Chichorro, F., Deacon, C., Fartmann, T., Fukushima, C.S., Gaigher, R., Habel, J.C. and Hallmann, C.A., 2020. Scientists warning to humanity on insect extinctions. *Biol. Conserv.*, **242**: 108426. <https://doi.org/10.1016/j.biocon.2020.108426>
- Choate, 1999. *Introduction to the identification of beetles (Coleoptera). Dichotomous keys to some families of Florida Coleoptera*, pp. 23-33.
- Choate, 2001. *Manual for the identification of Ground beetles (Coleoptera: Carabidae)(including tiger beetles) of Florida*. Department Entomology and Nematology, University of Florida, pp. 1-40.
- Clarke, K., 1993. Análises multivariadas não paramétricas de mudanças na estrutura da comunidade. *Rev. Brasil. Ecol.*, **18**: 117-143.
- Clarke, K. and Gorley, R., 2015. *Getting started with primer v7*. PRIMER-E: Plymouth, Plymouth Marine Laboratory, pp. 20.
- Clarke, K. and Warwick, R., 1994. An approach to statistical analysis and interpretation. *Change Mar. Commun.*, **2**: 117-143.
- District Sialkot, 2023. *Climate* [Online]. Lahore: Govt of Punjab Available: <https://sialkot.punjab.gov.pk/climate#:~:text=Sialkot%20is%20hot%20and%20humid,is%20generally%20plain%20and%20fertile>. [Accessed 04 August 2023 2023].
- Ernsting, G. and Isaaks, A., 2000. Ectotherms, temperature, and trade-offs: size and number of eggs in a carabid beetle. *Am. Natur.*, **155**: 804-813. <https://doi.org/10.1086/303361>
- Forister, M.L., Cousens, B., Harrison, J.G., Anderson, K., Thorne, J.H., Waetjen, D., Nice, C.C., De Parsia, M., Hladik, M.L. and Meese, R., 2016. Increasing neonicotinoid use and the declining butterfly fauna of lowland California. *Biol. Lett.*, **12**: 20160475. <https://doi.org/10.1098/rsbl.2016.0475>
- Ghannem, S., Bejaoui, M., Gahdab, C. and Boumaiza, M., 2016. Taxonomic notes on the ground beetles (Coleoptera: Carabidae) of Tunisia. *Arq. Entomol.*, pp. 65-82.
- Ghazanfar, M., Hussain, M., Abbas, Z. and Batool, M., 2017. Diversity, composition and distribution of dung beetle fauna in croplands and pastures of Jhelum, Punjab, Pakistan. *Pak. J. Sci.*, **69**: 369-374.
- Heino, J., Alahuhta, J., Fattorini, S. and Schmera, D., 2019. Predicting beta diversity of terrestrial and aquatic beetles using ecogeographical variables: Insights from the replacement and richness difference components. *J. Biogeogr.*, **46**: 304-315. <https://doi.org/10.1111/jbi.13485>
- Hussain, A., Khan, M.R., Tamkeen, A., Anwar, T., Tahir, S., Ahmad, I. and Qadri, M., 2010. Distributional diversity of hymenopterans pollinator bees from district Skardu, Northern areas of Pakistan. *Pak. J. Ent.*, **25**: 81-86.
- Hussain, A., Rahim, K.M., Ghffar, A., Alia, H. and Jamil, A., 2012. The hymenopterous pollinators of Himalayan foot hills of Pakistan (distributional diversity). *Afr. J. Biotechnol.*, **11**: 7263-7269.
- Hussain, M., Akbar, R., Malik, M.F., Kazam, S.N. and Zainab, T., 2017. Diversity, distribution and seasonal variations of grasshopper populations in Sialkot, Punjab, Pakistan. *Pure appl. Biol.*, **6**: 1372-1381. <https://doi.org/10.19045/bspab.2017.600148>

- Hussain, M., Kanwal, M., Aftab, K., Khalid, M., Liaqat, S., Iqbal, T., Rahman, G. and Umar, M., 2022a. Distribution patterns of dung beetle (Coleoptera: Scarabaeidae) assemblages in croplands and pastures across two climatic zones of Pakistan. *Orient. Insects*, **56**: 392-407. <https://doi.org/10.1080/00305316.2021.2010617>
- Hussain, M., Liaqat, H., Malik, M.F., Aftab, K., Batool, M., Iqbal, R. and Liaqat, S., 2023. Distribution patterns of insect pollinator assemblages at Deva Vatala National Park, Bhimber, Azad Jammu and Kashmir. *Pakistan J. Zool.*, (**Vol?**): 1-7.
- Hussain, M., Malik, M.F., Siddique, S., Umar, M., Zainab, T. and Zafar, F., 2018. Diversity and distribution of coccinellid beetles in irrigated and rainfed fields of Gujrat, Punjab, Pakistan. *Punjab Univ. J. Zool.*, **33**: 1-6. <https://doi.org/10.17582/pujz/2018.33.1.1.6>
- Hussain, M., Nazir, T. and Malik, M.F., 2022b. Community structure of ladybird beetles (Coccinellidae: Coleoptera) alongside the Chenab river at Wazirabad, Gujranwala, Punjab, Pakistan. *Pakistan J. Zool.*, **55**: 1-7. <https://doi.org/10.17582/journal.pjz/20221004171007>
- Hussain, M., Razaq, H., Malik, M.F., Aftab, K., Riaz, J. and Liaqat, S., 2022c. Taxonomic diversity of carabid beetles in the agricultural landscape under hot semi-arid climate in Mandi Bahauddin, Punjab, Pakistan. *Pakistan J. Zool.*, **55**: 1-8. <https://doi.org/10.17582/journal.pjz/20220710070739>
- Hussain, M. and Umar, M., 2021. Faunistic analysis of insects of Deva Vatala National Park and agroecosystem of Gujrat Pakistan: Faunistic analysis of insects orders. *Kuwait J. Sci.*, **50**: 1-25.
- Hussain, M., Younas, M., Malik, M. F., Umar, M., Kanwal, M. and Batool, M., 2020. Spatio-temporal diversity of dung beetles in selected locales of Sialkot, Punjab, Pakistan. *Punjab. Univ. J. Zool.*, **35**: 35-42. <https://doi.org/10.17582/journal.pujz/2020.35.1.35.42>
- Kazi, A., Khatri, I., Rustamani, M. and Wagan, M., 2016. Records of Carabidae in fauna of British India segregated for present boundaries of Pakistan. *Sindh Univ. Res. J. (Sci. Ser.)*, **48**: 833-838.
- Khan, H., Ullah, I., Ali, I., Muhammad, I., Ahmad, T. and Xing, L., 2021. Environmental and soil gradient effects on biodiversity of butterflies in Khyber Pakhtunkhwa, Pakistan. *Arthropods*, **10**: 1-17.
- Khan, M., Rafi, M., Nazir, N., Khan, M., Khan, I., Hayat, A., Ghaffar, A., Rahim, J. and Perveen, F., 2014. Biodiversity of butterflies from poonch division of Azad Kashmir, Pakistan. *J. Agric. Technol.*, **10**: 885-898.
- Khan, M., Ullah, H., Suleman, S., Khan, M., Muhammad, N., Zada, S. and Hussain, S., 2015. A review on diversity of butterfly fauna in Pakistan. *World J. Zool.*, **10**: 313-317.
- Khan, S., 2019. Climate classification of Pakistan. *Int. J. econ. Environ. Geol.*, **10**: 60-71. <https://doi.org/10.46660/ijeeg.Vol10.Iss2.2019.264>
- Khatri, I., Nizamuddin, B.M.A., Rustamani, W.A.P., Ghulam, R.S. and Lakho, M., 2016. Ground beetle fauna (Carabidae) of Mekran Division, Balochistan. *Pakistan J. Ent.*, **31**: 219-225.
- King, J. and Porter, S., 2004. Recommendations on the use of alcohols for preservation of ant specimens (Hymenoptera, Formicidae). *Insectes Soc.*, **51**: 197-202. <https://doi.org/10.1007/s00040-003-0709-x>
- Knapp, M., Seidl, M., Knappová, J., Macek, M. and Saska, P., 2019. Temporal changes in the spatial distribution of carabid beetles around arable field-woodlot boundaries. *Sci. Rep.*, **9**: 8967. <https://doi.org/10.1038/s41598-019-45378-7>
- Lovei, G.L. and Sunderland, K.D., 1996. Ecology and behavior of ground beetles (Coleoptera: Carabidae). *Annu. Rev. Ent.*, **41**: 231-256. <https://doi.org/10.1146/annurev.en.41.010196.001311>
- Luff, M., 1980. The biology of the ground beetle Harpalus rufipes in a strawberry field in Northumberland. *Ann. appl. Biol.*, **94**: 153-164. <https://doi.org/10.1111/j.1744-7348.1980.tb03907.x>
- Malik, R.N., Jadoon, W.A. and Husain, S.Z., 2010. Metal contamination of surface soils of industrial city Sialkot, Pakistan: A multivariate and GIS approach. *Environ. Geochem. Hlth.*, **32**: 179-191. <https://doi.org/10.1007/s10653-009-9274-1>
- Mangi, A., Panhwar, W.A., Shaikh, A.M., Sirohi, M.H., Soomro, S.R.S., Mirbhar, A.A. and Mustafa, S.B., 2023. Study on butterflies (Lepidoptera) from Taluka Larkana, Sindh Pakistan. *J. appl. Res. Pl. Sci.*, **4**: 385-389. <https://doi.org/10.38211/joarps.2023.04.01.46>
- Manzoor, M., Khan, S.W. and Shah, S.A., 2020. An annotated checklist of butterflies at elevated protected areas of Pakistan. *J. Bioresour. Manage.*, **7**: 5. <https://doi.org/10.35691/JBM.0202.0119>
- Maryam, H., Rafi, M.A., Zia, A., Rasul, G., Sheikh, M.K., Qasim, M. and Parveen, G., 2020. Insect pollinator fauna of apricot from Gilgit-Baltistan, Pakistan. *Pak. J. agric. Res.*, **33**: 202-211. <https://doi.org/10.17582/journal.pjar/2020/33.2.202.211>
- Mishkatullah, 2018. Ground beetles fauna of moist temperate forests of Mansehra Division, Khyber

- Pakhtunkhwa*. Pakistan Museum of Natural History, Garden Avenue, Shakarparian, Islamabad-44000.
- Nasir, A., Hussain, M., Fatima, S., Noureen, N. and Ghazanfar, M., 2016. New faunal records of dung beetles from district Sialkot, Punjab, Pakistan. *Biol. environ. Sci.*, **9**: 122-128.
- Noreen, F., Choudri, M., Noureen, S., Muhammad, A., Yaqoob, M., Mustafa, A.G., Cheema, F., Sajjad, F. and Mushtaq, U., 2020. Ethnomedicinal profile of Flora of District Sialkot, Punjab, Pakistan. *New Trends med. Sci.*, **1**: 65-83.
- Noureen, N., Hussain, M. and Malik, M.F., 2015a. Taxonomic account of dung beetles from Gujrat, Punjab (Pakistan). *Biol. environ. Sci.*, **7**: 20-26.
- Noureen, N., Hussain, M., Malik, M.F. and Afsheen, S., 2015b. New records of dung beetle fauna from Pakistan. *J. Ent. Zool. Stud.*, **3**: 428-430.
- Noureen, N., Hussain, M., Malik, M. F., Umar, M., Abbas, Z. and Munaward, S., 2021. Habitat types effect on diversity, distribution and abundance of dung beetles: Dung beetles diversity in different habitats. *Biol. Sci. Pak. J. scient. indust. Res.*, **64**: 217-224. <https://doi.org/10.52763/PJSIR.BIOL.SCI.64.3.2021.217.224>
- Perveen, F.K., 2015. Checklist of butterfly (Insecta: Lepidoptera) fauna of Tehsil Tangi, Khyber Pakhtunkhwa, Pakistan. *Arthropods*, **4**: 98-106.
- Pizzolotto, R., Mazzei, A., Bonacci, T., Scalercio, S., Iannotta, N. and Brandmayr, P., 2018. Ground beetles in Mediterranean olive agroecosystems: Their significance and functional role as bioindicators (Coleoptera, Carabidae). *PLoS One*, **13**: e0194551. <https://doi.org/10.1371/journal.pone.0194551>
- Post, R., Flook, P. and Millest, A., 1993. Methods for the preservation of insects for DNA studies. *Biochem. Syst. Ecol.*, **21**: 85-92. [https://doi.org/10.1016/0305-1978\(93\)90012-G](https://doi.org/10.1016/0305-1978(93)90012-G)
- Rafi, M.A., Carpenter, J.M., Qasim, M., Shehzad, A., Zia, A., Khan, M.R., Mastoi, M.I., Naz, F., Ilyas, M. and Shah, M., 2017. The vespidae fauna of Pakistan. *Zootaxa*, **4362**: 1-28. <https://doi.org/10.11646/zootaxa.4362.1.1>
- Rahim, J., Khan, M.R., Khan, M.A.G. and Nazir, N., 2013a. Biodiversity of ground beetles (Coleoptera: Carabidae) from District Poonch, Azad Kashmir. *Pakistan J. Zool.*, **45**: 1447-1451.
- Rahim, J., Khan, M.R. and Nazir, N., 2013b. Systematic and abundance of ground beetles (Carabidae: Coleoptera) from district Poonch Azad Kashmir, Pakistan. *J. Agric. Vet. Sci.*, **6**: 24-29. <https://doi.org/10.9790/2380-0622429>
- Rainio, J. and Niemelä, J., 2003. Ground beetles (Coleoptera: Carabidae) as bioindicators. *Biodivers. Conserv.*, **12**: 487-506. <https://doi.org/10.1023/A:1022412617568>
- Ramzan, U., Majeed, W., Rana, N. and Nargis, S., 2021. Occurrence of different insect species with emphasis on their abundance and diversity in different habitats of Faisalabad, Pakistan. *Int. J. trop. Insect Sci.*, **41**: 1237-1244. <https://doi.org/10.1007/s42690-020-00314-5>
- Raven, P. and Wagner, D., 2021. Agricultural intensification and climate change are rapidly decreasing insect biodiversity. *Proc. natl. Acad. Sci.*, **118**: e2002548117. <https://doi.org/10.1073/pnas.2002548117>
- Rebrina, F., Petek, M., Gulin, V. and Brigić, A., 2022. Ground beetle assemblages respond to motorway proximity through changes in functional rather than taxonomic diversity in a grassland ecosystem. *Glob. Ecol. Conserv.*, **38**: e02255. <https://doi.org/10.1016/j.gecco.2022.e02255>
- Ribera, I., Foster, G.N., Downie, I.S., McCracken, D.I. and Abernethy, V.J., 1999. A comparative study of the morphology and life traits of Scottish ground beetles (Coleoptera, Carabidae). *Annls Zool. Fenn*, 21-37.
- Roberts, T.J., 2001. *The butterflies of Pakistan*. Oxford University Press.
- Rounsevell, M., Fischer, M., Rando, A.T.M., Mader, A., Caplat, P., Ekroos, J. and Smith, H.G., 2018. *The IPBES regional assessment report on biodiversity and ecosystem services for Europe and Central Asia*. Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem. <https://doi.org/10.17011/conference/eccb2018/109102>
- Sahito, R., Memon, N. and Amur, A., 2020. 32 seasonal variation and diversity of carabid beetles fauna (Coleoptera: Carabidae) in coastal areas of Sindh, Pakistan. *Pure appl. Biol.*, **9**: 1556-1567. <https://doi.org/10.19045/bspab.2020.90163>
- Satpathi, C.R., 2021. *Ground beetles and their role in management of crop pests*, CRC Press. <https://doi.org/10.1201/9781003261087>
- Schaffers, A.P., Raemakers, I.P., Sýkora, K.V. and Ter Braak, C.J., 2008. Arthropod assemblages are best predicted by plant species composition. *Ecology*, **89**: 782-794. <https://doi.org/10.1890/07-0361.1>
- Schauff, M.E., 2001. *Collecting and preserving insects and mites: Techniques and tools*. National Museum of Natural History, NHB-168, Washington, DC 20560, Systematic Entomology Laboratory, USDA.
- Schwerk, A., Wińska-Krysiak, M., Przybysz, A., Zarsa-

- Januszkiewicz, E. and Sikorski, P., 2020. Carabid beetle (Coleoptera: Carabidae) response to soil properties of urban wastelands in Warsaw, Poland. *Sustainability*, **12**: 10673. <https://doi.org/10.3390/su122410673>
- Shannon, C.E., 1949. *The mathematical theory of communication, by CE Shannon (and recent contributions to the mathematical theory of communication), W. Weaver*, University of Illinois Press Champaign, IL, USA.
- Siddiqui, H., Zubair, A. and Khatri, I., 2014. Distributional notes and new records for the dung beetles (Coleoptera: Scarabaeidae: Scarabaeinae) of Pakistan. *Pakistan J. Zool.*, **46**: 295-307.
- Simpson, E.H., 1949. Measurement of diversity. *Nature*, **163**: 688-688. <https://doi.org/10.1038/163688a0>
- Skłodowski, J., 2017. Manual soil preparation and piles of branches can support ground beetles (Coleoptera, carabidae) better than four different mechanical soil treatments in a clear-cut area of a closed-canopy pine forest in northern Poland. *Scand. J. For. Res.*, **32**: 123-133. <https://doi.org/10.1080/02827581.2016.1195868>
- Soomro, F.A., Panhwar, W.A., Shaikh, A.M., Memon, K.H., Memon, Z.U.N., Ujjan, S.A. and Ujan, J.A., 2021. Ground beetles (Coleoptera: Carabidae) collected from different ecological zones of Sukkur Sindh-Pakistan with new records. *Int. J. Emerg. Technol.*, **12**: 103-107.
- Symondson, W., Sunderland, K. and Greenstone, M., 2002. Can generalist predators be effective biocontrol agents? *Annls Rev. Ent.*, **47**: 561-594. <https://doi.org/10.1146/annurev.ento.47.091201.145240>
- Szyszkó-Podgórska, K., Dymitryszyn, I., Jankiewicz, U., Kondras, M., Żyfką-Zagrodzińska, E. and Schwerk, A., 2021. Assemblage characteristics of butterflies and carabid beetles as a function of soil characteristics and plant diversity in differently managed fields, forests and ecotones: A case study in Tuczno Forest District, Poland. *Land*, **11**: 25. <https://doi.org/10.3390/land11010025>
- Ullah, M., Naeem, M., Mahmood, K. and Rafi, M.A., 2017. Faunistic studies of the tribe Brachinini (Carabidae: Coleoptera) from northern Pakistan. *Zootaxa*, **4232**: zootaxa. 4232. <https://doi.org/10.11646/zootaxa.4232.2.2>
- van der Meer, M., Kay, S., Lüscher, G. and Jeanneret, P., 2020. What evidence exists on the impact of agricultural practices in fruit orchards on biodiversity? A systematic map. *Environ. Evid.*, **9**: 1-27. <https://doi.org/10.1186/s13750-020-0185-z>
- Vichitbandha, P. and Wise, D.H., 2002. A field experiment on the effectiveness of spiders and carabid beetles as biocontrol agents in soybean. *Agric. For. Ent.*, **4**: 31-38. <https://doi.org/10.1046/j.1461-9563.2002.00118.x>
- Wang, M., Axmacher, J.C., Yu, Z., Zhang, X., Duan, M., Wu, P., Zou, Y. and Liu, Y., 2021. Perennial crops can complement semi-natural habitats in enhancing ground beetle (Coleoptera: Carabidae) diversity in agricultural landscapes. *Ecol. Indic.*, **126**: 107701. <https://doi.org/10.1016/j.ecolind.2021.107701>
- Wei, S., Huang, W., Zhu, M., Gao, L., Wang, Y., Zhang, R., Li, Z. and Zhao, Z., 2020. The asymmetric responses of carabid beetles to steppe fragmentation in Northwest China. *Glob. Ecol. Conserv.*, **23**: e01058. <https://doi.org/10.1016/j.gecco.2020.e01058>
- Wepprich, T., Adrion, J.R., Ries, L., Wiedmann, J. and Haddad, N.M., 2019. Butterfly abundance declines over 20 years of systematic monitoring in Ohio, USA. *PLoS One*, **14**: e0216270. <https://doi.org/10.1371/journal.pone.0216270>
- Worthen, W.B. and Merriman, D.C., 2013. Relationships between carabid beetle communities and forest stand parameters: Taxon congruence or habitat association? *SouthEast Nat.*, **12**: 379-386. <https://doi.org/10.1656/058.012.0211>
- Yu-Feng, M., Khan, Z., Perveen, F., Rafi, M.A., Shah, S.W., Xiao-Hong, S. and Xing, L., 2020. Biodiversity of butterflies in Tangi Charsadda, Khyber Pakhtunkhwa, Pakistan. *Pakistan J. Zool.*, **52**: 835. <https://doi.org/10.17582/journal.pjz/20190326090338>