



## Review Article

# Cytoplasmic and Nuclear Maturation of Intact and Reconstructed Oocytes Controlling the Developmental Competence of Embryos

Abd El-Nasser Ahmed Mohammed<sup>1\*</sup>, Shaker Al Suwaiegh<sup>1</sup>, Ibrahim Al-Gherair<sup>1</sup>, Sami Al-khamis<sup>1</sup>, Waleed Alhujaili<sup>2</sup>, Alhassan Mohammed<sup>3</sup> and Alhossein Mohammed<sup>3</sup>

<sup>1</sup>Department of Animal and Fish Production, King Faisal University, P.O. Box 402, Al-Ahsa 31982, Saudi Arabia

<sup>2</sup>College of Human Medicine, Assiut University 7111, Egypt

<sup>3</sup>Department of Biology, College of Science, Taibah University, Al-Madinah Al-Munawwarah, 41321, Saudi Arabia.

## ABSTRACT

Oocytes grow and develop within several distinct ovarian follicles during the fetal and postnatal periods, and they ovulate in each estrous or menstrual cycle from ovarian follicles after puberty. Ovarian follicles start to form when oogonia are surrounded by a single layer of pregranulosa follicular cells and are called primordial follicles. Primordial follicles grow and develop into further stages upon activation. The growth and development of primordial follicles to pre-ovulatory follicles in mammals lasts several months. Oocytes acquire developmental competence during the preantral and antral ovarian follicle stages. The final maturation stage of ovarian follicles occurs during estrous or menstrual cycles. Changes occur in the cytoplasm, nuclei, and the surrounding follicular cells of oocytes for completion of meiotic maturation, fertilization, and further embryo development either *in vivo* or *in vitro*. The cytoplasmic maturation of oocytes helps to surpass the maternal zygotic block and progress for further embryo development. Hence, the cytoplasm of enucleated germinal vesicle (GV) or mature (metaphase II; MII) oocytes control development after embryonic/somatic nuclear transfer. Therefore, this review was conducted to collect and consolidate the current knowledge of cytoplasm and nuclear material effects on maturation of intact and reconstructed oocytes and further developmental competence of embryos *in vivo* and *in vitro* and their importance for nucleus reprogramming and fertility.

## Article Information

Received 14 February 2024

Revised 06 May 2024

Accepted 11 May 2024

Available online 16 October 2024 (early access)

Published 29 August 2025

## Authors' Contribution

AAM and SA designed the study and wrote the manuscript. IA collected the data. SA and WA analyzed the results and wrote the manuscript. AM and AM prepared figures. All authors revised the manuscript, and approved the final version.

## Key words

Follicle, Cumulus, Cytoplasm, Nucleus, Nucleolus, Maturation, Embryo

## INTRODUCTION

Oocytes grow and develop during fetal and postnatal periods within several distinct ovarian follicles, and they ovulate from the ovarian follicles after puberty in each estrous or menstrual cycle. The formation of ovarian follicle begins with primordial follicles, where oogonia are surrounded by pregranulosa cells

(O'Connell and Pepling, 2021; Mohammed *et al.*, 2022). The growth and development of primordial follicles occur upon activation of pregranulosa cells to reach the preantral and antral follicle stages (Zhang *et al.*, 2014; Wu *et al.*, 2021) (Fig. 1).

The changes that occur during follicle development include granulosa cell proliferation, oocyte growth, and theca cell differentiation (Richards and Pangas, 2010). Preantral follicle growth is gonadotropin-independent, whereas antral follicle growth is gonadotropin-dependent (Chakravarthi *et al.*, 2021). Tang *et al.* (2012) investigated the role of follicle stimulating hormone (FSH) in combination with growth and differentiation factor-9 or basic fibroblast growth factor on the survival, activation, and growth of bovine primordial follicles, which improved after long-term culture of ovarian tissue.

Most of the information on ovarian follicle growth and development stems from studies on ruminants, swine,

\* Corresponding author: aamohammed@kfu.edu.sa  
0030-9923/2025/0005-2465 \$ 9.00/00



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Fig. 1. Ovarian follicle structures upon fallopian tube ligation in mice; preantral follicles (A), antral follicles (B) and corpora lutea (C) structures. N.B. Differences in sizes of oocytes and the layers of surrounding cells inside the follicles.




Fig. 2. The structures of ovarian follicle in control (A) and ovarian transplanted tissues in rats (B, C). The images show preantral follicles (A), antral follicles (B), and corpora lutea (C).

rodent species, and humans (Senosy *et al.*, 2017; Bunel *et al.*, 2020; Baerwald and Pierson, 2020; Pournaghi *et al.*, 2021; Ko *et al.*, 2021). Several factors such as species (Gordon, 2003; Patton *et al.*, 2021), nutrition and feed additives (Mohammed and Attaai, 2011; Fernandes *et al.*, 2016; Senosy *et al.*, 2017, 2018; Mohammed 2018, 2019; Liang *et al.*, 2012; Moulavi and Hosseini, 2019; Gutiérrez-Añez *et al.*, 2021; Saini *et al.*, 2022; Al-Mufarji *et al.*, 2022; Al-Masruri *et al.*, 2022a, b) and males (Taira *et al.*, 2022) have effect on ovarian follicle growth and development. In addition, recently adapted techniques of ovarian synchronization, ovarian superovulation, and

ovarian transplantation for dysfunction treatment and the maximizing of reproduction are known to change ovarian follicle growth and the development of the oocytes they contain (Gordon, 2003; Ata and Telek, 2021) (Fig. 2). Superovulation protocol via gonadotrophin stimulation might result in abnormal changes in cytoplasm and nucleus structure, lower the concentration of estradiol hormone, and disrupt cumulus cell mRNA (Algriany *et al.*, 2007; von Wolff *et al.*, 2022). Moreover, Kanitz *et al.* (2002) found that high doses of FSH could lead to defects in ovulation at the level of the pituitary gland and ovaries. Consequently, the transfer of GV nuclei, nucleoli, or cytoplasts of highly competent oocytes to abnormal oocytes through adapted techniques may treat abnormalities in oocytes.

Oocytes acquire developmental competence during ovarian follicle development through changes in the nucleus and cytoplasm. Therefore, the recipient cytoplasm and donor nucleus stages of artificial gametes affect developmental competence. Such artificial gametes may be used for different purposes, such as studying of embryonic/somatic nucleus reprogramming in recipient cytoplasts, production of cloned and transgenic animals, and treatment of infertility (Campbell *et al.*, 1996; Wilmut *et al.*, 1997; Mohammed *et al.*, 2019; Singh *et al.*, 2022; Dolma *et al.*, 2022). The roles of cytoplasmic maturation and nuclear progression to the MII stage of intact and reconstructed oocytes in controlling developmental competence of oocytes to embryos are compiled and discussed in this review.

### ASPIRATION AND DEVELOPMENTAL COMPETENCE OF CUMULUS-ENCLOSED GERMINAL VESICLE OOCYTE

Cumulus-enclosed germinal vesicle oocytes (COCs) can be aspirated or picked-up (OPU) from different animals, humans, or slaughterhouse ovaries (Mohammed *et al.*, 2005; Jin *et al.*, 2016; Ongaratto *et al.*, 2020; Girsh, 2021; Al-Zeidi *et al.*, 2022a, b) for *in vitro* embryo production. The OPU approach has been applied in humans to overcome abnormalities such as hidden ovaries, severe tubal diseases, or multiple adhesions with a success rate lower than 50%. Oocytes are picked up from genetically superior animals before puberty, early gestation, and early postpartum through laparoscopic and ultrasonic methods to shorten the interval between generations and increase animal production (Ongaratto *et al.*, 2020; Furukawa *et al.*, 2021). COCs have been used for *in vitro* embryo production through oocyte maturation (IVM), fertilization or activation (IVF), and culture (IVC) of different animal species and humans. The development of COCs to blastocyst stage does not exceed 30-40% (Mohammed

*et al.*, 2005; Somfai *et al.*, 2012). Several factors, such as ovary storage time, wave of estrous cycle, follicle diameter, oocyte quality, incubation time, maturation environment, nutrition and feed additives, and male effect, affect the developmental competence of aspirated oocytes to embryos (Schwartz *et al.*, 1998; Demyda-Peyrás *et al.*, 2013; Senosy *et al.*, 2017, 2018; Taira *et al.*, 2022) (Table I).

The development of oocytes to embryos increases when slaughterhouse ovaries are exposed to warm conditions for a few h before oocyte aspiration. Storage of ovaries at 30 °C for 3-4 h after slaughter has led to an

increase in the number of oocytes developed to embryos. It was suggested that a storage period of 4 h resulted in the formation of an environment within the 3-8 mm follicles, similar to that which occurs in preovulatory follicles. In our study, changes in follicular fluid (FF) composition at times of zero, 4, and 8 h in goat ovaries were investigated. The most notable changes in FF composition were the concentrations of glucose, cholesterol, and triglycerides, which are sources of energy for oocytes. This might partially explain the increase in the developmental competence of oocytes collected from the ovaries 4 h after slaughter but decrease thereafter.

**Table I. Factors affecting developmental competence of aspirated oocytes *in vitro*.**

| Ovaries                | Treatments                            | Effects                                                                                                                                              | References                                                                                                           |
|------------------------|---------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------|
| Slaughterhouse ovaries | Storage the ovaries 30 °C for 3-4 h   | Increase developmental competence of oocytes to embryos                                                                                              | Luu <i>et al.</i> , 2011                                                                                             |
|                        | Defined and undefined media           | Variable effects on oocyte maturation and embryo development                                                                                         | Mohammed <i>et al.</i> , 2005; Madkour <i>et al.</i> , 2016; Spacek and Carnevale, 2018; Abdoon <i>et al.</i> , 2018 |
|                        | IGF-1 and FF                          | Variable effects on oocyte maturation and embryo development                                                                                         | Oberlender <i>et al.</i> , 2013                                                                                      |
|                        | Fetal calf serum                      | Increase developmental competence of oocytes to embryos                                                                                              | Mohammed <i>et al.</i> , 2005; Moulavi and Hosseini, 2019                                                            |
|                        | Follicular fluid                      | Variable effects on oocyte maturation and embryo development depending on follicle sizes collected and added percentage to the maturation medium     | Mohammed <i>et al.</i> , 2005; Sinclair <i>et al.</i> , 2008                                                         |
|                        | Co-culture cumulus and oviduct cells  | Increase developmental competence of oocytes to embryos                                                                                              | Lee <i>et al.</i> , 2018                                                                                             |
|                        | Cumulus cells surrounding the oocytes | Increase developmental competence of oocytes to embryos                                                                                              | Mohammed 2006; Mohammed <i>et al.</i> , 2005, 2008, 2010, 2019; Al-Zeidi <i>et al.</i> , 2022                        |
|                        | Hormone                               | Increase developmental competence of oocytes to embryos                                                                                              | Moulavi and Hosseini, 2019                                                                                           |
|                        | Follicle size                         | Variable effects where the increase of follicle sizes enhance developmental competence of oocytes to embryos                                         | Shabankareh <i>et al.</i> , 2014; Gordon, 2003; Maylinda <i>et al.</i> , 2018                                        |
| Live organisms         | Follicle size                         | Variable effects on oocyte maturation and embryo development where the higher follicle size the higher developmental competence of oocyte to embryos | Hasler <i>et al.</i> , 1998; Maylinda <i>et al.</i> , 2018; Raj <i>et al.</i> , 2018                                 |
|                        | Follicular wave                       | Variable effects on oocyte maturation and embryo development                                                                                         | Gordon <i>et al.</i> , 2003; Cavalieri <i>et al.</i> , 2018; Baby and Bartlewski, 2011                               |
|                        | Follicular and luteal stages          | Variable effects on oocyte maturation and embryo development                                                                                         | Gordon <i>et al.</i> , 2003                                                                                          |
|                        | Nutrition                             | Variable effects on oocyte maturation and embryo development                                                                                         | Gordon <i>et al.</i> , 2003                                                                                          |
|                        | Feed additives                        | Variable effects on oocyte maturation and embryo development                                                                                         | Senosy <i>et al.</i> , 2017, 2018; Mohammed and Al-Hozab, 2018                                                       |

The more cumulus cells surround the oocytes and homogeneity of the oocytes cytoplasts, the greater the developmental competence of oocytes to embryos (Mohammed *et al.*, 2005). In addition, the male effect stimulation of super ovulated females during the last phase of the FSH regimen increase the number and diameter of large follicles, in addition to the number of corpora lutea and viable embryos, compared to unstimulated ewes (Taira *et al.*, 2022). Therefore, these factors should be taken into consideration for enhancement of oocyte quality, oocyte maturation, fertilization, and development of embryos.

### OOCYTE QUALITY AND GENE EXPRESSION

Oocyte quality is critical for maturation, fertilization, and embryonic and fetal development. The higher the oocyte quality, the higher the developmental competence of the resulting embryo and fetus. Oocytes are graded morphologically according to the cumulus cells surrounding the oocytes, diameter of the oocyte, and homogeneity of the cytoplasts (Mohammed *et al.*, 2005; Lopes *et al.*, 2010; Al-Zeidi *et al.*, 2022). Castaneda *et al.* (2013) found that both active mitochondria and lipid content were correlated with oocyte diameter. In addition, brilliant cresyl blue staining (BCB) were used to select developmentally competent GV oocytes of different species (Opieła and Kańska-Książkiewicz, 2013; Jewgenow *et al.*, 2019; Mohammed and Al-Hozab, 2020). Bovine GV oocytes stained with +BCB had a relatively higher developmental competence than non-stained BCB oocytes. A previous study investigated the relationship between BCB staining, lipid content, and active mitochondria. They concluded that higher lipid content of bovine oocytes stained with BCB might provide higher developmental competence. In addition, Sinclair *et al.* (2008) investigated the amino and fatty acid content in the follicular fluid of bovine follicles as a predictor of embryo development. They found that two amino acids (alanine and glycine) had the highest value for predicting early embryonic development, and this might serve as a useful tool in measuring COCs quality.

In recent years, transcriptome and proteome have been substantially explored as biomarkers for oocyte quality owing to the progression of the new technologies. The approach is an invaluable tool for clinical practice in assisted reproductive technologies, obstetrics, and gynecology (Li *et al.*, 2019). The expression patterns of glucose metabolism genes (G6PDH, LDH, and PDH) were investigated in buffalo oocytes matured under different glucose concentrations (Kumar *et al.*, 2013). The pattern of G6PDH expression during oocyte maturation and early embryonic development is predictive of oocyte quality

and developmental competence of embryos. However, Liu *et al.* (2021) found in good prognosis IVF patients that mitochondrial DNA copy number of cumulus cells is not linked to embryo implantation. Therefore, further studies on gene expression are required to determine the relation between oocyte quality and further development of oocytes into preimplantation and post implantation embryo stages.

### NUCLEAR AND CYTOPLASMIC MATURATION OF INTACT OOCYTES

Oocytes require several months to acquire developmental competence before maturation (Blondin *et al.*, 1997; Hashimoto *et al.*, 2002). Small follicles contain inhibiting maturation substances within the follicles, which disappear in the preovulatory follicle simultaneously with the LH surge. Therefore, oocytes resume the first meiotic division to the metaphase II (MII) stage in vivo after the LH surge and *in vitro* when they are removed from antral follicles and cultured under favorable *in vitro* conditions (38.5 °C, 90% humidity, 5.0% CO<sub>2</sub>).

Oocyte maturation *in vitro* was described nearly 85 years ago when rabbit and human oocytes were aspirated from follicles and cultured in a favorable medium (Pincus and Enzmann, 1935). It has been indicated that the developmental competence of *in-vitro* matured oocytes is lower than that of *in-vivo* matured oocytes (Leibfried-Rutledge *et al.*, 1987; Margalit *et al.*, 2019; Sakaguchi and Nagano, 2020) due to insufficient cytoplasmic maturity (Blondin *et al.*, 1997). The longer the maturation time of oocytes, the lower the developmental competence of the resulting embryos, as indicated in swine and camel oocytes, compared to oocytes of other species (Wani and Hong, 2020; Mohammed and Al-Hozab, 2020; Li *et al.*, 2022).




Fig. 3. The germinal vesicle karyoplasts placed under the zona pellucida of enucleated germinal vesicle oocytes (A) and fused with cytoplasts (B).

The morphological maturation stages of intact GV or reconstructed GV oocytes (Figs. 3 and 4) include oocyte nuclear and cytoplasmic maturation, in addition to cumulus expansion of the surrounding cells (Fig. 4). The germinal

vesicle of denuded oocytes is visible in rodent, rabbit, and human oocytes, whereas it is not visible in ruminant oocytes because of the presence of lipid droplets in the cytoplasm. [Gordon \(2003\)](#) described the chronology of events during bovine oocyte maturation as inductive and synthetic phases. The nuclear changes included germinal vesicle breakdown (GVBD), which occurred at 2-3 h in rodent oocytes and 6-8 h in ruminant oocytes, respectively. Rodent oocytes reached the MI stage at 11-13 h and MII stage at 15-17 h. The ruminant oocytes reached the MI stage at 15-21 h and MII stage at 24 h ([Gordon, 2003](#); [Virant-Klun et al., 2018](#); [Mohammed et al., 2008, 2010, 2019](#)).




Fig. 4. Oocyte maturation; denuded germinal vesicle oocyte (A), germinal vesicle breakdown (B), cumulus expansion (C), polar body extrusion (D), fluorescence staining of matured oocyte (E), hematoxylin stain of matured oocyte (F).

Follicle size, follicular wave during the estrous cycle, follicular and luteal stage of estrous, species and age ([Gordon, 2003](#); [Patton et al., 2021](#)), nutrition and feed additives ([Cavalieri et al., 2018](#); [Mohammed, 2018, 2019](#); [Mohammed and Al-Hozab, 2020](#); [Liang et al., 2012](#); [Moulavi and Hosseini, 2019](#); [Pournaghi et al., 2021](#); [Gutiérrez-Añez et al., 2021](#); [Saini et al., 2022](#)), and male effects ([Taira et al., 2022](#)) were found to affect oocyte maturation. In addition, super stimulation of ovarian follicles via gonadotropin injections resulted in changes in small, medium, and large follicles according to the number and dose of gonadotropin injections and side of the ovary ([Abdelnaby et al., 2021](#)). FSH stimulates transcription and translation in ovarian granulosa cells, which are essential for female reproductive endocrine regulation ([Dai et al., 2021](#)). The timing of fully grown germinal vesicle oocytes to reach the MII stage *in vitro* for different species and the various approaches for improving maturation are presented in [Table II](#) ([Gordon, 2003](#); [Baruffi et al., 2004](#);

[Somfai et al., 2012](#)). Enrichment of maturation media with cumulus cells, hormones, growth factors, and other factors improves oocyte maturation and embryo development.

## CYTOPLASMIC MATURATION AND NUCLEAR REPROGRAMMING OF RECONSTRUCTED OOCYTES AND DEVELOPMENTAL COMPETENCE

Germinal vesicle nucleus, nucleolar, or cytoplasmic dysfunction of oocytes may be associated with infertility in animal and human oocytes ([Fulka et al., 2004](#); [Miao et al., 2019](#); [Wang et al., 2022](#)). Oocyte GV nuclei, nucleoli, MII spindles, and cytoplasm transfer techniques have been used for the treatment of oocyte dysfunction due to cytoplasmic, nuclear, or nucleolar reasons or to increase their developmental competence ([Fulka et al., 2004](#); [Hoseini et al., 2016](#); [Benc et al., 2018](#)). [Hoseini et al. \(2016\)](#) concluded that cytoplasm transfer technique is not effective for cytoplasmic maturity of recipient GV oocytes; however, the presence of cumulus cells during oocyte maturation or GV nucleus or nucleolus transfer is effective in increasing the developmental competence of the resulting embryos. [Chang et al. \(2005\)](#) found that the developmental incompetency of denuded mouse GV oocytes matured *in vitro* is ooplasmic in nature and is associated with aberrant Oct-4 expression.

One of the fundamental factors for successful assisted reproductive technologies (ART) in reconstructed oocytes is the interaction between the recipient cytoplasm and donor nucleus ([Polanski et al., 2005](#); [Mohammed, 2006](#); [Mohammed et al., 2008, 2010](#); [Wani et al., 2018](#)). Oocytes at stages GV, ProMI, MI, and MII were enucleated to obtain recipient cytoplasts. The obtained cytoplasts can be used in cell biology for basic research and reprogramming of the introduced germ, embryonic, or somatic nuclei during transfer ([Grabarek et al., 2004](#); [Polanski et al., 2005](#); [Mohammed et al., 2008, 2010, 2019](#)). Such approaches might solve the problem of low developmental competence of oocytes from prepubertal or advanced maternal age ([Hassold and Chiu, 2003](#)) due to developmental incompetency of ooplasmic or nuclear nature ([Bao et al., 2003](#)). The following questions arise: Does the transfer of nuclei from non-growing, growing, or aging GV oocytes into fully grown GV cytoplasts solve infertility problem? Does the transfer of MII spindles to competent mature MII cytoplasts rescue infertility in animals and humans? Does the transfer of competent nucleoli from donor fully grown GV oocytes to recipient GV oocytes containing abnormal nucleoli rescues infertility?

**Table II. Duration of oocyte maturation *in vitro* from germinal vesicle to metaphase II stage using different supplements to maturation media in different animal species and human.**

| Species | Duration, h | Treatments                                         | References                                |
|---------|-------------|----------------------------------------------------|-------------------------------------------|
| Rodents | 17          | Cumulus cells                                      | Grabarek <i>et al.</i> , 2004             |
|         |             | Cumulus cells                                      | Mohammed 2006, 2008                       |
|         |             | Cumulus cells                                      | Mohammed <i>et al.</i> , 2006, 2008, 2010 |
|         |             | Perfluorooctane                                    | Wei <i>et al.</i> , 2021                  |
|         |             | CRH and ACTH                                       | Gong <i>et al.</i> , 2021                 |
| Goat    | 24          | Cysteamine, leukemia inhibitory factor, and Y27632 | An <i>et al.</i> , 2018                   |
|         |             | Physiological oocyte maturation                    | Suresh <i>et al.</i> , 2021               |
|         |             | Leptin                                             | de Senna Costa <i>et al.</i> , 2022       |
| Sheep   | 24          | Resveratrol                                        | Zabihi <i>et al.</i> , 2020               |
|         |             | Oxygen tension                                     | Sánchez-Ajofrín <i>et al.</i> , 2020      |
|         |             | C-type natriuretic peptide                         | Zhang <i>et al.</i> , 2018                |
| Cattle  | 24          | FF and FCS                                         | Mohammed <i>et al.</i> , 2005             |
|         |             | Retinoic acid                                      | Borges <i>et al.</i> , 2021               |
|         |             | Lycopene                                           | Residiwati <i>et al.</i> , 2021           |
| Buffalo | 24          | -                                                  | Marin <i>et al.</i> , 2019                |
|         |             | Brain-derived neurotrophic factor                  | Zhao <i>et al.</i> , 2019                 |
|         |             | Retinoic acid                                      | Gad <i>et al.</i> , 2018                  |
| Rabbit  | 24          | Gonadotropin-releasing hormone                     | Yoshimura <i>et al.</i> , 1991            |
|         |             | -                                                  | Arias-Álvarez <i>et al.</i> , 2017        |
| Human   | 24          | -                                                  | Chian <i>et al.</i> , 2004                |
|         |             | -                                                  | Baruffi <i>et al.</i> , 2004              |
|         |             | Coenzyme Q10                                       | Ma <i>et al.</i> , 2020                   |
|         |             | A fertilin-derived peptide                         | Sallem <i>et al.</i> , 2022               |
| Camel   | 42-48       | -                                                  | Wani <i>et al.</i> , 2018                 |
|         |             | Macromolecule                                      | Moulavi and Hosseini, 2019                |
|         |             | Roscovitine                                        | Wani and Hong 2018; 2020                  |
| Pig     | 48          | Dihydroartemisinin                                 | Luo <i>et al.</i> , 2018                  |
|         |             | Zearalenone                                        | Wang <i>et al.</i> , 2022                 |
|         |             | CRH and ACTH                                       | Gong <i>et al.</i> , 2021                 |
|         |             | Allicin                                            | Li <i>et al.</i> , 2022                   |

Therefore, the techniques of enucleation of denuded GV, cumulus-enclosed GV, and enucleation of GV oocytes were adapted for such studies (Grabarek *et al.*, 2004; Mohammed *et al.*, 2008, 2010, 2019). Our unique enucleation and enucleation techniques (Figs. 5 and 6) of denuded and cumulus-enclosed GV oocytes enabled us to confirm the role of cumulus cells, nuclear material, and nucleoli on oocyte maturation and embryo development. Complete enucleation of denuded GV oocytes removes the entire GV nucleus. Therefore, the resulting cytoplasm does not contain any GV components or cumulus effects. Selective enucleation of denuded GVs involves the removal of nuclear membrane, leaving the nucleolus and nuclear sap in the cytoplasm; however, selective enucleation of cumulus-enclosed GVs leaves the effect of nucleolus, nuclear sap, and cumulus cells in the cytoplasm (Fig. 5). Therefore, the aforementioned techniques

and enucleation confirmed the role of cumulus cells, nuclear material, and nucleolus in oocyte maturation and developmental competence.

Our results of selective enucleation of GV oocytes yielded GV cytoplasm in similar quantity as did complete enucleation of GV oocytes (Mohammed, 2006; Mohammed *et al.*, 2008, 2010). Therefore, the aforementioned techniques and enucleation confirmed the role of cumulus cells, nuclear material, and nucleolus in oocyte maturation and further developmental competence of embryos. Our results indicated that GVsel cytoplasm provided proper timing of maturation after reconstruction with GV, contrary to GV cytoplasm, which accelerated timing of maturation. However, this did not improve haploidization, as measured with abnormal nuclear morphology. Abnormal acceleration of maturation cannot be rescued by any particular cell cycle stage or developmental stage of blastomeres 1/2, 1/4

or 1/8 used as nuclear donor. Highest maturation efficiency of GV cytoplasts together with their least disturbance from abnormal meiosis were provided by G2 donor blastomere nuclei (Mohammed, 2006; Mohammed *et al.*, 2008, 2010, 2019, 2022).




Fig. 5. Obtaining types of recipient germinal vesicle cytoplasts, donor nuclei transfer and further meiotic maturation and developmental competence; GV germinal vesicle nucleus of fully-grown oocyte, G1 gap 1 stage, S S-phase stage, G2 gap 2 stage.




Fig. 6. Enucleation of fully grown germinal vesicle oocyte (A and B) the nucleolus-free germinal vesicles are visible (red arrows) in the cytoplasm. In the enucleation pipettes, the nucleolus of GV oocyte undergoing enucleation (A, yellow arrow) and several nucleoli of previously enucleated GV oocytes (B yellow arrows) are visible.

The nucleolus and cumulus cells were confirmed to be dispensable for GV oocyte maturation, but the resulting embryos are blocked at the two-cell stage with

an anucleolated nucleus (Fig. 7). Therefore, selective enucleation as countered by enucleolation revealed the influence of GV contents on pronuclear formation. Furthermore, complete enucleation of denuded GV oocytes followed by G2/M embryonic or fetal fibroblast nuclear transfer indicate maturation with abnormalities such as acceleration of polar body extrusion, large polar body, and misalignment of chromosomes over the MII spindle, in addition to blocked embryo development at the one-cell stage after activation or fertilization. Surprisingly, no nucleoli were observed in the developed embryos. In addition, upon fertilization, one pronucleus was formed in the cytoplasm, and was expected to be somatic, whereas the male pronucleus was not formed and was found as scattered chromosomes in the cytoplasm (Fig. 8). This expectation is based on the fact that the pronucleus close to the polar body is somatic and the pronucleus farthest from the polar body is spermic. This was confirmed through maturation of GV cytoplasts for 17 h followed by fertilization. Fertilizing sperm in the cytoplasm remained in a highly condensed head form (Mohammed *et al.*, 2008, 2010, 2019, 2022).




Fig. 7. Pronuclei (arrow) in the zygote (A) and nuclei (arrow) in 2-cell embryo (B) obtained from enucleolated and fertilized fully grown germinal vesicle oocytes. Nucleoli are not visible both in pronuclei and in blastomere nuclei. In the zygote the second pronucleus is out of focus.




Fig. 8. Pronuclear morphology in GV (A) and GVsel cytoplasts (B and C) reconstructed with embryonic/somatic nuclei and fertilized after *in vitro* maturation (light microscope, DIC). (A) One pronucleus formed of somatic origin without nucleoli (arrow) and extruded large PB; (B) Two pronuclei were formed of somatic and sperm origin and both contained nucleoli; (C) stained two pronuclei were formed of somatic (green arrow) and sperm (red arrow) origin and both contained nucleoli.




Fig. 9. Effects of nuclear material and cumulus cells on developmental competence of cytoplasts reconstructed with  $\frac{1}{2}$ -blastomere nuclei. GV removal the whole germinal vesicle of GV oocyte, GVsel removal nuclear material and membrane leaving nucleolus and nuclear sap in the cytoplasm of denuded GV oocytes, GVCsel removal nuclear material and membrane leaving nucleolus and nuclear sap in the cytoplasm of cumulus enclosed GV oocytes.




Fig. 10. Hatching blastocysts developed from GVCsel cytoplasts reconstructed with  $\frac{1}{2}$ -blastomere nuclei at G2/M phase (A) two hatching blastocysts under light microscope (DIC) and (B) one hatching blastocyst after Hoechst staining (fluorescence microscope).

Upon selective enucleation of denuded GV oocytes, reconstructed with G2/M embryonic nucleus, followed by maturation and fertilization, both pronuclei were formed in addition to further developmental competence to morula/blastocyst (Fig. 9 and 10). Although activated reconstructed GVsel cytoplasts were also cleaved and few of them developed to the blastocyst stage the frequency of cleavage to the 4-cell stage and blastocyst formation was evidently lower than those of reconstructed GVCsel cytoplasts (38.5% vs 71.4% and 7.7% and 23.8%, respectively). Also, the mean cell number in the obtained blastocysts was slightly higher in blastocysts derived from activated reconstructed GVCsel cytoplasts. What is deserved to observe is the absence or lower cells of inner cell mass in the cloned developed blastocysts over somatic or embryonic nuclear transfer to GVCsel cytoplasts compared to normal developed blastocysts. The number

and quality of inner cell mass is the most important character of the developed blastocysts to give birth. Zhu *et al.* (2022) reported that the poor morphology of inner cell mass (grade C) of the developed embryos increases birth weight and large for gestational age compared to inner cell mass of grade A. In addition, Plana-Carmona *et al.* (2022) confirmed that the trophectoderm acts as a niche for the inner cell mass through C/EBP $\alpha$ -regulated IL-6 signaling. Therefore, further studies are required for investigating the reasons of absence or poor quality inner cell mass of the developed blastocysts from GV cytoplasts after embryonic nuclear transfer.

This very interesting technique of enucleation denuded or cumulus-enclosed GV oocytes could be applied with visible GV nucleus as in human, rodent and rabbit oocytes. In ruminant GV oocytes, ultra-centrifugation to 15,000 rpm is required for visualization of GV nucleus due to lipid droplets and the cumulus is stripped off during such centrifugation. In addition, the GV nucleus could be visible approximately in two third of the centrifuged GV oocytes.

Concerning the cell cycle stage of the introduced donor nuclei, our results and others indicate that G0/G1 cells or nuclei could be transferred to metaphase II cytoplasts, whereas G2/M cells or nuclei could be transferred to germinal vesicle cytoplasts (Wilmot *et al.*, 1997; Mohammed *et al.*, 2008, 2010, 2019, 2022). Donor cells or nuclei introduced in the S phase to the germinal vesicle cytoplasm result in abnormalities in oocyte maturation and block further developmental competence of embryos. To the best of our knowledge, embryonic or somatic nuclear transfer to metaphase II cytoplasts results in deliveries with low efficiency (Campbell *et al.*, 1996; Wani *et al.*, 2018), whereas with embryonic or somatic nuclear transfer to GVCsel cytoplasm, the embryonic development reached to the hatched blastocyst stage (Mohammed *et al.*, 2008, 2010). Upon morula and blastocysts transferred to the uterine of surrogate mothers, no delivery was confirmed in mice (Mohammed unpublished data).

Our complete and selective enucleation of GV yielded GV cytoplasts with high efficiency (Wilmot *et al.*, 1997; Mohammed *et al.*, 2008, 2010, 2019, 2022). The sole success of the recipient germinal vesicle cytoplasm is the meiotic maturation of germinal vesicle karyoplasts; germ cells; primary or secondary spermatogonia (Ogura *et al.*, 1998; Zhang *et al.*, 2015). Germinal vesicle recipient cytoplasts are required in basic research for investigations of cell biology and donor cell reprogramming. Further studies are still required for improvement the outcomes of manipulation germinal vesicle oocytes in different animal species and human as well.

## CONCLUSIONS

Oocytes grow and develop during fetal and postnatal periods within several distinct ovarian follicles, and they ovulate in each estrous or menstrual cycle from ovarian follicles after puberty. Trials were unsuccessful when germ cells were used for *in vitro* embryo production during the fetal period; however, they succeeded when germ cells were used of the prepubertal and post pubertal periods for *in vitro* embryo production. The higher the follicle diameter and oocyte quality, the higher the maturation, fertilization, and developmental competence of the embryo. Estrous or menstrual cycle synchronization and superovulation protocols for ovarian follicles have been used in different species for embryo transfer. In addition, ovarian tissue transplantation has been adopted for the treatment of infertility and conservation of species. Furthermore, defects in oocytes can be repaired via the nucleolus, germinal vesicle, and spindle transfer to the recipient oocyte or cytoplasm. Primary and secondary spermatocytes and spermatids have been successfully used to fertilize GV and MII oocytes. Furthermore, germinal vesicle, embryonic, and somatic cells at different cell cycle stages (G0, S, and G/M stages) have been transferred to different cytoplasts (GV, ProMI, MI, and MII stages) with variable results. Assisted reproductive techniques (ART) are used to treat infertility, save endangered species, enhance meat and milk production through *in vitro* manipulations of germ cells, follicles, oocytes, and sperm. Studies on the kinetics or progression of oocyte maturation and further developmental competence of embryos are required, in addition to ovarian follicle gene expression. Although assisted reproductive techniques are still relatively rare and expensive, especially in third-world countries, their use has doubled over the past decade and thus necessitates further development.

## DECLARATIONS

### Acknowledgments

The authors would like to acknowledge the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University, Saudi Arabia (GrantA051).

### Funding

The study was funded by Funding of Scientific Research Deanship of King Faisal University (GrantA051)

### IRB approval

The approval of the study was granted by the The Ethical Research Committee of King Faisal University.

### Ethical statement

The procedures and material of the current study was approved of the Scientific Research Deanship Ethical Standards of King Faisal University.

### Statement of conflict of interest

The authors have declared no conflict of interest.

## REFERENCES

- Abdelnaby, E.A., El-Maaty, A.M.A. and El-Badry, D.A., 2021. Evaluation of ovarian hemodynamics by color and spectral Doppler in cows stimulated with three sources of follicle-stimulating hormone. *Reprod. Biol.*, **21**: 100478. <https://doi.org/10.1016/j.repbio.2020.100478>
- Abdoon, A.S., Fathalla, S.I., Shawky, S.M., Kandil, O.M., Kishta, A.A. and Masoud, S.R., 2018. *In vitro* maturation and fertilization of donkey oocytes. *J. Equine Vet. Sci.*, **65**: 118–122. <https://doi.org/10.1016/j.jevs.2018.04.007>
- Al-Masruri, H., Al-Zeidi, R., Al-Mufarji, A., Mohammed, A.A., Al-Madani, A. and Mohammed, H., 2022. Leverage of *Moringa oleifera* supplementation on performances, biochemical, and milk profiles in mammals. *Adv. Anim. Vet. Sci.*, **10**: 2043-2050. <https://doi.org/10.17582/journal.aavs/2022/10.9.2043.2050>
- Al-Masruri, H., Al-Zeidi, R., Al-Mufarji, A. and Mohammed, A.A., 2022a. *Moringa oleifera* leaves supplementation to anaesthetized mice associated with changes of thermo-tolerance parameters, blood and plasma indices. *Fresen. Environ. Bull.*, **8**: 8179-8187.
- Al-Mufarji, A., Mohammed, A.A., Al-Zeidi, R., Al-Masruri, H. and Mohammed, A., 2022b. Effects of *Moringa oleifera* on follicular development, blood and metabolic profiles of subtropical ewes during peripartum. *Adv. Anim. Vet. Sci.*, **10**: 1706-1712. <https://doi.org/10.17582/journal.aavs/2022/10.8.1706.1712>
- Al-Zeidi, R., Al-Masruri, H., Al-Mufarji, A. and Mohammed, A.A., 2022a. Role of cumulus cells and follicular fluid on oocyte maturation and developmental competence of embryos: Intact and reconstructed oocytes. *Adv. Anim. Vet. Sci.*, **10**: 1219-1226. <https://doi.org/10.17582/journal.aavs/2022/10.6.1219.1226>
- Al-Zeidi, R., Al-Masruri, H., Al-Mufarji, A., Mohammed, A.A. and Mohammed, H., 2022b. Changes in nucleus and cytoplasm during oocyte maturation: Involvement in embryo production.

- Adv. Anim. Vet. Sci.*, **10**: 2081-2089. <https://doi.org/10.17582/journal.aavs/2022/10.9.2081.2089>
- Algriany, O.A., Vos, P., Groenendaal, H., van Gastel, A., Sirard, M.A. and Dieleman, S.J., 2007. *Effects of oFSH stimulation on steroid concentrations in the fluid of preovulatory follicles during final maturation in relation to the LH surge in the cow*. Expression analysis of genes implicated in meiotic resumption in vivo and developmental competence of bovine oocytes, pp. 33.
- An, L., Liu, J., Du, Y., Liu, Z., Zhang, F., Liu, Y., Zhu, X., Ling, P., Chang, S. and Hu, Y. 2018. Synergistic effect of cysteamine, leukemia inhibitory factor, and Y27632 on goat oocyte maturation and embryo development *in vitro*. *Theriogenology*, **108**: 56–62. <https://doi.org/10.1016/j.theriogenology.2017.11.028>
- Arias-Álvarez, M., García-García, R.M., López-Tello, J., Rebollar, P.G., Gutiérrez-Adán, A. and Lorenzo, P.L., 2017. *In vivo* and *in vitro* maturation of rabbit oocytes differently affects the gene expression profile, mitochondrial distribution, apoptosis and early embryo development. *Reprod. Fertil. Dev.*, **29**: 1667–1679. <https://doi.org/10.1071/RD15553>
- Ata, B. and Telek, S.B., 2021. Assisted reproductive technology for women with endometriosis, a clinically oriented review. *Curr. Opin. Obstet. Gynecol.*, **33**: 225-231. <https://doi.org/10.1097/GCO.0000000000000710>
- Baby, T.E. and Bartlewski, P.M., 2011. Circulating concentrations of ovarian steroids and follicle-stimulating hormone (FSH) in ewes with 3 or 4 waves of antral follicle emergence per estrous cycle. *Reprod. Biol.*, **11**: 19–36. [https://doi.org/10.1016/S1642-431X\(12\)60061-8](https://doi.org/10.1016/S1642-431X(12)60061-8)
- Baerwald, A. and Pierson, R., 2020. Ovarian follicular waves during the menstrual cycle: Physiologic insights into novel approaches for ovarian stimulation. 2020. *Fertil. Steril.*, **114**: 443–457. <https://doi.org/10.1016/j.fertnstert.2020.07.008>
- Bao, S., Hitoshi, U., Hirose, A., Aono, F., Ono, Y. and Kono, T., 2003. Development of bovine oocytes reconstructed with a nucleus from growing stage oocytes after fertilization *in vitro*. *Theriogenology*, **59**: 1231-1239. [https://doi.org/10.1016/S0093-691X\(02\)01174-3](https://doi.org/10.1016/S0093-691X(02)01174-3)
- Baruffi, R.L., Avelino, K.B., Petersen, C.G., Mauri, A.L., Garcia, J.M. and Franco, J.G., 2004. Nuclear and cytoplasmic maturation of human oocytes cultured *in vitro*. *Fertil. Steril.*, **82**: S265–S266. <https://doi.org/10.1016/j.fertnstert.2004.07.709>
- Benc, M., Strejček, F., Morović, M., Murín, M., Martínková, S., Jettmarová, D. and Laurinčík, J., 2019. Enucleolation and nucleolus transfer in mammalian oocytes and zygotes. *Int. J. Dev. Biol.*, **63**: 253–258. <https://doi.org/10.1387/ijdb.190002mb>
- Blondin, P., Coenen, K., Guilbault, L.A. and Sirard, M.A., 1997. *In vitro* production of bovine embryos: Developmental competence is acquired before maturation. *Theriogenology*, **47**: 1061–1075. [https://doi.org/10.1016/S0093-691X\(97\)00063-0](https://doi.org/10.1016/S0093-691X(97)00063-0)
- Borges, M.A., Sousa, F.S.S., Paschoal, J.D., Lopes, I.A.R., Feijó, A.L.S., Neto, A.C.P.S., Silva, P.L., Seixas, F.K. and Collares, T., 2021. Effect of supplementation of medium with Bauhinia for ficata recombinant lectins on expression of oxidative stress genes during *in vitro* maturation of bovine oocytes. *Reprod. Toxicol.*, **103**: 64–70. <https://doi.org/10.1016/j.reprotox.2021.05.012>
- Bunel, A., Nivet, A.L., Blondin, P., Vigneault, C., Richard, F.J. and Sirard, M.A., 2020. The effects of LH inhibition with cetrorelix on cumulus cell gene expression during the luteal phase under ovarian coasting stimulation in cattle. *Domest. Anim. Endocrinol.*, **72**: 106429. <https://doi.org/10.1016/j.domaniend.2019.106429>
- Campbell, K.H., Loi, P., Otaegui, P.J. and Wilmut, I., 1996. Cell cycle co-ordination in embryo cloning by nuclear transfer. *Rev. Reprod.*, **1**: 40–46. <https://doi.org/10.1530/ror.0.0010040>
- Castaneda, C.A., Kaye, P., Pantaleon, M., Phillips, N., Norman, S., Fry, R. and Michael, J.D., 2013. Lipid content, active mitochondria and brilliant cresyl blue staining in bovine oocytes. *Theriogenology*, **79**: 417–422. <https://doi.org/10.1016/j.theriogenology.2012.10.006>
- Cavalieri, F.L.B., Morotti, F., Seneda, M.M., Colombo, A.H.B., Andreazzi, M.A., Emanuelli, I.P. and Rigolon, L.P., 2018. Improvement of bovine *in vitro* embryo production by ovarian follicular wave synchronization prior to ovum pick-up. *Theriogenology*, **117**: 57–60. <https://doi.org/10.1016/j.theriogenology.2017.11.026>
- Chakravarthi, V.P., Ratri, A., Masumi, S., Borosha, S., Ghosh, S., Christenson, L.K., Roby, K.F., Wolfe, M. W. and Rumi, M.A.K., 2021. Granulosa cell genes that regulate ovarian follicle development beyond the antral stage: The role of estrogen receptor  $\beta$ . *Mol. Cell. Endocrinol.*, **528**: 111212. <https://doi.org/10.1016/j.mce.2021.111212>
- Chang, H.C., Liu, H., Zhang, J., Grifo, J. and Krey, L.C., 2005. Developmental incompetency of denuded mouse oocytes undergoing maturation in vitro is

- ooplasmic in nature and is associated with aberrant Oct-4 expression. *Hum. Reprod.*, **20**: 1958–1968. <https://doi.org/10.1093/humrep/dei003>
- Chian, R.C., Buckett, W.M. and Tan, S.L., 2004. *In-vitro* maturation of human oocytes. *Reprod. Biomed. Online*, **8**: 148–166. [https://doi.org/10.1016/S1472-6483\(10\)60511-1](https://doi.org/10.1016/S1472-6483(10)60511-1)
- Dai, X.X., Jiang, Z.Y., Wu, Y.W., Sha, Q.Q., Liu, Y., Ding, J.Y., Xi, W.D., Li, J. and Fan, H.Y. 2021. CNOT6/6L-mediated mRNA degradation in ovarian granulosa cells is a key mechanism of gonadotropin-triggered follicle development. *Cell Rep.*, **37**: 110007. <https://doi.org/10.1016/j.celrep.2021.110007>
- de Senna Costa, J.A., Cezar, G.A., Monteiro, P.L.J., Silva, D.M.F., Silva, R.A.J.A., Bartolomeu, C.C., dos Santos Filho, A.S., Wischral, A. and Batista, A.M., 2022. Leptin improves *in-vitro* maturation of goat oocytes through MAPK and JAK2/STAT3 pathways and affects gene expression of cumulus cells. *Reprod. Biol.*, **22**: 100609. <https://doi.org/10.1016/j.repbio.2022.100609>
- Demyda-Peyrás, S., Dorado, J., Hidalgo, M., Anter, J., de Luca, L., Genero, E. and Moreno-Millán, M., 2013. Effects of oocyte quality, incubation time and maturation environment on the number of chromosomal abnormalities in IVF-derived early bovine embryos. *Reprod. Fertil. Dev.*, **25**: 1077–1084. <https://doi.org/10.1071/RD12140>
- Dolma, P., Ali, A., Sobti, R.C. and Chauhan, A., 2022. Animal cloning: Perspectives for futuristic medicine. In: *Advances in animal experimentation and modeling*, Elsevier. pp. 53–67. <https://doi.org/10.1016/B978-0-323-90583-1.00008-8>
- Fernandes, C.C.L., Martins, L.T., Gaudêncio Neto, S., Tavares, K.C.S., Aguiar, L.H., Calderón, C.E.M., Silva, A.M., Alves, J.P.M., Silva, C.M.G., Rossetto, R., Bertolini, L.R., Bertolini, M. and Rondina, D., 2016. Gene expression, oocyte quality and embryo production by cloning in goats supplemented with different diets. *Small Rumin. Res.*, **144**: 255–262. <https://doi.org/10.1016/j.smallrumres.2016.09.023>
- Fulka, H., Mrazek, M. and Fulka Jr, J., 2004. Nucleolar dysfunction may be associated with infertility in humans. *Fertil. Steril.*, **82**: 486–487. <https://doi.org/10.1016/j.fertnstert.2003.12.042>
- Furukawa, E., Chen, Z., Ueshiba, H., Wu, Y., Chiba, H., Yanagawa, Y., Katagiri, S., Nagano, M. and Hui, S.P., 2021. Postpartum cows showed high oocyte triacylglycerols concurrently with high plasma free fatty acids. *Theriogenology*, **176**: 174–182. <https://doi.org/10.1016/j.theriogenology.2021.09.034>
- Gad, A., Hamed, S.A., Khalifa, M., Amin, A., El-Sayed, A., Swiefy, S.A. and El-Assal, S., 2018. Retinoic acid improves maturation rate and upregulates the expression of antioxidant-related genes in *in vitro* matured buffalo (*Bubalus bubalis*) oocytes. *Int. J. Vet. Sci. Med.*, **6**: 279–285. <https://doi.org/10.1016/j.ijvsm.2018.09.003>
- Girsh, E., 2021. Chapter 8- ovum pickup (OPU) from section 2- assisted reproductive procedures. In: *A textbook of clinical embryology*. Cambridge University Press. pp. 89–94. <https://doi.org/10.1017/9781108881760.009>
- Gong, S., Jiang, L., Cheng, H., Pan, L.Z., Xu, M.T., Zhang, M., Yuan, H.J. and Tan, J.H., 2022. Effects of CRH and ACTH exposure during *in vitro* maturation on competence of pig and mouse oocytes. *Theriogenology*, **173**: 211–220. <https://doi.org/10.1016/j.theriogenology.2021.06.025>
- Gordon, I., 2003. Establishing pregnancies with IVP Embryos. In: *Laboratory production of cattle embryos*. (2<sup>nd</sup>), pp. 303–321. <https://doi.org/10.1079/9780851996660.0303>
- Grabarek, J.B., Płusa, B., Modliński, J.A. and Karasiewicz, J., 2004. Reconstruction of enucleated mouse germinal vesicle oocytes with blastomere nuclei. *Zygote*, **12**: 163–172. <https://doi.org/10.1017/S0967199404002746>
- Gutiérrez-Añez, J.C., Lucas-Hahn, A., Haderer, K.G., Aldag, P. and Niemann, H., 2021. Melatonin enhances *in vitro* developmental competence of cumulus-oocyte complexes collected by ovum pick-up in prepubertal and adult dairy cattle. *Theriogenology*, **161**: 285–293. <https://doi.org/10.1016/j.theriogenology.2020.12.011>
- Hashimoto, S., Minami, N., Takakura, R. and Imai, H., 2002. Bovine immature oocytes acquire developmental competence during meiotic arrest *in vitro*. *Biol. Reprod.*, **66**: 1696–1701. <https://doi.org/10.1095/biolreprod66.6.1696>
- Hasler, J.F., 1998. The current status of oocyte recovery, *in vitro* embryo production, and embryo transfer in domestic animals, with an emphasis on the bovine. *J. Anim. Sci.*, **76**: 52–74. [https://doi.org/10.2527/1998.76suppl\\_352x](https://doi.org/10.2527/1998.76suppl_352x)
- Hassold, T. and Chiu, D., 1985. Maternal age-specific rates of numerical chromosome abnormalities with special reference to trisomy. *Hum. Genet.*, **70**: 11–17. <https://doi.org/10.1007/BF00389450>
- Hoseini, F.S., Salsabili, N., Akbari-Asbagh, F., Aflatoonian, R. and Aghaee-Bakhtiari, S.H., 2016. Comparison of gene expression profiles in human germinal vesicle before and after cytoplasmic

- transfer from mature oocytes in Iranian infertile couples. *J. Family Reprod. Hlth.*, **10**: 71.
- Jewgenow, K., Fernandez-Gonzalez, L., Jänsch, S., Viertel, D. and Zahmel, J., 2019. Brilliant cresyl blue staining allows the selection for developmentally competent immature feline oocytes. *Theriogenology*, **126**: 320–325. <https://doi.org/10.1016/j.theriogenology.2018.12.021>
- Jin, J.I., Ghanem, N., Kim, S.S., Choi, B.H., Ha, A.N., Lee, K.L., Sun, D.W., Lim, H.T., Lee, J.G. and Kong, I.K., 2016. Interaction of donor age, parity and repeated recovery of cumulus–oocyte complexes by ovum pick-up on *in vitro* embryo production and viability after transfer. *Livest. Sci.*, **188**: 43–47. <https://doi.org/10.1016/j.livsci.2016.04.002>
- Kanitz, W., Becker, F., Schneider, F., Kanitz, E., Leiding, C., Nohner, H.P. and Pöhlend, R., 2002. Superovulation in cattle: Practical aspects of gonadotropin treatment and insemination. *Reprod. Nutr. Dev.*, **42**: 587–599. <https://doi.org/10.1051/rnd:2002045>
- Ko, Y., Kim, J.H., Lee, S.R., Kim, S.H. and Chae, H.D., 2021. Influence of pretreatment of insulin on the phosphorylation of extracellular receptor kinase by gonadotropin-releasing hormone and gonadotropins in cultured human granulosa cells. *Eur. J. Obstet. Gynecol. Reprod. Biol.*, **262**: 113–117. <https://doi.org/10.1016/j.ejogrb.2021.05.016>
- Korzekwa, A.J. and Kotlarczyk, A.M., 2021. Artificial reproductive technology (ART) applied to female cervids adapted from domestic ruminants. *Animals*, **11**: 2933. <https://doi.org/10.3390/ani11102933>
- Kumar, P., Rajput, S., Verma, A., De, S. and Datta, T.K., 2013. Expression pattern of glucose metabolism genes in relation to development rate of buffalo (*Bubalus bubalis*) oocytes and *in vitro*–produced embryos. *Theriogenology*, **80**: 914–922. <https://doi.org/10.1016/j.theriogenology.2013.07.018>
- Lee, S.H., Oh, H.J., Kim, M.J., Kim, G.A., Choi, Y. Bin, Jo, Y.K. and Lee, B.C. 2018. Effect of co-culture canine cumulus and oviduct cells with porcine oocytes during maturation and subsequent embryo development of parthenotes *in vitro*. *Theriogenology*, **106**: 108–116. <https://doi.org/10.1016/j.theriogenology.2017.09.015>
- Leibfried-Rutledge, M.L., Critser, E.S., Eyestone, W.H., Northey, D.L. and First, N.L., 1987. Development potential of bovine oocytes matured *in vitro* or *in vivo*. *Biol. Reprod.*, **36**: 376–383. <https://doi.org/10.1095/biolreprod36.2.376>
- Li, M., Yan, J., Zhi, X., Wang, Y., Hang, J. and Qiao, J., 2019. Gene expression during oogenesis and oocyte development. In: *The Ovary*. Elsevier, pp. 205–216. <https://doi.org/10.1016/B978-0-12-813209-8.00013-3>
- Li, Q., Zhang, Y., Li, W., Yan, K., Liu, Y., Xu, H., Lu, Y., Liang, X. and Yang, X., 2022. Allicin protects porcine oocytes against LPS-induced defects during maturation *in vitro*. *Theriogenology*, **182**: 138–147. <https://doi.org/10.1016/j.theriogenology.2022.02.007>
- Liang, S., Kang, J., Jin, H., Liu, X., Li, J., Li, S., Lu, Y., Wang, W. and Yin, X.J., 2012. The influence of 9-cis-retinoic acid on nuclear and cytoplasmic maturation and gene expression in canine oocytes during *in vitro* maturation. *Theriogenology*, **77**: 1198–1205. <https://doi.org/10.1016/j.theriogenology.2011.10.027>
- Liu, W., Guo, J., Li, C., Liao, H., Qin, Y. and Huang, G., 2021. Mitochondrial DNA copy number of cumulus cells is not linked to embryo implantation in good prognosis IVF patients. *Reprod. Biomed. Online*, **42**: 901–908. <https://doi.org/10.1016/j.rbmo.2021.01.002>
- Lopes, G., Rocha, A. and Sousa, M., 2010. Comparative ultrastructural analysis of diestrous and anestrous canine Grade 1 cumulus–oocyte complexes. *Anim. Reprod. Sci.*, **122**: 244–252. <https://doi.org/10.1016/j.anireprosci.2010.08.018>
- Luo, Y., Che, M.J., Liu, C., Liu, H.G., Fu, X.W. and Hou, Y.P., 2018. Toxicity and related mechanisms of dihydroartemisinin on porcine oocyte maturation *in vitro*. *Toxicol. appl. Pharmacol.*, **341**: 8–15. <https://doi.org/10.1016/j.taap.2018.01.002>
- Luu, V.V., Namula, Z., Kaedei, Y., Tanihara, F. and Otoi, T., 2011. Effects of ovary storage time on the quality and meiotic competence of cat oocytes. *Reprod. Fertil. Dev.*, **24**: 205–206. <https://doi.org/10.1071/RDv24n1Ab187>
- Ma, L., Cai, L., Hu, M., Wang, J., Xie, J., Xing, Y., Shen, J., Cui, Y., Liu, X.J. and Liu, J., 2020. Coenzyme Q10 supplementation of human oocyte *in vitro* maturation reduces postmeiotic aneuploidies. *Fertil. Steril.*, **114**: 331–337. <https://doi.org/10.1016/j.fertnstert.2020.04.002>
- Madkour, A., Bouamoud, N., Kaarouch, I., Louanjli, N., Saadani, B., Assou, S., Aboulmaouahib, S., Sefrioui, O., Amzazi, S. and Copin, H., 2018. Follicular fluid and supernatant from cultured cumulus–granulosa cells improve *in vitro* maturation in patients with polycystic ovarian syndrome. *Fertil. Steril.*, **110**: 710–719. <https://doi.org/10.1016/j.fertnstert.2018.04.038>
- Margalit, T., Ben-Haroush, A., Garor, R., Kotler, N.,

- Shefer, D., Krasilnikov, N., Tzabari, M., Oron, G., Shufaro, Y. and Sapir, O., 2019. Morphokinetic characteristics of embryos derived from *in-vitro* matured oocytes and their *in-vivo* matured siblings after ovarian stimulation. *Reprod. Biomed. Online*, **38**: 7–11. <https://doi.org/10.1016/j.rbmo.2018.10.002>
- Marin, D.F.D., da Costa, N.N., Santana, P.P.B., de Souza, E.B. and Ohashi, O.M., 2019. Importance of lipid metabolism on oocyte maturation and early embryo development: Can we apply what we know to buffalo? *Anim. Reprod. Sci.*, **211**: 106220. <https://doi.org/10.1016/j.anireprosci.2019.106220>
- Maylinda, S., Sarah, O.L. and Busono, W., 2018. Role of seasons on the production and reproduction performance of Kacang goats (*Kambing kacang*) in north middle Timor regency. *Ind. J. Anim. Res.*, **52**: 1-5. <https://doi.org/10.18805/ijar.B-668>
- Miao, J.K., Liu, Y.H., Liu, S., Liu, X.M., Wang, P.C., Du, Z.Q. and Yang, C.X., 2019. Lysosomal dysfunction disturbs porcine oocyte maturation and developmental capacity by disorganizing chromosome/cytoskeleton and activating autophagy/apoptosis. *Theriogenology*, **140**: 44-51. <https://doi.org/10.1016/j.theriogenology.2019.08.019>
- Mohammed, A.A. and Al-Hozab, A.A., 2020. + (-) catechin raises body temperature, changes blood parameters, improves oocyte quality and reproductive performance of female mice. *Ind. J. Anim. Res.*, **54**: 543–548. <https://doi.org/10.18805/ijar.B-981>
- Mohammed, A.A. and Attaai, A.H., 2011. Effects of dietary urea on timing of embryo cleavages and blood components in mice. *Vet. World*, **4**: 360. <https://doi.org/10.5455/vetworld.2011.360-363>
- Mohammed, A.A., Karasiewicz, J., Kubacka, J., Gręda, P. and Modliński, J.A., 2010. Enucleated GV oocytes as recipients of embryonic nuclei in the G1, S, or G2 stages of the cell cycle. *Cell. Reprogram.*, **12**: 427–435. <https://doi.org/10.1089/cell.2009.0107>
- Mohammed, A.A., Karasiewicz, J. and Modliński, J.A., 2008. Developmental potential of selectively enucleated immature mouse oocytes upon nuclear transfer. *Mol. Reprod. Dev.*, **75**: 1269–1280. <https://doi.org/10.1002/mrd.20870>
- Mohammed, A.A., Karasiewicz, J., Papis, K. and Modliński, J.A., 2005. Oocyte maturation in the presence of randomly pooled follicular fluid increases bovine blastocyst yield *in vitro*. *J. Anim. Feed Sci.*, **14**: 501. <https://doi.org/10.22358/jafs/67048/2005>
- Mohammed, A.A., 2006. Meiotic maturation and developmental competence of mouse oocytes reconstructed with G2/M somatic nuclei. In: *Biotechnology*. Scientific Redagogical Publishing, C. Budejovice, pp. 1117-1120.
- Mohammed, A.A., 2018. Development of oocytes and preimplantation embryos of mice fed diet supplemented with dunaliella salina. *Adv. Anim. Vet. Sci.*, **6**: 33-39. <https://doi.org/10.17582/journal.aavs/2018/6.1.33.39>
- Mohammed, A.A., 2019. *Nigella sativa* oil improves physiological parameters, oocyte quality after ovarian transplantation, and reproductive performance of female mice. *Pakistan J. Zool.*, **51**: 2225-2231. <https://doi.org/10.17582/journal.pjz/2019.51.6.2225.2231>
- Mohammed, A.A., Al-Mufarji, A. and Alawaid, S., 2022. Developmental potential of ovarian follicles in mammals: involvement in assisted reproductive techniques. *Pakistan J. Zool.*, **54**: 2919-2929. <https://doi.org/10.17582/journal.pjz/20220128140127>
- Mohammed, A.A., Al-Suwaiegh, S. and Al-Shaheen, T., 2019. Do the cytoplasm and nuclear material of germinal vesicle oocyte support developmental competence upon reconstruction with embryonic/somatic nucleus. *Cell. Reprogram.*, **21**: 163–170. <https://doi.org/10.1089/cell.2019.0032>
- Moulavi, F. and Hosseini, S.M., 2019. Effect of macromolecule supplement on nuclear and cytoplasmic maturation, cryosurvival and *in vitro* embryo development of dromedary camel oocytes. *Theriogenology*, **132**: 62–71. <https://doi.org/10.1016/j.theriogenology.2019.03.027>
- O'Connell, J.M. and Pepling, M.E., 2021. Primordial follicle formation. Some assembly required. *Curr. Opin. Endocr. Metab. Res.*, **18**: 118–127. <https://doi.org/10.1016/j.coemr.2021.03.005>
- Oberlander, G., Murgas, L.D.S., Zangeronimo, M.G., da Silva, A.C., de Alcantara Menezes, T., Pontelo, T.P., and Vieira, L.A., 2013. Role of insulin-like growth factor-I and follicular fluid from ovarian follicles with different diameters on porcine oocyte maturation and fertilization *in vitro*. *Theriogenology*, **80**: 319–327. <https://doi.org/10.1016/j.theriogenology.2013.04.018>
- Ogura, A., Suzuki O., Tanemura K., Mochida K., Kobayashi Y. and Matsuda J., 1998. Development of normal mice from metaphase I oocytes fertilized with primary spermatocytes. *Proc. natl. Acad. Sci. USA*, **12**: 5611–5615. <https://doi.org/10.1073/pnas.95.10.5611>
- Ongaratto, F.L., Cedeño, A.V., Rodriguez-Villamil,

- P., Tribulo, A. and Bó, G.A., 2020. Effect of FSH treatment on cumulus oocyte complex recovery by ovum pick up and in vitro embryo production in beef donor cows. *Anim. Reprod. Sci.*, **214**: 106274. <https://doi.org/10.1016/j.anireprosci.2020.106274>
- Opiela, J. and Kątska-Książkiewicz, L., 2013. The utility of Brilliant Cresyl Blue (BCB) staining of mammalian oocytes used for *in vitro* embryo production (IVP). *Reprod. Biol.*, **13**: 177–183. <https://doi.org/10.1016/j.repbio.2013.07.004>
- Patton, B.K., Madadi, S. and Pangas, S.A., 2021. Control of ovarian follicle development by TGF- $\beta$  family signaling. *Curr. Opin. Endocr. Metab. Res.*, **18**: 102–110. <https://doi.org/10.1016/j.coemr.2021.03.001>
- Pincus, G. and Enzmann, E.V., 1935. The comparative behavior of mammalian eggs *in vivo* and *in vitro*: I. The activation of ovarian eggs. *J. exp. Med.*, **62**: 665–675. <https://doi.org/10.1084/jem.62.5.665>
- Plana-Carmona, M., Stik, G., Bulteau, R., Segura-Morales, C., Alcázar, N., Wyatt, C., Klonizakis, A., Andrés-Aguayo, L., Gasnier, M., Tian, T.V., Garcia, G.T., Vila-Casadesús, M., Plachta, N., Serrano, M., Francesconi, M. and Graf, T., 2022. The trophectoderm acts as a niche for the inner cell mass through C/EBP $\alpha$ -regulated IL-6 signaling. *Stem Cell Reports*, **17**: 1991–2004. <https://doi.org/10.1016/j.stemcr.2022.07.009>
- Polanski, Z., Hoffmann, S. and Tsurumi, C., 2005. Oocyte nucleus controls progression through meiotic maturation. *Dev. Biol.*, **281**: 184–195. <https://doi.org/10.1016/j.ydbio.2005.02.024>
- Pournaghi, M., Khodavirdilou, R., Saadatlou, M.A.E., Nasimi, F.S., Yousefi, S., Mobarak, H., Darabi, M., Shahnazi, V., Rahbarghazi, R. and Mahdipour, M., 2021. Effect of melatonin on exosomal dynamics in bovine cumulus cells. *Process Biochem.*, **106**: 78–87. <https://doi.org/10.1016/j.procbio.2021.03.008>
- Raj, M.P., Naidu, G.V., Srinivas, M., Raghunath, M. and Rao, K.A., 2018. Effect of preovulatory follicle on fertility in Graded murrah buffaloes (*Bubalus bubalis*). *Ind. J. Anim. Res.*, **52**: 834–838. <https://doi.org/10.18805/ijar.B-3302>
- Residiwati, G., Azari-Dolatabad, N., Tuska, H.S.A., Sidi, S., van Damme, P., Benedetti, C., Montoro, A.F., Luceno, N.L., Pavani, K.C. and Opsomer, G., 2021. Effect of lycopene supplementation to bovine oocytes exposed to heat shock during *in vitro* maturation. *Theriogenology*, **173**: 48–55. <https://doi.org/10.1016/j.theriogenology.2021.07.014>
- Richards, J.S. and Pangas, S.A., 2010. The ovary: Basic biology and clinical implications. *J. clin. Investig.*, **120**: 963–972. <https://doi.org/10.1172/JCI41350>
- Saini, S., Sharma, V., Ansari, S., Kumar, A., Thakur, A., Malik, H., Kumar, S. and Malakar, D., 2022. Folate supplementation during oocyte maturation positively impacts the folate-methionine metabolism in pre-implantation embryos. *Theriogenology*, **182**: 63–70. <https://doi.org/10.1016/j.theriogenology.2022.01.024>
- Sakaguchi, K. and Nagano, M., 2020. Follicle priming by FSH and pre-maturation culture to improve oocyte quality *in vivo* and *in vitro*. *Theriogenology*, **150**: 122–129. <https://doi.org/10.1016/j.theriogenology.2020.01.023>
- Sallem, A., Denizot, A.L., Ziyat, A., L'Hostis, A., Favier, S., Burlet, P., Lapierre, J.M., Dimby, S.F., Patrat, C. and Sifer, C., 2022. A fertilin-derived peptide improves *in vitro* maturation and ploidy of human oocytes. *F and S Sci.*, **3**: 21–28. <https://doi.org/10.1016/j.xfss.2021.10.004>
- Sánchez-Ajofrín, I., Iniesta-Cuerda, M., Sánchez-Calabuig, M.J., Peris-Frau, P., Martín-Maestro, A., Ortiz, J. A., del Rocío Fernández-Santos, M., Garde, J.J., Gutiérrez-Adán, A. and Soler, A.J., 2020. Oxygen tension during *in vitro* oocyte maturation and fertilization affects embryo quality in sheep and deer. *Anim. Reprod. Sci.*, **213**: 106279. <https://doi.org/10.1016/j.anireprosci.2020.106279>
- Schwartz, J., Schneider, M.R., Rodrigues, J.L. and Reichenbach, H.D., 1998. Effect of short-term storage of bovine oocytes in different media and temperatures on the subsequent *in vitro* embryo development. *Theriogenology*, **1**: 217. [https://doi.org/10.1016/S0093-691X\(98\)90570-2](https://doi.org/10.1016/S0093-691X(98)90570-2)
- Senosy, W., Kassab, A.Y. and Mohammed, A.A., 2017. Effects of feeding green microalgae on ovarian activity, reproductive hormones and metabolic parameters of Boer goats in arid subtropics. *Theriogenology*, **96**: 16–22. <https://doi.org/10.1016/j.theriogenology.2017.03.019>
- Senosy, W., Kassab, A.Y., Hamdon, H.A. and Mohammed, A.A., 2018. Influence of organic phosphorus on reproductive performance and metabolic profiles of anoestrous Farafra ewes in subtropics at the end of breeding season. *Reprod. Domest. Anim.*, **53**: 904–913. <https://doi.org/10.1111/rda.13183>
- Shabankareh, H.K., Azimi, G. and Torki, M., 2014. Developmental competence of bovine oocytes selected based on follicle size and using the brilliant cresyl blue (BCB) test. *Iran. J. Reprod. Med.*, **12**: 771.
- Sinclair, K.D., Lunn, L.A., Kwong, W.Y., Wonnacott,

- K., Linforth, R.S.T. and Craigon, J., 2008. Amino acid and fatty acid composition of follicular fluid as predictors of *in-vitro* embryo development. *Reprod. Biomed. Online*, **16**: 859–868. [https://doi.org/10.1016/S1472-6483\(10\)60153-8](https://doi.org/10.1016/S1472-6483(10)60153-8)
- Singh, P., Singh, P.K. and Singh, R.L., 2022. Transgenic animals. In: *Emerging issues in climate smart livestock production* (eds. S. Mondal, R.L. Singh), Academic Press, pp. 171–208. <https://doi.org/10.1016/B978-0-12-822265-2.00001-6>
- Somfai, T., Inaba, Y., Watanabe, S., Geshi, M. and Nagai, T., 2012. Follicular fluid supplementation during *in vitro* maturation promotes sperm penetration in bovine oocytes by enhancing cumulus expansion and increasing mitochondrial activity in oocytes. *Reprod. Fertil. Dev.*, **24**: 743–752. <https://doi.org/10.1071/RD11251>
- Spacek, S.G. and Carnevale, E.M., 2018. Impact of equine and bovine oocyte maturation in follicular fluid from young and old mares on embryo production *in vitro*. *J. Equine Vet. Sci.*, **68**: 94–100. <https://doi.org/10.1016/j.jevs.2018.04.009>
- Suresh, A., Shukla, M.K., Kumar, D., Shrivastava, O.P. and Verma, N., 2021. Simulated physiological oocytematuration (SPOM) improves developmental competence of *in vitro* produced goat embryos. *Theriogenology*, **172**: 193–199. <https://doi.org/10.1016/j.theriogenology.2021.06.003>
- Taira, A.R., Brandão, F.Z., Brair, V.L., Cosentino, I.O., Leal, F.S.C., Ribeiro, A.C.S., Balaro, M.F.A., Batista, R.I.T.P., Souza-Fabjan, J.M.G. and da Fonseca, J.F., 2022. Biostimulation with the ram effect increases the follicle recruitment, ovulatory diameter, and embryo viability rate in super ovulated ewes. *Theriogenology*, **181**: 140–146. <https://doi.org/10.1016/j.theriogenology.2022.01.021>
- Tang, K., Yang, W.C., Li, X., Wu, C.J., Sang, L. and Yang, L.G., 2012. GDF-9 and bFGF enhance the effect of FSH on the survival, activation, and growth of cattle primordial follicles. *Anim. Reprod. Sci.*, **131**: 129–134. <https://doi.org/10.1016/j.anireprosci.2012.03.009>
- Tian, H., Qi, Q., Yan, F., Wang, C., Hou, F., Ren, W., Zhang, L. and Hou, J., 2021. Enhancing the developmental competence of prepubertal lamb oocytes by supplementing the *in vitro* maturation medium with sericin and the fibroblast growth factor 2-leukemia inhibitory factor-insulin-like growth factor 1 combination. *Theriogenology*, **159**: 13–19. <https://doi.org/10.1016/j.theriogenology.2020.10.019>
- Virant-Klun, I., Bauer, C., Ståhlberg, A., Kubista, M. and Skutella, T., 2018. Human oocyte maturation *in vitro* is improved by co-culture with cumulus cells from mature oocytes. *Reprod. BioMed. Online*, **36**: 508–523. <https://doi.org/10.1016/j.rbmo.2018.01.011>
- von Wolff, M., Eisenhut, M., Stute, P. and Bersinger, N.A., 2022. Gonadotrophin stimulation reduces follicular fluid hormone concentrations and disrupts their quantitative association with cumulus cell mRNA. *Reprod. Biomed. Online*, **44**: 193–199. <https://doi.org/10.1016/j.rbmo.2021.08.018>
- Wang, Y., Xing, C.H., Chen, S. and Sun, S.C., 2022. Zearalenone exposure impairs organelle function during porcine oocyte meiotic maturation. *Theriogenology*, **177**: 22–28. <https://doi.org/10.1016/j.theriogenology.2021.10.008>
- Wani, N.A. and Hong, S.B., 2018. Source, treatment and type of nuclear donor cells influences *in vitro* and *in vivo* development of embryos cloned by somatic cell nuclear transfer in camel (*Camelus dromedarius*). *Theriogenology*, **106**: 186–191. <https://doi.org/10.1016/j.theriogenology.2017.10.010>
- Wani, N.A., Hong, S. and Vettical, B.S., 2018. Cytoplasm source influences development of somatic cell nuclear transfer (SCNT) embryos *in vitro* but not their development to term after transfer to synchronized recipients in dromedary camels (*Camelus dromedarius*). *Theriogenology*, **118**: 137–143. <https://doi.org/10.1016/j.theriogenology.2018.05.021>
- Wani, N.A. and Hong, S.B., 2020. Effect of roscovitine pretreatment on *in vitro* maturation of oocytes and their subsequent developmental after chemical activation in dromedary camel (*Camelus dromedarius*). *Theriogenology*, **157**: 176–180. <https://doi.org/10.1016/j.theriogenology.2020.07.025>
- Wei, K.N., Wang, X.J., Zeng, Z.C., Gu, R.T., Deng, S.Z., Jiang, J., Xu, C.L., Li, W. and Wang, H.L., 2021. Perfluorooctane sulfonate affects mouse oocyte maturation *in vitro* by promoting oxidative stress and apoptosis induced by mitochondrial dysfunction. *Ecotoxicol. Environ. Saf.*, **225**: 112807. <https://doi.org/10.1016/j.ecoenv.2021.112807>
- Wilmut, I., Schnieke, A.E., McWhir, J., Kind, A.J. and Campbell, K.H.S., 1997. Viable offspring derived from fetal and adult mammalian cells. *Nature*, **385**: 810–813. <https://doi.org/10.1038/385810a0>
- Wu, Z., Liu, K. and Zhang, C., 2021. Clinical applications of *in vitro* primordial follicle activation. *Curr. Opin. Endocr. Metab. Res.*, **18**: 194–200. <https://doi.org/10.1016/j.coemr.2021.03.020>

- Yoshimura, Y., Nakamura, Y., Yamada, H., Nanno, T., Ubukata, Y., Ando, M. and Suzuki, M., 1991. Gonadotropin-releasing hormone agonists induce meiotic maturation and degeneration of oocytes in the *in vitro* perfused rabbit ovary. *Fertil. Steril.*, **55**: 177–183. [https://doi.org/10.1016/S0015-0282\(16\)54079-5](https://doi.org/10.1016/S0015-0282(16)54079-5)
- Zabihi, A., Shabankareh, H.K., Hajarian, H. and Foroutanifar, S., 2021. *In vitro* maturation medium supplementation with resveratrol improves cumulus cell expansion and developmental competence of Sanjabi sheep oocytes. *Livest. Sci.*, **243**: 104378. <https://doi.org/10.1016/j.livsci.2020.104378>
- Zhang, H., Risal, S., Gorre, N., Busayavalasa, K., Li, X., Shen, Y., Bosbach, B., Brännström, M. and Liu, K., 2014. Somatic cells initiate primordial follicle activation and govern the development of dormant oocytes in mice. *Curr. Biol.*, **24**: 2501–2508. <https://doi.org/10.1016/j.cub.2014.09.023>
- Zhang, J. Hui Liu, H., 2015. Cytoplasm replacement following germinal vesicle transfer restores meiotic maturation and spindle assembly in meiotically arrested oocytes. *Reprod. BioMed. Online*, **31**: 71–78. <https://doi.org/10.1016/j.rbmo.2015.03.012>
- Zhang, T., Fan, X., Li, R., Zhang, C. and Zhang, J., 2018. Effects of pre-incubation with C-type natriuretic peptide on nuclear maturation, mitochondrial behavior, and developmental competence of sheep oocytes. *Biochem. biophys. Res. Commun.*, **497**: 200–206. <https://doi.org/10.1016/j.bbrc.2018.02.054>
- Zhao, X., Du, F., Liu, X., Ruan, Q., Wu, Z., Lei, C., Deng, Y., Luo, C., Jiang, J. and Shi, D., 2019. Brain-derived neurotrophic factor (BDNF) is expressed in buffalo (*Bubalus bubalis*) ovarian follicles and promotes oocyte maturation and early embryonic development. *Theriogenology*, **130**: 79–88. <https://doi.org/10.1016/j.theriogenology.2019.02.020>
- Zhu, J., Lian, Y., Zhuang, X., Lin, S., Zheng, X., Li, J., Rong, L. and Ping, L.P., 2022. Poor morphology of inner cell mass increases birth weight and large for gestational age. *Reprod. BioMed. Online*, **45**: 890–897. <https://doi.org/10.1016/j.rbmo.2022.06.025>