



Molecular Depiction of Vermicompost Associated Bacteria Possessed Agricultural Traits

Saiqa Andleeb^{1*}, Anum Naseer¹, Fahad Kiyani¹, Fayaz Ahmed², Muhammad Fiaz Khan², Wajid Arshad Abbasi³ and Shaukat Ali⁴

¹Microbial Biotechnology and Vermi-technology Laboratory, Vermi-tech Unit, Department of Zoology, University of Azad Jammu and Kashmir, Muzaffarabad, 13100, Pakistan

²Department of Zoology, Hazara University, Mansehra, KPK, Pakistan

³Department of CS&IT, Computational Biology and Data Analysis Laboratory, University of Azad Jammu and Kashmir, Muzaffarabad, Pakistan

⁴Department of Zoology, GC University, Lahore, Pakistan

Fahad Kiyani and Fayaz Ahmed made equal contribution.

ABSTRACT

Vermicomposting is an environmentally cordial process in which different organic wastes are converted into compost by earthworms. Vermicompost is the end product of vermicomposting which is a peat-like material, highly porous with maximum aeration and drainage capacity. It is rich in essential nutrients and beneficial microbes which are very useful for plant development and growth. The current research was conducted to isolate and identify some beneficial bacteria from the vermicompost which could be used as plant growth-promoting bacteria (PGPB) in the future to enhance crop production in Pakistan. Cattle dung, organic waste materials (egg shells, coconut peels, paper waste, raw vegetables, and fruits), and *Eisenia fetida* were used for vermicompost production. Vermicompost-associated bacteria were identified using microscopic techniques, various biochemical tests, cultural media, and ribotyping. Plant growth-promoting traits i.e. Indole acetic acid, siderophore, ammonia, hydrogen cyanide, and potassium hydroxide were also performed. Antibioassay analysis was also done to screen the multi-drug-resistant bacteria. Results revealed that the isolated vermicompost-associated bacteria were identified as *Achromobacter ruhlandii*, *Bacillus wiedmannii*, *Pseudomonas* sp., *Achromobacter xylosoxidans*, *Oceanobacillus oncorhynchi*, *Bacillus mycoides*, *Serratia nematodiphila*, *Serratia marcescens*, and *Paenibacillus dendritiformis* possessed plant growth promoting traits, non-pathogenic, and non-multi-drug resistant bacteria. It was concluded that vermicompost-associated bacteria could be used as a potential source of biofertilizers not only to enhance plant growth and development, and soil fertility, but also to be useful for the control of pests and pathogens for sustainable agriculture in Pakistan.

Article Information

Received 25 May 2023

Revised 25 June 2024

Accepted 07 July 2024

Available online 07 October 2024 (early access)

Published 25 September 2025

Authors' Contribution

SA conceived the idea, designed the experiments, analysed and interpreted the results, and wrote manuscript. AN helped in the designing of experiments. FK and FA performed the experiments and wrote the first draft. WAB contributed in molecular depiction using bioinformatics tools. MFK and SA did formal analysis, interpreted results and manuscript review.

Key words

Vermicompost, Vermibacteria, Biochemical tests, Ribotyping, Azad Kashmir, *E. fetida*

INTRODUCTION

Soil microorganisms i.e., bacteria, fungi, and Actinomycetes involved in the soil formation, improving

the soil fertility via degradation and detoxification of organic materials in soil, and maintain the ecological balance (Emperor *et al.*, 2015; Zhu *et al.*, 2017). Vermicomposting is an ecofriendly, bio-oxidative, and non-thermophilic organic decomposition technology to recycle the organic waste into fine granular product called vermicompost via the joint action of earthworms and microorganisms (Lazcano *et al.*, 2008; Chitrapriya *et al.*, 2013) (Supplementary Fig. 1). Various earthworm species

* Corresponding author: drsaiqa@gmail.com, drsaiqa@ajku.edu.pk
0030-9923/2025/0006-2625 \$ 9.00/00



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abbreviations

PGPB, plant growth-promoting bacteria; VAB, vermicompost associated bacteria; TMP, Trimethoprim; ATM, Aztreonam; K, Kanamycin, E, Erythromycin, S, Sulfonamide, TOB, Tobramycin; P, Penicillin; NOR, Norfloxacin, AMC, Amoxicillin and AMP, Ampicillin.

i.e., *Metaphire californica*, *Eudrilus eugeniae*, *Eisenia andrei*, *Perionyx excavates*, and *Eisenia fetida* were used for vermicomposting (Deka *et al.*, 2011; Varma *et al.*, 2016; Boruah *et al.*, 2019; González-Moreno *et al.*, 2022).

Earthworms ingest soil microbes, their population increases *via* passage through the intestinal track due to favorable environment, and excrete them along with nutrients in their vermicast (Huang *et al.*, 2013; Pathma and Sakthivel, 2012). Several antibiotics and enzymes play an important role in the decomposition and degradation of organic molecules which were released by microbes in the gastrointestinal track of earthworms (Lazcano *et al.*, 2008).

Vermicompost is not only rich in micro- and macronutrients, enzymes, plant growth hormones, plant growth regulators, and antibiotics (Ravindran *et al.*, 2016; Shafique *et al.*, 2021) but also improves the plant growth, plant nutrition and productivity, soil fertility, soil structure as well as growth of soil beneficial microbes (Pathma and Sakthivel, 2012; Zhu *et al.*, 2017) (Supplementary Fig. 1). In addition, vermicompost improves soil aeration, moisture contents, pH, electrical conductivity, and water holding capacity (Shafique *et al.*, 2021). Vermicompost suppresses plant diseases and increased microbial population like phosphate solubilizers, nitrogen fixers, Indole acetic acid producers, siderophore producers, which could significantly enhance the vegetative plant growth of ornamental plants (Shafique *et al.*, 2021; Chauhan and Singh, 2015). Previous literature reported the presence of bacteria in vermicompost produced *via* *L. mauritii*, *P. excavatus* and *E. eugeniae* (Upadhyay *et al.*, 2015; Selvi and Koilraj, 2015; Begum and Bora, 2018). Devi *et al.* (2009) and Pathma and Sakthivel (2012) illustrated the presence of Proteobacteria, Bacteroidetes Planctomycetes, Actinobacteria, and Firmicutes in vermicompost (Yasir *et al.*, 2009a).

In view of above the objectives of the current research were to isolate, characterize, and identify vermicompost associated bacteria from vermicompost produced *via* *E. fetida*, to explore the agricultural traits and to screen the prevalence of antibiotic resistance among vermicompost associated bacteria. No research has been done before on the beneficial vermi-bacterium isolation and identification from vermicompost in Pakistan. Hopefully, this identified bacterial isolates could be used as microbial biofertilizer to enhance the seed germination and plant growth of various crops in future.

MATERIALS AND METHODS

Chemicals used

Nutrient broth (lennox), Luria Bertani (LB) broth (lennox), nutrient agar medium (sigma-aldrich), hydrogen

peroxide, oxidase reagent, gram staining kit (merck), skim milk agar (neogen), Wattman No.1 disc, MacConkey agar (sigma-aldrich), mannitol salt agar (oxoid), Nessler's reagent (sigma-aldrich), Kovac's reagent, phenol, 0.5% Picric acid (sigma-aldrich), King's B medium (Sigma), 2% sodium carbonate (merck), dilute Iodine, 3% KOH starch (sigma-aldrich) and bacteriological Peptone (oxoid). The antibiotics used were trimethoprim (TMP 5µg), aztreonam (ATM 30µg), kanamycin (K 30µg), erythromycin (E 15µg), sulfonamide (S 300µg), tobramycin (TOB 10µg), penicillin (P 10µg), norfloxacin (NOR 10µg), amoxicillin (AMC 30µg) and ampicillin (AMP 25µg). All these antibiotics were taken from Oxide company.

Glassware and types of equipment used

Steam sterilizer (autoclave), Laminar flow (ESO Prod Model; EQU/03-EHC; Serial # 2000-0052), 37°C incubator (MMM group Medcenter Enrich tungsten GmbH), Analytical balance (SARTORIUS GMBM GOTTINGEN, Germany), sterile distilled water, sterile bottles, sterile dissecting pins, dissecting box, dissecting board, gloves, 37°C shaker (Irmeco GmbH, Germany), digital weighing machine (Jeweler Precision Balance Model: DH-V600A), 70% Ethanol, 500 ml beakers, 250 ml conical flasks, test tubes, petri plates, L shaped glass rod, microscope, bacteriological wire loop, micropipette, glass slides, glycerol, coverslips, spirit lamp and toothpicks.

Isolation, enumeration, and purification of vermi-bacteria

Vermicompost associated bacteria (VAB) were isolated from the vermicompost collected from Vermitech Unit, Department of Zoology, The University of Azad Jammu and Kashmir, Muzaffarabad. Vermicompost was prepared using cow dung, raw vegetables and fruits, coconut peel, wheat straw and rice straw, egg shells and waste papers. *Eisenia fetida* was used for the vermicomposting (Shafique *et al.*, 2021). Hundred mg vermicompost was added in 200 ml of distilled water, mixture was shaken for 20 min, and kept at room temperature for two days. After incubation, serial dilution method was used as illustrated by Somasegaran and Hoben (2012) for the isolation of vermicompost associated bacteria. Nutrient broth medium was used for bacterial culturing. After incubation, serial dilution 10^{-2} , 10^{-3} , 10^{-4} , and 10^{-5} were made and again incubated for 24 h at room temperature to pick the single colony of VAB. After incubation, 10 µl sample was spread on nutrient agar medium and incubated at 37°C for overnight. Next day, several colonies were observed in the case of 10^{-2} , 10^{-3} , 10^{-4} , and 10^{-5} diluted samples. From these plates, twenty bacterial isolates were taken in nutrient broth medium and incubated for 24 h at 37°C. After incubation,

14 VAB isolates (VAB-1 to VAB-14) were purified after sub-culturing and then kept at -20 °C in the form of 60% glycerol.

Morphological and biochemical characteristics of VAB

For the identification of VAB different techniques and methods i.e. microscopic (Gram staining), cultural techniques (nutrient agar medium, MacConkey agar), biochemical tests (Plant growth promoting traits like production of Indole acetic acid, siderophore, ammonia, hydrogen cyanide, potassium hydroxide), hydrolytic enzymes production like catalase, oxidase, amylase, protease, and lipase, and hemolytic test were applied (Okon *et al.*, 1977; Dawwam *et al.*, 2013; Kumar *et al.*, 2015). The different colors, size, elevations, textures, shapes, and edges of vermicompost associated bacteria were recorded.

Molecular characterization

Genomic DNA was extracted from bacterial isolates using chloroform: Isoamyl alcohol method with slight modifications. The isolates were grown in Luria broth medium for 24 h at 37 °C. After incubation, medium was centrifuged at 10,000 rpm for 5 min to harvest cells in pellet form. The pellet was suspended in lysis buffer-1 (Tris EDTA and SDS; pH 4.0) and centrifuged at 10,000 rpm for 10 min. After centrifugation, 500 µL of chloroform: isoamyl alcohol (24:1) was added, mixed, and centrifuged at 10,000 rpm for 10 min. To the collected supernatant 1/10th volume of sodium acetate and 2.5 volumes of chilled 100% absolute ethanol were added, incubated at -20 °C for overnight, Next day, samples were centrifuged for 10 min at 10,000 rpm, and pellet was washed with 70% ethanol. After centrifugation, pellet was dried for 3 h at room temperature (25±2 °C) and DNA was suspended in distilled water (20 µL). For the identification of bacterial diversity, 16S rRNA primers; universal bacterial primer (27F; 5'-AGAGTTTGTATCCTGGCTCAG-3') was taken to amplify 1500 bps sequence using following PCR conditions (initial denaturation 95 °C for 10 min; cyclic denaturation at 95 °C for 30 sec; annealing at 50°C for 1 min; cyclic extension 72°C for 1 min 30 sec; and final extension 72°C for 7 min; 30 cycles) (de Lillo *et al.*, 2006). After PCR analysis all PCR products were dispatch to Macrogen, Korea for sequence analysis. Obtained nucleotide sequences were further preceded for homology through BLAST at National Center for Biotechnology Information (NCBI) platform. The evolutionary history was concluded through Tamura-Nei model and neighbour-joining method (Tamura *et al.*, 2007). This analysis contained 32 nucleotide sequences. There were 1573 positions in the final dataset. Evolutionary studies were accompanied in MEGA X (Kumar *et al.*, 2018). After

homology prediction and phylogenetic analysis, amplified sequences were submitted to Genbank, NCBI for obtaining accession numbers.

Antibiogram analysis

The emergence of multi-drug resistant among microbial biofertilizers is serious concern in developing countries now a days due to anthropogenic activities. Antibiotic resistant biofertilizers carry antibiotic resistant genes (ARGs) which aggravate negative impact not only on environment as well as on wildlife and public health (Mahdi *et al.*, 2022). So, in the current study, various antibiotics like kanamycin (30µg), norfloxacin (10µg), erythromycin (15µg), amoxicillin (30µg), ampicillin (25µg), trimethoprim (5µg), penicillin (10µg), tobramycin (10µg), sulfonamide (300µg) and aztreonam (30µg) were used to display the sensitivity/resistivity of vermibacteria using agar disc diffusion method (Bauer *et al.*, 1959; Brown and Kothari, 1975). Nutrient agar (Oxoid: CMOO3) and nutrient broth media (Oxoid: CM1) were used for bacterial growth. The bacteria were added to a nutrient broth medium for the growth and incubated for 24 h on a rotary shaker at 37°C. The incubated culture was mixed in a freshly prepared nutrient agar medium (NAM) at 45°C. The mixture was poured into sterilized Petri dishes, solidified in laminar flow at room temperature. After solidification, antibiotic discs were placed on the surface and then incubated for 24 h at 37°C. According to Seeley *et al.* (2001), the growth of bacteria was determined in 24-48 h, and the diameter of the inhibition zone in mm was also measured with the help of ruler (Hammer *et al.*, 1999).

Statistical analysis

Each experiment was repeated in triplicates and Mean ± Standard Deviation from absolute data was calculated: (<http://easycalculation.com/statistics/standard-deviation.php>).

RESULTS

Morphological features

The isolated bacterial isolates were characterized by morphological features, microscopic studies, and biochemical characterizations (Table I). Results revealed that different colonies were appeared on the various agar culture media plates i.e., MacConkey agar and nutrient agar medium. The colonies of vermicompost associated bacteria were found with different colors, size, elevations, texture, shape, and edge, respectively. The colonies of VAB showed yellow, opaque, white, cream, and cherry in color; the surface and shape of colonies were recorded as spherical, rough, oval, smooth, circular, and oval. On the

other hand, colony elevations (convex, flat, umbonate), margins (entire, irregular, filiform), and wet texture were recorded. Gram staining indicated that VAB-1, VAB-3, VAB-4, VAB-10, and VAB-12 were observed as Gram-negative rod bacteria while VAB-2, VAB-5, VAB-6, VAB-7, VAB-8, VAB-9, VAB-11, and VAB-14 are rod shaped Gram-positive bacteria.

Biochemical characterization

Results revealed that all vermicompost associated bacteria were shown catalase and oxidase positive results. In case of proteolytic assay, VAB-1, VAB-4, VAB-5, VAB-6, VAB-8, VAB-9, VAB-10, VAB-11, VAB-12, and VAB-14 were shown positive hydrolysis of casein in the range of 2.0 ± 0.0 mm to 8.0 ± 0.0 mm, while VAB-2, VAB-3, VAB-7 showed negative results. Similarly, positive lipolytic activity was shown by all tested VAB isolates. All VAB isolates showed amylolytic activity except VAB-3. VAB-1, VAB-3, VAB-4, and VAB-9 showed urease and citrate positive results (Table I). All VAB showed the mannitol fermentation when streaked on the mannitol salt agar medium except VAB-1, VAB-2, and VAB-3 (Table I). Hemolytic test results revealed that all isolated VABs are non-pathogenic in nature.

Plant growth promoting traits

Results revealed that all VAB isolates were involved

in the production of siderophore and Indole acetic acid (except VAB-6), and act as phosphate solubilizers (Table II). Ammonia production test revealed that VAB-1, VAB-6, VAB-7, VAB-8, VAB-9, VAB-10, VAB-12, and VAB-14 were positive for this test. Yellow color was observed in the medium, denoting the production of ammonia by these isolates. All VAB involved in the HCN production except VAB-6 (Table II). In case of KOH test, VAB-3, VAB-4, and VAB-6 showed negative results.

Molecular identification

The 16S rDNA genes from isolated bacteria were amplified using 27F bacterial universal primer and the size of amplified genes was recorded in the range of 1100 bps to 1769 bps. The partial nucleotide sequences were proceeded for further BLAST analysis at National Center for Biotechnology Information (NCBI) and results reveal that no significant similarities of VAB-8, VAB-9, and VAB-11 were found (Table III). On the other hand, VAB-1 indicated 91% similarity with *Achromobacter insolitus* (AB642187.1), *Achromobacter denitrificans* (KY475779.1), *Achromobacter anxifer* (MN197585.1), *Achromobacter ruhlandii* (KY087962.1), *Achromobacter agilis* (MK791141.1), and *Achromobacter xylosoxidans* (MK587674.1); VAB-2 showed 99.55% homology with *Bacillus cereus* (JX971533.1) and 99.46% with

Table I. Morphological and biochemical characterization of vermicompost associated bacteria.

Characteristics → Bacterial strains ID↓	GS	Sha	Color	Colony sha/sur	Ele	Mar	Tex	MSA	Mot	Cata	Oxid	Amy	Pro	Lip	Ure	Cit
<i>Achromobacter ruhlandii</i> (VAB-1)	-	Rod	Yel	Sph/Rou	Con	Ent	Wet	-	+	+	+	+	+	+	+	+
<i>Bacillus wiedmannii</i> (VAB-2)	+	Rod	Opa/W	Oval/Sm	Flat	Irg	Wet	-	+	+	+	+	-	+	-	+
<i>Pseudomonas</i> sp. (VAB-3)	-	Rod	Yel	Sph/Rou	Flat	Ser	Dry	-	+	+	+	-	-	+	+	+
<i>Achromobacter xylosoxidans</i> (VAB-4)	-	Rod	Opa	Oval/Sm	Con	Ent	Wet	+	+	+	+	+	+	+	+	+
<i>Oceanobacillus oncorhynchi</i> (VAB-5)	+	Rod	Yel	Cir/Sm	Con	Ent	Wet	+	+	+	+	+	+	+	-	-
<i>Bacillus mycoides</i> (VAB-6)	+	Rod	W	Cir/Sm	Flat	Irg	Wet	+	-	+	+	+	+	+	-	+
<i>Serratia nematodiphila</i> (VAB-7)	+	Rod	Cre	Cir/Sm	Con	Ent	Wet	+	+	+	+	+	-	+	-	+
Unknown (VAB-8)	+	Rod	Yel	Oval/Sm	Con	Ent	Wet	+	+	+	+	+	+	+	-	+
Unknown (VAB-9)	+	Rod	Yel	Oval/Sm	Con	Ent	Wet	+	+	+	+	+	+	+	+	+
<i>Serratia marcescens</i> (VAB-10)	-	Rod	Cre	Cir/Sm	Umb	Filli	Wet	+	+	+	+	+	+	+	-	+
Unknown (VAB-11)	+	Rod	Cre	Cir/Rou	Con	Ent	Wet	+	+	+	+	+	+	+	+	+
<i>Serratia marcescens</i> (VAB-12)	-	Rod	Cre	Cir/Sm	Umb	Filli	Wet	+	+	+	+	+	+	+	-	+
<i>Paenibacillus dendritiformis</i> (VAB-14)	+	Rod	Che	Oval/Rou	Con	Ent	Wet	+	+	+	+	+	+	+	-	+

GS, Gram staining; Sha, Shape of bacteria; Sur, Surface; Ele, Elevation; Mar, Margin; Tex, Texture; MSA, Mannitol salt agar; Mot, Motility; Cata, Catalase; Oxid, Oxidase; Amy, Amylase; Pro, Proteases; Lip, Lipase; Ure, Urease; Cit, Citrate; Yel, Yellow; W, White; Opa, Opaque; Cre, Cream; Sm, Smooth; Umb, Umbonate; Filli, Filiform; Con, Convex; Ent, Entire; W, White; Irg, Irregular; Cir, Circular; Rou, Rough; Sph, Spherical; +, positive; -, negative.

Table II. Plant growth promoting traits of vermicompost associated bacteria.

Agricultural traits → Bacterial strains ID↓	Sidero- phore	Phos- phate	Potassium hydroxide	Indole acetic acid	Hydrogen cyanide	Ammono- nia
<i>Achromobacter ruhlandii</i> (VAB-1)	+	+	+	+	+	+
<i>Bacillus wiedmannii</i> (VAB-2)	+	+	+	+	+	-
<i>Pseudomonas</i> sp. (VAB-3)	+	+	-	+	+	-
<i>Achromobacter xylosoxidans</i> (VAB-4)	+	+	-	+	+	-
<i>Oceanobacillus oncorhynchi</i> (VAB-5)	+	+	+	+	+	-
<i>Bacillus mycoides</i> (VAB-6)	+	+	-	-	-	+
<i>Serratia nematodiphila</i> (VAB-7)	+	+	+	+	+	+
Unknown (VAB-8)	+	+	+	+	+	+
Unknown (VAB-9)	+	+	+	+	+	+
<i>Serratia marcescens</i> (VAB-10)	+	+	+	+	+	+
Unknown (VAB-11)	+	+	+	+	+	-
<i>Serratia marcescens</i> (VAB-12)	+	+	+	+	+	+
<i>Paenibacillus dendritiformis</i> (VAB-14)	+	+	+	+	+	+

Table III. BLAST outcomes of vermicompost associated bacteria.

Microbe coding	Amplified length	Percent identity (%)	Scientific name	Sequence id	Accession length
VAB-1	1170	91%	<i>Achromobacter insolitus</i>	AB642187.1	1295
		91%	<i>Achromobacter denitrificans</i>	KY475779.1	1339
		91%	<i>Achromobacter anaxifer</i>	MN197585.1	1420
		91%	<i>Achromobacter ruhlandii</i>	KY087962.1	1448
		91%	<i>Achromobacter agilis</i>	MK791141.1	1398
		91%	<i>Achromobacter xylosoxidans</i>	MK587674.1	1426
VAB-2	1171	99.55	<i>Bacillus cereus</i>	JX971533.1	1374
		99.46	<i>Bacillus subtilis</i>	MG593998.1	1188
		99.46	<i>Bacillus tropicus</i>	ON878141.1	1351
		99.46	<i>Bacillus thuringiensis</i>	ON974260.1	1423
		99.46	<i>Bacillus paranthracis</i>	ON939702.1	1419
		99.46	<i>Bacillus pacificus</i>	ON885910.1	1417
		99.46	<i>Bacillus wiedmannii</i>	ON340605.1	1370
		99.46	<i>Bacillus paramycoides</i>	ON000574.1	1448
		99.46	<i>Bacillus firmus</i>	LC385949.1	710
VAB-3	1679	89	<i>Escherichia coli</i>	MH793567.1	1371
		93	<i>Stenotrophomonas maltophilia</i>	MK493713.1	643
		93	<i>Pseudomonas</i> sp.	MT793102.1	1498
		93	<i>Pseudomonas</i> sp.	MT793102.1	1498
VAB-4	1224	96.82	<i>Achromobacter insolitus</i>	MN044760.1	1300
		96.52	<i>Achromobacter denitrificans</i>	MT275805.1	1374
		96.52	<i>Achromobacter ruhlandii</i>	KY087962.1	1448
		96.56	<i>Achromobacter xylosoxidans</i>	MK587674.1	1426
		96.09	<i>Achromobacter aegrifaciens</i>	NR_117707.1	1483

Table continues on next page.....

Microbe coding	Amplified length	Percent identity (%)	Scientific name	Sequence id	Accession length
VAB-5	1100	78.95	<i>Bacillus pumilus</i>	KM222185.1	1448
		78	<i>Bacillus subtilis</i>	OM760707.1	1454
		83	<i>Bacillus cereus</i>	FJ404785.1	539
		77	<i>Oceanobacillus oncorhynchi</i>	JX280491.1	1485
		75	<i>Pediococcus pentosaceus</i>	ON545700.1	1102
VAB-6	1100	81	<i>Bacillus anthracis</i>	KP813652.1	1221
		81	<i>Bacillus toyonensis</i>	LN995802.1	1111
		81	<i>Bacillus thuringiensis</i>	HM032789.1	948
		81	<i>Bacillus cereus</i>	MF726967.1	1267
		80	<i>Bacillus proteolyticus</i>	MK418822.1	1028
		80	<i>Bacillus albus</i>	MN658863.1	1235
		81	<i>Bacillus manliponensis</i>	MW519873.1	1083
		81	<i>Bacillus pseudomyoides</i>	MT573519.1	1040
		82	<i>Bacillus mycoides</i>	KU687331.1	1257
VAB-7	1100	87.03	<i>Serratia marcescens</i>	MF462933.1	1479
		87	<i>Staphylococcus aureus</i>	MH603394.1	1161
		87	<i>Serratia surfactantfaciens</i>	NR_169468.1	1495
		87	<i>Enterobacter cloacae</i>	KJ850207.1	1434
		87	<i>Raoultella planticola</i>	MZ203710.1	1363
		87	<i>Serratia nematodiphila</i>	KM099143.1	1392
		87	<i>Klebsiella</i> sp.	MZ203699.1	1363
VAB-8	1100	No significant similarity found			
VAB-9	1100	No significant similarity found			
VAB-10	1100	83.16	<i>Serratia marcescens</i>	MT386168.1	1385
		83.16	<i>Pseudomonas</i> sp. strain jx-18	KY780232.1	1454
		83.16	<i>Serratia nematodiphila</i>	ON651725.1	1261
VAB-11		No significant similarity found			
VAB-12	1591	94	<i>Serratia marcescens</i>	KM099142.1	1411
		92	<i>Serratia nematodiphila</i>	MN691576.1	1251
VAB-14	1100	89	<i>Paenibacillus dendritiformis</i>	MG592704.1	1414
		89	<i>Paenibacillus popilliae</i>	KC107790.1	1380
		89	<i>Paenibacillus thiaminolyticus</i>	EU330645.1	1474

Bacillus subtilis (MG593998.1), *Bacillus tropicus* (ON878141.1), *Bacillus thuringiensis* (ON974260.1), *Bacillus paranthracis* (ON939702.1), *Bacillus pacificus* (ON885910.1), *Bacillus wiedmannii* (ON340605.1), *Bacillus paramycoides* (ON000574.1); VAB-3 showed 94% resemblance with *Bacillus firmus* (LC385949.1), 89% with *Escherichia coli* (MH793567.1), 93% with *Stenotrophomonas maltophilia* (MK493713.1), 93% with *Pseudomonas* sp. (MT793102.1); VAB-4 showed 96.82% similarity with *Achromobacter insolitus* (MN044760.1), 96.52% with *Achromobacter denitrificans* (MT275805.1),

96.52% with *Achromobacter ruhlandii* (KY087962.1), 96.56% with *Achromobacter xylosoxidans* (MK587674.1), 96.09% with *Achromobacter aegrifaciens* (NR_117707.1); VAB-5 indicated 78.95% similarity with *Bacillus pumilus* (KM222185.1), 78% with *Bacillus subtilis* (OM760707.1), 83% with *Bacillus cereus* (FJ404785.1), 77% with *Oceanobacillus oncorhynchi* (JX280491.1), 75% with *Pediococcus pentosaceus* (ON545700.1); VAB-6 showed 81% homology with *Bacillus anthracis* (KP813652.1), *Bacillus toyonensis* (LN995802.1), *Bacillus thuringiensis* (HM032789.1), *Bacillus cereus* (MF726967.1), *Bacillus*

manliponensis (MW519873.1), *Bacillus pseudomycoloides* (MT573519.1), 80% with *Bacillus proteolyticus* (MK418822.1), *Bacillus albus* (MN658863.1), 82% with *Bacillus mycoloides* (KU687331.1); VAB-7 displayed 87.03% resemblance with *Serratia marcescens* (MF462933.1), 87% with *Staphylococcus aureus* (MH603394.1), *Serratia surfactantifaciens* (NR_169468.1), *Enterobacter cloacae* (KJ850207.1), *Raoultella planticola* (MZ203710.1), *Serratia nematodiphila* (KM099143.1), *Klebsiella* sp. (MZ203699.1); VAB-10 indicated 83.16% similarity with *Serratia marcescens* (MT386168.1), *Pseudomonas* sp. strain jx-18 (KY780232.1), *Serratia nematodiphila* (ON651725.1); VAB-12 showed 94% resemblance with *Serratia marcescens* (KM099142.1) and 92% with *Serratia nematodiphila* (MN691576.1); and VAB-14 indicated 89% homology with *Paenibacillus dendritiformis* (MG592704.1), *Paenibacillus popilliae* (KC107790.1), and *Paenibacillus thiaminolyticus* (EU330645.1). The results of phylogenetic tree using the NJ method were recorded as VAB-1 indicated relationship with *Achromobacter ruhlandii* (81%), VAB-2 with *Bacillus wiedmannii* (81%), VAB-3 with *Pseudomonas* sp. (82%); VAB-4 with *Achromobacter xylosoxidans* (96%), VAB-5 with *Oceanobacillus oncorhynchi* (74%), VAB-6 with *Bacillus mycoloides* (87%), VAB-7 with *Serratia nematodiphila* (74%), VAB-10 with *Serratia marcescens* (95%), VAB-12 with *Serratia marcescens* (79%), and

VAB-14 with *Paenibacillus dendritiformis* (75%).

Antibiogram analysis

Erythromycin and norfloxacin showed the maximum inhibition of tested vermicompost associated bacteria (Table IV). Erythromycin showed the maximum inhibition of *Achromobacter ruhlandii* (18.0±0.0 mm), *Pseudomonas* sp. (14.0±0.0 mm), *Oceanobacillus oncorhynchi* (14.0±0.0 mm), *Bacillus mycoloides* (12.0±0.0 mm), unknown (18.0±0.0 mm), unknown (18.0±0.0 mm), *Serratia marcescens* (20.0±0.0 mm), unknown (16.0±0.0 mm), *Serratia marcescens* (20.0±0.0 mm). Similarly, norfloxacin indicated the maximum inhibition of *Achromobacter ruhlandii* (14.0±0.0 mm), *Achromobacter xylosoxidans* (13.0±0.0 mm), *Bacillus mycoloides* (11.0±0.0), Unknown (15.0±0.0 mm), unknown (14.0±0.0 mm), *Serratia marcescens* (13.0±0.0 mm), unknown (17.0±0.0 mm), *Serratia marcescens* (11.0±0.0 mm) *Paenibacillus dendritiformis* (17.0±0.0 mm). Trimethoprim, kanamycin, sulfonamide showed the maximum inhibition of *Oceanobacillus oncorhynchi* and VAB-11 with 17.0±0.0 mm, 15.0±0.0 mm, 11.0±0.0 mm, 19.0±0.0 mm, 14.0±0.0 mm, 12.0±0.0 mm. On the other hand, all tested VAB isolates displayed resistant towards trimethoprim, aztreonam, ampicillin, and amoxocillin (Table IV). Tobramycin and penicillin showed lowest and moderate inhibition of tested vermicompost associated bacterial isolates.

Table IV. Antibigram analysis of vermicompost associated bacteria.

Used Antibiotics → Bacterial strains↓	Zone of inhibition (mm) M±SD									
	TMP (5µg)	ATM (30µg)	K (30µg)	E (15µg)	S (300µg)	TOB (10µg)	P (10µg)	NOR (10µg)	AMC (30µg)	AMP (25µg)
<i>Achromobacter ruhlandii</i> (VAB-1)	0.0±0.0	0.0±0.0	9.0±0.0	18.0±0.0	10.0±0.0	5.0±0.0	5.0±0.0	14.0±0.0	0.0±0.0	0.0±0.0
<i>Bacillus wiedmannii</i> (VAB-2)	0.0±0.0	0.0±0.0	9.0±0.0	6.0±0.0	14.0±0.0	7.0±0.0	0.0±0.0	9.0±0.0	0.0±0.0	0.0±0.0
<i>Pseudomonas</i> sp. (VAB-3)	0.0±0.0	0.0±0.0	7.0±0.0	14.0±0.0	12.0±0.0	6.0±0.0	0.0±0.0	10.0±0.0	0.0±0.0	0.0±0.0
<i>Achromobacter xylosoxidans</i> (VAB-4)	0.0±0.0	0.0±0.0	5.0±0.0	0.0±0.0	0.0±0.0	5.0±0.0	0.0±0.0	13.0±0.0	0.0±0.0	0.0±0.0
<i>Oceanobacillus oncorhynchi</i> (VAB-5)	17.0±0.0	0.0±0.0	15.0±0.0	14.0±0.0	11.0±0.0	6.0±0.0	19.0±0.0	7.0±0.0	0.0±0.0	0.0±0.0
<i>Bacillus mycoloides</i> (VAB-6)	0.0±0.0	0.0±0.0	6.0±0.0	12.0±0.0	12.0±0.0	6.0±0.0	7.0±0.0	11.0±0.0	0.0±0.0	0.0±0.0
<i>Serratia nematodiphila</i> (VAB-7)	0.0±0.0	0.0±0.0	5.0±0.0	3.0±0.0	4.0±0.0	5.0±0.0	10.0±0.0	10.0±0.0	0.0±0.0	0.0±0.0
Unknown (VAB-8)	0.0±0.0	0.0±0.0	6.0±0.0	18.0±0.0	6.0±0.0	6.0±0.0	2.0±0.0	15.0±0.0	0.0±0.0	0.0±0.0
Unknown (VAB-9)	0.0±0.0	0.0±0.0	5.0±0.0	18.0±0.0	7.0±0.0	6.0±0.0	0.0±0.0	14.0±0.0	0.0±0.0	0.0±0.0
<i>Serratia marcescens</i> (VAB-10)	0.0±0.0	0.0±0.0	3.0±0.0	20.0±0.0	6.0±0.0	5.0±0.0	0.0±0.0	13.0±0.0	0.0±0.0	0.0±0.0
Unknown (VAB-11)	19.0±0.0	0.0±0.0	14.0±0.0	16.0±0.0	12.0±0.0	15.0±0.0	0.0±0.0	17.0±0.0	0.0±0.0	0.0±0.0
<i>Serratia marcescens</i> (VAB-12)	0.0±0.0	0.0±0.0	5.0±0.0	20.0±0.0	4.0±0.0	4.0±0.0	10.0±0.0	11.0±0.0	0.0±0.0	4.0±0.0
<i>Paenibacillus dendritiformis</i> (VAB-14)	0.0±0.0	0.0±0.0	16.0±0.0	8.0±0.0	8.0±0.0	5.0±0.0	12.0±0.0	17.0±0.0	5.0±0.0	5.0±0.0

DISCUSSION

Vermicompost associated bacteria

Based on the morphological, microscopic, biochemical tests, and molecular characterization, 10 VAB strains were identified viz. *Achromobacter ruhlandii*, *Bacillus wiedmannii*, *Pseudomonas* sp., *Achromobacter xylosoxidans*, *Oceanobacillus oncorhynchi*, *Bacillus mycoides*, *Serratia nematodiphila*, *Serratia marcescens*, and *Paenibacillus dendritiformis*. The current results agreed with previous literature (Upadhyay *et al.*, 2015; Selvi and Koilraj, 2015; Begum and Bora, 2018). Khyade (2018) and Vaz-Moreira *et al.* (2008) also demonstrated the presence of *B. cereus*, *B. pumilus*, *B. macrolides*, *B. licheniformis*, *Bacillus benzoovorans*, *B. megaterium*, and *B. subtilis* from vermicompost. Several bacteria i.e. *Pseudomonas*, *Klebsiella*, *Bacillus*, *Serratia*, *Azospirillum*, *Acetobacter*, *Burkholderia*, and *Azotobacter*, have been recorded as plant growth-promoting bacteria (PGPB) while *Pseudomonas* and *Bacillus* spp. have been identified as the predominant communities. Similar results were found by Kang *et al.* (2014), and our findings agreed with them. Andleeb *et al.* (2022) isolated and identified eleven vermibacteria from the gut of *E. fetida* such as *B. megaterium*, *Bacillus mycoides*, *Staphylococcus hominis*, *B. aryabhattai*, *B. subtilis*, *B. licheniformis*, *B. spizizenii*, *B. mojavensis*, *B. cereus*, *B. toyonensis*, *B. thuringiensis*, *B. anthracis*, and *B. paranthracis*. which enhanced the quality of vermicompost during vermicomposting?

Vermicompost associated bacteria possessed agricultural traits

Vermibacteria associated with the gut of *E. fetida* possessed agricultural traits and their impact was done on ornamental plants. These plant growth promoting vermibacteria (PGPVB) not only used for the develop of plant growth but also enhance the plant nutrition value either directly or indirectly mechanisms (Andleeb *et al.* 2022). The direct assistance of plant growth is carried out by various ways such as supplementation of essential nutrients such as phosphate, nitrogen, potassium, zinc, and iron (Divjot *et al.*, 2020; Fasciglione *et al.*, 2015). Similarly, PGPB also enhanced the plant growth by the production of phytohormones (Al-Kahtani *et al.*, 2020). On the other side, the indirect plant growth promotion occurs through the prevention of deleterious effects of pathogens and pest on plants by releasing compounds or defense enzymes/ proteins by PGPBs (Simon *et al.*, 2019; Frampton *et al.*, 2012). Several reports have also been known for the imperative role of PGPBs in mitigating salt stress in different crop plants tomato, groundnut, wheat, rice and red pepper (Upadhyay and Singh, 2015; Bal *et al.*, 2013; Shukla *et al.*, 2012).

The current results showed that vermicompost associated bacterial strains i.e., *Achromobacter ruhlandii*, *Bacillus wiedmannii*, *Pseudomonas* sp., *Achromobacter xylosoxidans*, *Oceanobacillus oncorhynchi*, *Bacillus mycoides*, *Serratia nematodiphila*, *Serratia marcescens*, and *Paenibacillus dendritiformis* have ability to produce siderophores, hydrolytic enzymes, hydrogen cyanide, ammonia, and IAA, and solubilize phosphates and our findings agreed with the outcomes of Mahdi *et al.* (2020) and Ahemad and Kibret (2014). *Achromobacter* spp. are endophytic bacteria, showed plant growth promoting properties like production of IAA, HCN, and ammonia, and have phosphate solubilizing activity. *Oceanobacillus oncorhynchi* is gram-positive rod shaped endophytic bacteria, and also possessed agricultural traits. Our outcomes agreed with previous reports indicated that *Oceanobacillus oncorhynchi* and *Achromobacter* spp could be used as PGPB (Mapelli *et al.*, 2013; Orhan, 2016; Jha and Kumar, 2009).

Key role of Vermicompost associated bacteria as PGPB

The current study reveals that vermicompost associated bacteria possessed osphate solubilizing capability and phosphorous solubilizing microbes play an important role directly and indirectly in biological, physical, and chemical soil properties, and agreed with previous studies (Guo *et al.*, 2015; Alori *et al.*, 2017). The presence of phosphate solubilizing bacterium in soils as well as vermicompost may be considered a positive indicator of utilizing the microbes as biofertilizers for crop production and beneficial for sustainable agriculture. In addition, phosphorous solubilization, VAB involved in the IAA and siderophore production which promotes the seed germination, enhance root development, biosynthesis of various metabolites, initiate florescence, enhances root surface area and root length, resistance to biotic and abiotic stresses, and nodule formation. Bacterial isolates also able to produce ammonia which indicates that VAB supply nitrogen to their plants and not only capable to promote shoot and root elongation as well as biomass of the plant, and our findings are consistent with the outcomes of Etesami and Beattie (2018), Mahanty *et al.* (2017), and Dutta *et al.* (2015). The nitrogen fixation ability of associative bacteria provides the essential nitrogen content during the growth phase of the plant (Jha and Kumar, 2009). It was observed that *Bacillus* species such as *B. licheniformis*, *B. cereus*, and *B. subtilis* involved in the enhancement of root colonization, phytohormone production, enhanced the plant growth and possessed biocontrol properties due to presence of agricultural traits. Our findings agreed with previous (Radhakrishnan and Lee, 2016; Allard-Massicotte *et al.*, 2016; Rijavec and

Lapanje, 2016; Islam *et al.*, 2014; Beauregard *et al.*, 2013).

VABs are source of spread of antibiotic resistant genes

The major concern on the development of multi-drug resistant microbes is increasing day by day due to extensive use of antibiotics in livestock. These antibiotics alter the gut microbiota of the animals that started to attain antibiotic resistance and raises the transfer of antibiotic resistant genes to rhizosphere bacteria through horizontal gene transfer (Tariq *et al.*, 2022). These bacteria might be involved in the spread of resistome to human pathogens or can infect humans through contaminated vegetables and fruits (Checcucci *et al.*, 2020; Zeng *et al.*, 2018). According to Gonzalez *et al.* (2017), some PGPB involved in the human infections on exposure of contaminated soil, water and farm products. So, in the current study, screening of antibiotic susceptibility tests was necessary to know the status of VABs that could be used as microbial biofertilizers. Results revealed that all vermibacterial isolates are not multidrug resistant bacteria and their presence in the fields reduced the possibility of transfer of resistome in foodchain.

Current study reveals that vermicompost associated bacteria i.e., *Achromobacter* spp., *Bacillus* spp., *Pseudomonas* sp., *Oceanobacillus oncorhynchi*, *Serratia* spp., and *Paenibacillus dendritiformis* could be used as microbial biofertilizers to enhance crop production, involved in vermicomposting, improves the vermicompost quality and soil health, and used as a biocontrol agent due to presence of agricultural traits.

DECLARATIONS

Acknowledgements

Authors are thankful to Transfer Technology Funds (TDF), HEC for providing funds (TDF-02-006) for current research work and also thankful to Department of Biochemistry, Islamia University Bahawalpur, Bahawalpur for the identification of vermibacterial isolates.

Funding

Whole research work is done under HEC funded project No. TDF-02-006 titled: "Establishment of the vermi-tech unit at Azad Jammu and Kashmir, Muzaffarabad for vermi product development". The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Ethical statement and approval

All experiments conducted during the research were carefully designed to prevent distress, suffering, and unnecessary pain to the experimental animals.

All procedures were carried out in accordance with international regulations, specifically following Article 9 of the Dutch Law on animal experimentation (Wet op de dierproeven). The current study was approved by Institutional Review Committee of Office of Research, Innovation, and Commercialization (ORIC), The University of Azad Jammu and Kashmir, Muzaffarabad vide No. 246/ORIC/2022; Dated: 3-10-2022.

Availability of data and materials

All the data generated during this study are included in this article.

Supplementary material

There is supplementary material associated with this article. Access the material online at: <https://dx.doi.org/10.17582/journal.pjz/20230525060540>

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Ahemad, M. and Kibret, M., 2014. Mechanisms and applications of plant growth promoting rhizobacteria: Current perspective. *J. King Saud Uni. Sci.*, **26**: 1–20. <https://doi.org/10.1016/j.jksus.2013.05.001>
- Alkahtani, M.D.F., Fouda, A., Attia, K., Al-Otaibi, F., Eid, A.M., Ewais, E., Hijri, M., St-Arnaud, M., Hassan, S., Khan, N., Yaser M.H. and Khaled, A.A.A., 2020. Isolation and characterization of plant growth promoting endophytic bacteria from desert plants and their application as bioinoculants for sustainable agriculture. *Agronomy*, **10**: 1325. <https://doi.org/10.3390/agronomy10091325>
- Allard-Massicotte R.L., Tessier, F., Lecuyer, V., Lakshmanan, J.F., Lucier, D., Garneau, L., Caudwell, H., Vlamakis, H.P. and Bais, P.B., 2016. Beauregard. *Bacillus subtilis* early colonization of *Arabidopsis thaliana* roots involves multiple chemotaxis receptors. *MBio*, **7**: e01664-16. <https://doi.org/10.1128/mBio.01664-16>
- Alori, E.T. Dare, M.O. and Babalola, O.O., 2017. Microbial inoculants for soil quality and plant health. *Sustainable agriculture reviews*. Springer, pp. 281-307. https://doi.org/10.1007/978-3-319-48006-0_9
- Andleeb, S, Shafique, I., Naseer, A., Abbasi, W.A., Ejaz, S., Liaqat, I., Ali, S., Khan, M.F., Ahmed, F. and Ali, N.M., 2022. Molecular characterization of plant growth-promoting vermi-bacteria associated

- with *Eisenia fetida* gastrointestinal tract. *PLoS One*, **17**: e0269946. <https://doi.org/10.1371/journal.pone.0269946>
- Bal, H.B., Nayak, L., Das, S. and Adhya, T.K., 2013. Isolation of ACC deaminase PGPR from rice rhizosphere and evaluating their plant growth promoting activity under salt stress. *Pl. Soil*, **366**: 93-105. <https://doi.org/10.1007/s11104-012-1402-5>
- Bauer, A.W., Perry, D.M. and Kirby, W.M., 1959. Single-disk antibiotic-sensitivity testing of staphylococci: An analysis of technique and results. *Arch. Intern. Med.*, **104**: 208-216. <https://doi.org/10.1001/archinte.1959.00270080034004>
- Beauregard, P.B., Chai, Y.R., Vlamakis, H., Losick, R. and Kolter, R., 2013. *Bacillus subtilis* biofilm induction by plant polysaccharides. *Proc. natl. Acad. Sci. U.S.A.*, pp. E1621-E1630. <https://doi.org/10.1073/pnas.1218984110>
- Begum, M. and Bora, P., 2018. Isolation and identification of bacterial strains in vermicompost and vermiwash. *Int. J. Rec. Res. Rev.*, **1**: 3-567
- Boruah, T., Barman, A., Kalita, P., Lahkar, J. and Deka, H., 2019. Vermicomposting of citronella bagasse and paper mill sludge mixture employing *Eisenia fetida*. *Bioresour. Technol.*, **294**: 122147. <https://doi.org/10.1016/j.biortech.2019.122147>
- Brown, D.F. and Kothari, D., 1975. Comparison of antibiotic discs from different sources. *J. clin. Pathol.*, **28**: 779-783. <https://doi.org/10.1136/jcp.28.10.779>
- Chauhan, H.K. and Singh, K., 2015. Potancy of Vermiwash with neem plant parts on the infestation of *Earias vittella* (Fabricius) and productivity of okra (*Abelmoschus esculentus* L.) Moench. *Asian J. Res. Pharma. Sci.*, **5**: 36-40. <https://doi.org/10.5958/2231-5659.2015.00006.5>
- Checucci, A., Trevisi, P., Luise, D., Modesto, M., Blasioli, S., Braschi, I. and Mattarelli, P., 2020. Exploring the animal waste resistome: The spread of antimicrobial resistance genes through the use of livestock manure. *Front. Microbiol.*, **11**: 1416. <https://doi.org/10.3389/fmicb.2020.01416>
- Chitrapriya, K., Asokan, S. and Nagarajan, R., 2013. Estimating the level of phosphate solubilizing bacteria and Azotobacter in the vermicompost of *Eudrilus eugeniae* and *Perionyx excavatus* with various combinations of cow-dung and saw-dust. *Int. J. Sci. Res. Publ.*, **3**: 1-4.
- Dawwam, G.E., Elbeltagy, A., Emara, H.M., Abbas, I.H. and Hassan, M.M., 2013. Beneficial effect of plant growth promoting bacteria isolated from the roots of potato plant. *Ann Appl. Stat.*, **58**: 195-201.
- de Lillo, A., Ashley, F.P., Palmer, R.M., Munson, M.A., Kyriacou, L., Weightman, A.J. and Wade, W.G., 2006. Novel subgingival bacterial phylotypes detected using multiple universal polymerase chain reaction primer sets. *Oral Microbiol. Immunol.*, **21**: 61-68. <https://doi.org/10.1111/j.1399-302X.2005.00255.x>
- Deka, H., Deka, S., Baruah, C.K., Das, J., Hoque, S., Sarma, H. and Sarma, N.S., 2011. Vermicomposting potentiality of perionyx excavatus for recycling of waste biomass of java citronella-an aromatic oil yielding plant. *Bioresour. Technol.*, **102**: 11212-11217. <https://doi.org/10.1016/j.biortech.2011.09.102>
- Devi, S.H., Vijayalakshmi, K., Jyotsna, K.P., Shaheen, S.K., Jyothi, K. and Rani, M.S., 2009. Comparative assessment in enzyme activities and microbial populations during normal and vermicomposting. *J. environ. Biol.*, **30**: 1013-1017.
- Divjot, K., Kusam, L.R., Yadav, A.N., Sheikh, I., Kumar, V., Dhaliwal, H.S. and Saxena, A.K., 2020. Amelioration of drought stress in Foxtail millet (*Setaria italica* L.) by P solubilizing drought tolerant microbes with multifarious plant growth promoting attributes. *Environ. Sustain.*, **3**: 23-34. <https://doi.org/10.1007/s42398-020-00094-1>
- Dutta J., Handique, P.J. and Thakur, D., 2015. Assessment of culturable tea rhizobacteria isolated from tea estates of Assam, India for growth promotion in commercial tea cultivars. *Front. Microbiol.*, **6**: 1252. <https://doi.org/10.3389/fmicb.2015.01252>
- Emperor, G.N. and Kumar, K., 2015. Microbial population and activity on vermicompost of *Eudrilus eugeniae* and *Eisenia fetida* in different concentrations of tea waste with cow dung and kitchen waste mixture. *Int. J. Curr. Microbiol. appl. Sci.*, **4**: 496-507.
- Etesami, H. and Beattie, G.A., 2018. Mining halophytes for plant growth-promoting halotolerant bacteria to enhance the salinity tolerance of non-halophytic crops. *Front. Microbiol.*, **9**: 148. <https://doi.org/10.3389/fmicb.2018.00148>
- Fasciglione, G., Casanovas, E.M., Quillehauquy, V., Yommi, A.K., Goñi, M.G., Roura, S.I. and Barassi, C.A., 2015. Azospirillum inoculation effects on growth, product quality and storage life of lettuce plants grown under salt stress. *Sci. Hortic.*, **195**: 154-162. <https://doi.org/10.1016/j.scienta.2015.09.015>
- Frampton, R.A., Pitman, A.R. and Fineran, P.C., 2012. Advances in bacteriophage-mediated control of

- plant pathogens. *Int. J. Microbiol.*, **2012**: 326452. <https://doi.org/10.1155/2012/326452>
- González, O., Ortiz-Castro, R., DiAz-Pe'Rez, C., Di 'Az-Pe'Rez, A.L., Magan~aduen~as, V., Lo'Pez-Bucio, J. and Campos-García, J., 2017. Nonribosomal peptide synthases from *Pseudomonas aeruginosa* play a role in cyclodipeptide biosynthesis, quorum-sensing regulation, and root development in a plant host. *Microb. Ecol.*, **73**: 616–629. <https://doi.org/10.1007/s00248-016-0896-4>
- González-Moreno, M.Á., García Gracianteparaluceta, B., Marcelino Sádaba, S., Prieto Cobo, E. and Seco Meneses, A., 2022. Vermicomposting of lavender waste: A biological laboratory investigation. *Agronomy*, **12**: 2957. <https://doi.org/10.3390/agronomy12122957>
- Guo, L., Wu, G., Li, C., Liu, W., Yu, X., Cheng, D. and Jiang, G., 2015. Vermicomposting with maize increases agricultural benefits by 304%. *Agron. Sustain. Dev.*, **35**: 1149–1155. <https://doi.org/10.1007/s13593-015-0307-0>
- Huang, K., Li, F., Wei, Y., Chen, X. and Fu, X., 2013. Changes of bacterial and fungal community compositions during vermicomposting of vegetable wastes by *Eisenia foetida*. *Bioresour. Technol.*, **150**: 235–241. <https://doi.org/10.1016/j.biortech.2013.10.006>
- Islam, F., Yasmeen, T., Ali, Q., Ali, S., Arif, M.S., Hussain, S. and Rizvi, H., 2014. Influence of *Pseudomonas aeruginosa* as PGPR on oxidative stress tolerance in wheat under Zn stress. *Ecotoxicol. environ. Saf.*, **104**: 285–293. <https://doi.org/10.1016/j.ecoenv.2014.03.008>
- Jha P. and Kumar, A., 2009. Characterization of novel plant growth promoting endophytic bacterium *Achromobacter xylosoxidans* from wheat plant. *Microbiol. Ecol.*, **58**: 179–188. <https://doi.org/10.1007/s00248-009-9485-0>
- Kang, S.M., Radhakrishnan, R., You, Y.H., Joo, G.J., Lee, I.J., Lee, K.E. and Kim, J.H., 2014. Phosphate solubilizing *Bacillus megaterium* mjl212 regulates endogenous plant carbohydrates and amino acids contents to promote mustard plant growth. *Ind. J. Microbiol.*, **54**: 427–433. <https://doi.org/10.1007/s12088-014-0476-6>
- Khyade, V.B., 2018. Bacterial diversity in the alimentary canal of earthworms. *J. Bact. Mycol. Open Access.*, **6**: 183–185. <https://doi.org/10.15406/jbmoa.2018.06.00200>
- Kumar, J., Bahadur, A., Maurya, I., Raghuwanshi, B., Meena, R.S., Singh, D. and Dixit, J., 2015. Does a plant growth promoting rhizobacteria enhance agricultural Sustainability. *J Pure Appl Microbiol* 2015a; 9: 715–724.
- Kumar, S., Stecher, G., Li, M., Knyaz, C. and Tamura, K., 2018. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol.*, **35**: 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Lazcano, C., Gomez-Brandon, M. and Dominguez, J., 2008. Comparison of the effectiveness of composting and vermicomposting for the biological stabilization of cattle manure. *Chemosphere*, **72**: 1013–1019. <https://doi.org/10.1016/j.chemosphere.2008.04.016>
- Mahanty, T., Bhattacharjee, S., Goswami, M., Bhattacharyya, P., Das, B., Ghosh, A. and Tribedi, P., 2017. Biofertilizers: A potential approach for sustainable agriculture development. *Environ. Sci. Pollut. Res.*, **24**: 3315–3335. <https://doi.org/10.1007/s11356-016-8104-0>
- Mahdi, I., Fahsi, N., Hafidi, M., Allaoui, A. and Biskri, L., 2020. Plant growth enhancement using rhizospheric halotolerant phosphate solubilizing bacterium *Bacillus licheniformis* QA1 and *Enterobacter asburiae* QF11 isolated from *Chenopodium quinoa* Willd. *Microorganisms*, **8**: 948. <https://doi.org/10.3390/microorganisms8060948>
- Mahdi, I., Fahsi, N., Hijri, M. and Sobeh, M., 2022. Antibiotic resistance in plant growth promoting bacteria: A comprehensive review and future perspectives to mitigate potential gene invasion risks. *Front. Microbiol.*, **13**: 999988. <https://doi.org/10.3389/fmicb.2022.999988>
- Mapelli, F., Marasco, R., Rolli, E., Barbato, M., Cherif, H., Guesmi, A., Ouzari, I., Daffonchio, D. and Borin, S., 2013. Potential for plant growth promotion of rhizobacteria associated with *Salicornia* growing in Tunisian hypersaline soils. *Biomed. Res. Int.*, **2013**: 1–13. <https://doi.org/10.1155/2013/248078>
- Okon, L.S. and Albercht, B.R.H., 1977. Methods of growing *Spirillum lipoferum* and for counting it in pure culture and in association with plants. *Appl. Environ. Microbiol.*, **66**: 2445–2450. <https://doi.org/10.1128/aem.33.1.85-88.1977>
- Orhan, F., 2016. Alleviation of salt stress by halotolerant and halophilic plant growth-promoting bacteria in wheat (*Triticum aestivum*). *Braz. J. Microbiol.*, **47**: 621–627. <https://doi.org/10.1016/j.bjm.2016.04.001>
- Pathma, J. and Sakthivel, N., 2012. Microbial diversity of vermicompost bacteria that exhibit useful agricultural traits and waste management potential.

- Springer Plus, **1**: 26. <https://doi.org/10.1186/2193-1801-1-26>
- Radhakrishnan, R. and Lee, I.J., 2016. Gibberellins producing *Bacillus methylotrophicus* KE2 supports plant growth and enhances nutritional metabolites and food values of lettuce. *Pl. Physiol. Biochem.*, **109**: 181–189. <https://doi.org/10.1016/j.plaphy.2016.09.018>
- Ravindran, B., Wong, J.W., Selvam, A. and Sekaran, G., 2016. Influence of microbial diversity and plant growth hormones in compost and vermicompost from fermented tannery waste. *Bioresour. Technol.*, **217**: 200–204. <https://doi.org/10.1016/j.biortech.2016.03.032>
- Rijavec, T. and Lapanje, A., 2016. Hydrogen cyanide in the rhizosphere: Not suppressing plant pathogens, but rather regulating availability of phosphate. *Front. Microbiol.*, **7**: 1785. <https://doi.org/10.3389/fmicb.2016.01785>
- Seeley, H.W., Vandemark, P.J. and Lee, J.J., 2001. *Microbes in action: A laboratory manual of microbiology*. 4th edition, W.H. Freeman and Co., New York, USA. pp. 57–130.
- Selvi, C.P. and Koilraj, A.J., 2015. Bacterial diversity in compost and vermicompost of cotton waste at Courtallam, Nellai District in Tamilnadu, India. *Int. J. Curr. Microbiol. appl. Sci.*, **4**: 582–585.
- Shafique, I., Andleeb, S., Aftab, M.S., Naeem, F., Ali, S., Yahya, S., Fayaz, A., Tauseef, T., Sultan, T., Shahid, B., Khan, A.B., Ilsam, G. and Abbasi, W.A., 2021. Efficiency of cow dung based vermicompost on seed germination and plant growth parameters of *Tagetes erectus* (Marigold). *Heliyon*, **7**: e05895. <https://doi.org/10.1016/j.heliyon.2020.e05895>
- Shukla, P.S., Agarwal, P.K. and Jha, B., 2012. Improved salinity tolerance of *Arachis hypogaea* (L.) by the interaction of halotolerant plant-growth-promoting rhizobacteria *J. Pl. Growth Regul.*, **31**: 195–206. <https://doi.org/10.1007/s00344-011-9231-y>
- Simon, J.C., Marchesi, J., Mougel, C. and Selosse, M.A., 2019. Host-microbiota interactions: From holobiont theory to analysis. *Microbiome*, **7**: 5. <https://doi.org/10.1186/s40168-019-0619-4>
- Somasegaran, P., Hoben, H.J., 1994. *Handbook for rhizobia methods in legume-rhizobium technology*. Springer, Heidelberg, New York. <https://doi.org/10.1007/978-1-4613-8375-8>
- Tamura, K., Dudley, J., Nei, M. and Kumar, S., 2007. MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol. Biol. Evol.*, **24**: 1596–1599. <https://doi.org/10.1093/molbev/msm092>
- Tariq, M., Jameel, F., Ijaz, U., Abdullah, M. and Rashid, K., 2022. Biofertilizer microorganisms accompanying pathogenic attributes: A potential threat. *Physiol. Mol. Biol. Pl.*, **28**: 77–90. <https://doi.org/10.1007/s12298-022-01138-y>
- Upadhyay, S.K. and Singh, D.P., 2015. Effect of salt-tolerant plant growth-promoting rhizobacteria on wheat plants and soil health in a saline environment. *Pl. Biol.*, **17**: 288–293
- Upadhyay, S., Kumar, N., Singh, V.K. and Singh, A., 2015. Isolation, characterization and morphological study of *Azotobacter* isolates. *J. appl. Natl. Sci.*, **7**: 984–990.
- Varma, V.S., Kalamdhad, A.S. and Khwairkham, M., 2016. Feasibility of *eudrilus eugeniae* and *perionyx excavatus* in vermicomposting of water hyacinth. *Ecol. Eng.*, **94**: 127–135. <https://doi.org/10.1016/j.ecoleng.2016.05.058>
- Vaz-Moreira, I., Maria, E., Silva, C.M., Manaia, C.M. and Nunes, O.C., 2008. Diversity of bacterial isolates from commercial and homemade composts. *Microb. Ecol.*, **55**: 714–722. <https://doi.org/10.1007/s00248-007-9314-2>
- Yasir, M., Aslam, Z., Kim, S.W., Lee, S.W., Jeon, C.O. and Chung, Y.R., 2009. Bacterial community composition and chitinase gene diversity of vermicompost with antifungal activity. *Bioresour. Technol.*, **100**: 4396–4403. <https://doi.org/10.1016/j.biortech.2009.04.015>
- Zeng, Q., Xie, J., Li, Y., Gao, T., Xu, C. and Wang, Q., 2018. Comparative genomic and functional analyses of four sequenced *Bacillus cereus* genomes reveal conservation of genes relevant to plant growth promoting traits. *Sci. Rep.*, **8**: 1–10. <https://doi.org/10.1038/s41598-018-35300-y>
- Zhu, F., Hou, J., Xue, S., Wu, C., Wang, Q. and Hartley, W., 2017. Vermicompost and gypsum amendments improve aggregate formation in bauxite residue. *Land Degrad. Dev.*, **28**: 2109–2120. <https://doi.org/10.1002/ldr.2737>