



Antibiogram and Bacterial Profiling of Reproductive Tract Infections in Female Cattle and Buffaloes

Abdul Jabar¹, Ishtiaq Ahmed^{2*}, Aziz ur Rehman², Muhammad Zishan Ahmad³, Muhammad Kamran Rafique², Asim Aslam⁴, Aftab Ahmed Anjum⁵, Ammar Ghani⁴, Irfan Irshad⁶, Arbab Sikandar⁷, Muhammad Tahir Meraj², Qamar un Nisa⁴, Sarfraz-ur-Rahman⁸, Yasmeen Iqbal⁹ and Hafiz Qadeer Ahmed¹

¹Livestock and Dairy Development Department, Government of Punjab, Lahore, 54000, Pakistan.

²Department of Pathobiology, University of Veterinary and Animal Sciences Lahore, Sub-campus Jhang 35200, Pakistan.

³Department of Pathology, Peer Mehr Ali Shah Arid Agriculture University, Rawalpindi, 46000, Pakistan.

⁴Department of Pathology, University of Veterinary and Animal Sciences, Lahore, 54000, Pakistan.

⁵Institute of Microbiology, University of Veterinary and Animal Sciences, Lahore, 54000, Pakistan.

⁶Institute of Continuing Education and Extension, University of Veterinary and Animal Sciences, Lahore, 54000, Pakistan.

⁷Department of Basic Sciences, College of Veterinary and Animal Sciences, Jhang, 35200, Pakistan.

⁸Department of Parasitology, University of Veterinary and Animal Sciences, Lahore, 54000, Pakistan.

⁹Department of Chemistry, University of Agriculture, Faisalabad, 38000, Pakistan.

ABSTRACT

Reproductive tract disorders are considered a major cause of low conception rate and poor productivity of livestock, leading to infertility or sterility of animals, and thus, results in high economic losses. The current study was designed to investigate the role of bacteria in reproductive tract disorders of cattle and buffalo, and the efficacy of different antibiotics against different bacterial isolates. A total of 100 reproductive tracts of slaughtered cattle and buffaloes were randomly collected from different abattoirs of District Lahore and examined for the presence of any gross abnormality. The swab samples were processed for bacteriology on suitable culture media. The bacterial isolates were identified based on colony characteristics and biochemical tests and subjected to *in vitro* antibiotic sensitivity assay. For microscopic examination, tissue samples of reproductive tracts were processed for histopathology. Out of 100 collected samples, 54 samples showed the presence of gross lesions including endometritis (21%), pyometra (17%), Para-ovarian cysts (8%), hydrosalpinx (3%), luteal cysts (3%), and follicular cysts (2%). *E. coli* was the major contaminant isolated from all the samples (100%) followed by *Staphylococcus aureus*, *Micrococcus* species, *Staphylococcus epidermidis*, *Lactobacillus* species, *Corynebacterium* species, and *Aeromonas* species. Histopathological examination showed infiltration of polymorph nuclear (PMN) cells in the endometrium, degeneration of endometrial glands, congestion, hemorrhage, and a proliferation of connective tissue in a few cases. The isolates showed higher sensitivity to norfloxacin, ciprofloxacin, and chlortetracycline while resistant to bacitracin, sulphaphenazole, and chloramphenicol. A significant association was found between the antibiotics and the pattern of their sensitivity. In conclusion, endometritis and pyometra are the leading pathological disorders of the reproductive tract of cattle and buffalo. *E. coli* has been found as the most frequently occurring bacteria in the reproductive tract of female buffaloes and cattle. Though Ciprofloxacin, Norfloxacin, and Chlortetracycline have been investigated as the most effective drugs in the treatment of bacterial infections in female bovine reproductive tracts, further explorations on antimicrobial sensitivity and continuous monitoring of antimicrobial resistance are needed as future considerations.

* Corresponding author: ishtiaqahmed@uvas.edu.pk
0030-9923/2026/0001-0235 \$ 9.00/0



Copyright 2026 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Article Information

Received 09 April 2023

Revised 10 September 2024

Accepted 17 September 2024

Available online 16 January 2025
(early access)

Published 27 December 2025

Authors' Contribution

IA, AA, and AAA designed the research project, supervised the research work, collected and analysed data. AJ, AG, IL, HQA and QN performed sample collection and laboratory procedures. MKR and AUR provided research material and validated laboratory procedures. MTM and MZA performed data analysis and drafted the manuscript. AS, SR, and YI contributed to Lab work and revised the manuscript.

Key words

Antibiotic, Bovine, Microbial profiling, Pathology, Reproduction, Resistance

INTRODUCTION

Pakistan has 29.6 million cattle and 27.3 million buffaloes which are playing a vital role in the national economy by producing a significant proportion of milk and meat (Rehman *et al.*, 2017). Reproductive health is very important for a profitable dairy enterprise. Animals may suffer from acquired and congenital anomalies of the

reproductive tract. Disorders of the reproductive tract cause economic losses to dairy farmers due to increased risk of stillbirth, retained placenta, prolonged calving intervals, extra service charges, treatment expenditure, and culling of animals (Azawi, 2010). The reproductive tract may get infected with specific pathogens via contaminated semen or opportunistic microbes may proliferate under favorable circumstances (Azawi, 2008; Sheldon *et al.*, 2009). Dystocia, uterine prolapse, and retained placenta also predispose animals to get the infection. The most common problems of all commercial dairy farms are endometritis and pyometra (Hosseini-Zadeh and Ardalan, 2011).

Normal microflora in the reproductive tracts of cattle and buffaloes include a variety of beneficial microorganisms. These microbes include, for instance, *E. coli*, *S. epidermidis*, *Y. enterocolitica*, *Lactobacillus* species, and *Micrococcus* species (Azawi, 2010). Some bacteria like *A. pyogenes*, *F. necrophorum*, *C. pyogenes*, *S. aureus* and *S. pyogenes* may lead to severe inflammation of the uterus resulting in endometritis, metritis and pyometra (Bajaj *et al.*, 2018; Al-Dori *et al.*, 2020). Bacterial profiling of the uterine tract in repeat breeders have revealed the presence of *Bacillus* species and *Staphylococcus* species while *E. coli*, *S. aureus*, *A. pyogenes* and *Klebsiella* species have been reported as fatal pathogens during ovarian inactivity (Ahuja *et al.*, 2017).

During the uterine disease process, microorganisms, or pathogen-associated molecules, as well as inflammatory response, cause disruption of uterine tissue and endocrine dysfunction in the reproductive tracts of cattle (Price *et al.*, 2013). Several antibiotics are being used by clinicians to treat post-partum infections of the reproductive tract in bovines (Dubuc *et al.*, 2011). A large variety of antimicrobials are available in the market, being used in routine but excessive and prolonged usage resulted in multidrug resistance (Ma *et al.*, 2018). Antimicrobial susceptibility testing has been conducted worldwide, and different kinds of tetracyclines, cephalosporins, and penicillins are widely used drugs for reproductive tract diseases (Naz *et al.*, 2012; Verma *et al.*, 2018). However, the efficacy of such antimicrobial agents needs to be evaluated continuously due to the emergence of antibiotic resistance of bacterial strains (Frieri *et al.*, 2017).

As there could be several microbes involved in impairing the reproductive efficiency of animals and due to uncontrolled use of antimicrobials in animals in Pakistan which could lead to the emergence of multiple drug-resistant bacteria, continuous monitoring of the reproductive tract pathogens and their susceptibility to various antibiotics are essential. The current investigation has been undertaken to study the gross and histopathological lesions of the female bovine reproductive tract, the bacterial profile of

reproductive tract infections, and the efficacy of different antibiotics against bacteria isolated from reproductive tract infections.

MATERIALS AND METHODS

A total of 100 reproductive tracts were collected randomly from buffaloes (n = 60) and cattle (n = 40) slaughtered at Shera Kot and Shah Pur Kaanran abattoirs, Lahore, Pakistan. Abattoirs were visited twice a week and reproductive tracts were collected in sterilized polythene bags. Samples were brought to the Department of Pathology, University of Veterinary and Animal Sciences, Lahore, for further processing.

The gross lesions in any part of reproductive tracts were visually inspected by incising them completely with sterilized scissors, scalpel, and blade, and any type of pus, hemorrhages, edema, abscess, etc. was recorded. Both ovaries of each sample were examined for any lesions like para-ovarian cysts, follicular cysts (FC), and luteal cysts (LC) and results were noted.

Different culture media (MacConkey's agar, Nutrient agar, and EMB agar) were prepared by following the procedures given in Bergey's Manual of Determinative Bacteriology (Bergey *et al.*, 1994) to isolate and identify the bacterial organisms. The primary culture of microorganisms was obtained by using the streak plate method (Servais and Passerat, 2009) and from this primary culture; the purification of bacteria was done by using the three-dimensional streaking technique (Krishna *et al.*, 2019).

To study the morphology of bacteria, isolated bacteria were stained by the Gram staining method (Beveridge, 2001) and subjected to different biochemical tests to differentiate bacteria at genus and species level. Lactose fermentation test, Indole production test, Methyl red test, Voges Proskauer test, Citrate utilization test (Bergey *et al.*, 1994), Catalase test (Gruner *et al.*, 2007), Mannitol fermentation test (Makwana *et al.*, 2012), Glucose fermentation test (Abbott *et al.*, 2003), Starch hydrolysis test (Iverson and Millis, 1974), and test for the requirement of sodium for bacterial growth was used for differentiation among different bacterial isolates. Moreover, in addition to Gram's staining, spore staining and acid-fast staining were used for the identification of certain spore-forming and acid-fast bacteria respectively (Reynolds *et al.*, 2009).

Antibiotic sensitivity of *E. coli*, *S. aureus*, *S. epidermidis*, *Corynebacterium* species, *Aeromonas* species, *Micrococcus* species, and *Lactobacillus* species to different antibiotics was determined by the disk diffusion method in Mueller Hinton agar. The following antimicrobial drugs were used: penicillin (P; 10 µg), streptomycin (S; 10 µg), ticarcillin/clavulanic acid (TC;

75/10 µg), norfloxacin (NOR; 10 µg), sulphaphenazole (Sp; 200 µg), chloramphenicol (C; 20 µg), bacitracin (B; 10 U), trimethoprim (Tr; 5 µg), ciprofloxacin (CIP; 5 µg), gentamicin (CN; 10 µg), chlortetracycline (CT; 30 µg). The results were declared as sensitive (S), intermediate (I), and resistant (R), based on the zone of inhibition given by the Clinical and Laboratory Standards Institute (CLSI).

For histopathology, samples of uterine tissue were preserved in 10% buffered formalin and processed through routine paraffin embedding technique as described by Bancroft and Gamble (2008). Tissue sections were cut at 5 µm and stained with H and E stains for examination of changes in the tissues.

The obtained data (2012 to 2013) was analyzed using the Chi-Square test through IBM SPSS statistics Ver. 22.0. Armonk, NY.

RESULTS

A total of 100 randomly collected reproductive tracts of female buffaloes ($n=60$) and cattle ($n=40$) were inspected for gross lesions, of which 54% showed lesions. Out of the total of 60 reproductive tracts of buffaloes investigated, 34 reproductive tracts had gross pathological lesions including endometritis (13/60), pyometra and pyosalpinx (11/60), para-ovarian cyst (5/60), hydrosalpinx (2/60), luteal cyst (2/60), follicular cyst (1/34). In cattle ($n = 40$), gross lesions were found in 20 samples including endometritis (8/40), pyometra and pyosalpinx (6/40), para-ovarian cyst (3/40), hydrosalpinx (1/40), luteal cyst (1/40), and follicular cyst (1/40). Statistical analysis showed that the prevalence of different gross lesions in the reproductive tract of female bovines is not significantly associated ($p > 0.05$) with the species of the animal. The percentage prevalence of the different lesions found in the reproductive tract of female bovines is presented in Table I.

In endometritis, uterine tissues showed lesions such as hemorrhages in the stratum compactum, congestion, necrosis, and degeneration of endometrial glands, glandular tissue filled with mononuclear cells, destroyed secretory epithelial cells while in metritis, intact endometrial mucosa, the proliferation of connective tissue, infiltration of inflammatory cells and mononuclear cells in the myometrium were observed.

In pyometra samples, a thick layer of exudate on the surface of the endometrium, severe infiltration of mononuclear cells, sloughing of endometrial mucosa, necrosis of endometrial glands and glands filled with a heavy population of mononuclear cells, the proliferation of fibrous connective tissues in between endometrial glands,

many macrophages are also present, and infiltration of polymorphs mononuclear cells was observed in stratum functional.

Table I. Prevalence percentage (%) of gross lesions observed in the reproductive tract of the buffaloes and cattle.

Gross lesions	Buffaloes (n= 60)	Cattle (n = 40)
	Number ¹ (%)	Number ² (%)
Endometritis	13 (21.26 %)	8 (20 %)
Pyometra and pyosalpinx	11 (18.33 %)	6 (15 %)
Para-ovarian cysts	5 (8.33 %)	3 (7.50 %)
Hydrosalpinx	2 (3.33 %)	1 (2.5 %)
Luteal cysts	2 (3.33 %)	1 (2.50 %)
Follicular cysts	1 (1.67 %)	1 (2.50 %)
Total	34 (56.67 %)	20 (50 %)
p-value	0.996	

¹Buffalo showing gross lesions in the reproductive tract. ²Cattle showing of gross lesions in the reproductive tract.

The collected samples were processed for bacterial isolation and identification. All the samples were found positive for the presence of bacteria. A total of 301 bacterial isolates from 100 reproductive tracts of female cattle and buffaloes were identified. These isolates were recognized as *E. coli*, *Staphylococcus* sp., *Micrococcus* sp., *Lactobacillus* sp., *Corynebacterium* sp. (*C. xerosis* and *C. kutscheri*), and *Aeromonas* sp.

Out of 60 reproductive tracts of buffaloes examined for bacteriological examination, *E. coli* (60/60) had the highest prevalence i.e., 100% followed by *S. aureus* (31/60), *Micrococcus* sp. (29/60), *S. epidermidis* (23/60), *Lactobacillus* sp. (19/60), *Corynebacterium* sp. (*C. xerosis* and *C. kutscheri*) (15/60) and *Aeromonas* sp. (3/60) as 51.6%, 48.3%, 38.3%, 31.6%, 20% and 5%, respectively (Fig. 1). Out of 40 reproductive tracts of cattle examined for bacteriological examination, *E. coli* (40/40) had the highest prevalence i.e., 100% followed by *S. aureus* (20/40), *Micrococcus* sp. (18/20), *S. epidermidis* (15/40), *Lactobacillus* sp. (15/40), *Corynebacterium* sp. (15/40), *Aeromonas* sp. (2/40) as 50%, 45%, 42.5%, 37.5%, 22.5%, and 5%, respectively. Statistically, the association between the host animal species (buffalo and cattle) and the presence of certain bacteria in the female bovine's reproductive tract was not found ($p > 0.05$). The percentage prevalence of different bacteria isolated from the reproductive tract of female buffaloes and cattle is shown in Table II.

Table II. Prevalence (%) of different bacteria isolated from the female buffalo and cattle reproductive tracts.

Bacterial isolates	Buffalo (n = 60)	Cattle (n = 40)
	Number (%)	Number (%)
<i>E. coli</i>	60 (100.0%)	40 (100.0 %)
<i>S. aureus</i>	31 (51.6 %)	20 (50.0 %)
<i>Micrococcus</i> spp.	29 (48.3 %)	18 (45.0 %)
<i>S. epidermidis</i>	23 (38.3 %)	17 (42.5 %)
<i>Lactobacillus</i> spp.	19 (31.6 %)	15 (37.5 %)
<i>Corynebacterium</i> spp.	15 (20.0 %)	9 (22.5 %)
<i>Aeromonas</i> spp.	3 (5.0 %)	2 (5.0 %)
p-value	0.998	




Fig. 1. Percentage prevalence (%) of different bacteria isolated from the female buffalo and cattle reproductive tracts.

All species of isolated bacteria including *C. xerosis*, *C. kutscheri*, *S. aureus*, *S. epidermidis*, *Lactobacillus* species, *E. coli*, *Aeromonas* sp., and *Micrococcus* sp. were sensitive to norfloxacin (NOR; 10µg), ciprofloxacin (CIP; 5 µg), and chlortetracycline (CT; 30µg). All species were sensitive to gentamycin (CN; 10µg) except only *Aeromonas* species that were resistant to gentamycin. *C. xerosis*, *C. kutscheri*, *S. aureus*, *S. epidermidis*, and *Lactobacillus* species were found sensitive to penicillin (P; 10µg) whereas *E. coli* and *Aeromonas* species were found resistant to the penicillin (P; 10µg). However, *Micrococcus* species were found intermediately sensitive to penicillin (P; 10 µg). *C. kutscheri*, *S. aureus*, *Lactobacillus* sp., and *Micrococcus* sp. were found sensitive to the streptomycin (S; 10 µg) whereas *C. xerosis*, *E. coli*, and *Aeromonas* sp., were found resistant to the streptomycin (S; 10 µg). However, *S. epidermidis* was found intermediate sensitive against streptomycin (S; 10 µg). *E. coli*, *S. epidermidis*, *Lactobacillus* sp., *Aeromonas* sp. and *Micrococcus* sp. were found sensitive to ticarcillin/clavulanic acid (TC;

75/10 µg), whereas *S. aureus* was found intermediately sensitive to the same drug. Only *S. epidermidis* and *Micrococcus* sp. were sensitive against suphaphenazole (Sp; 200 µg) while all other bacteria were found resistant to the same drug. Chloramphenicol (C; 20µg) was effective against all isolates except *E. coli* and *S. aureus* which were found intermediate sensitive to the same drugs. *S. epidermidis* was found unaffected by chloramphenicol. *Lactobacillus* sp., *Micrococcus* sp. and *S. aureus* were found sensitive to trimethoprim (Tr; 5 µg) whereas the rest of the bacterial isolates were found resistant. All bacterial isolates were resistant to bacitracin (B; 10 U).

Table III presents the efficacy of antibiotics against various strains of gram-negative and gram-positive bacteria. The results indicate the effectiveness, intermediate effectiveness, or ineffectiveness of antibiotics. The analysis reveals a significant correlation ($p < 0.05$) between the bacterial Gram staining characteristics and sensitivity against antibiotics, specifically highlighting that Gram-positive bacteria exhibit significantly higher sensitivity to antibiotics compared to Gram-negative counterparts.

Table III. The sensitivity of different gram bacteria against different anti-biotic discs applied on MHA medium.

Bacteria	Effective antibiotics (n)	Intermediately effective antibiotics (n)	Non-effective antibiotics (n)	Sig.
Gram-negative				
<i>E. coli</i>	4	1	6	0.025*
<i>Aeromonas</i> spp.	4	0	7	
Gram-positive				
<i>Lactobacillus</i> spp.	8	0	3	0.025*
<i>Micrococcus</i> spp.	8	1	2	
<i>C. kutscheri</i>	8	0	3	
<i>C. xerosis</i>	7	0	4	
<i>S. aureus</i>	7	2	2	
<i>S. epidermidis</i>	6	1	4	

*A significant difference ($p < 0.05$) exists between sensitivity of gram negative and positive bacteria i.e., gram positive bacteria are more sensitive to the antibiotics applied than gram negative bacteria

DISCUSSION

Reproductive tract lesions may affect animal breeding either by causing infertility or sterility resulting in high economic losses (Fareed *et al.*, 2017). Gross lesions observed in the current study (54%) agree with the findings of Lazim *et al.* (2008) who reported that 53.3% of animals were affected with genital abnormalities in Iraq. Current

results differ in terms of prevalence from the study of [Mohammad \(2013\)](#) who reported 13.13% of cattle suffering from macroscopic pathologies in Yemen. Two different studies from Ethiopia reported that 36.8% and 35.9% of cattle had gross lesions ([Abalti et al., 2006](#); [Mekibib et al., 2013](#)). However, in another study, the percentage of gross lesions observed in the reproductive tract of buffaloes was slightly higher i.e. 67.3% than that of the present study ([Al-Fahad et al., 2004](#)). These variations in the occurrence of gross lesions may be attributable to differences in breed, dam's age, parity, nutrition, management, geographic location/environment, and number of examined animals ([Sharma et al., 2017](#)).

Gross lesions in the case of endometritis were almost identical to those previously reported research findings ([Lazim et al., 2008](#)). The results of the current study agree with [Ghanem et al. \(2002\)](#) who found 22.4% of cows suffer from endometritis. However, in contrast, a higher prevalence of endometritis in Iranian and Indian buffaloes were found i.e. 44.73% and 72%, respectively ([Vala et al., 2019](#)). Similarly, a clinical study on buffalo reported a higher percentage (48.79%) of animals suffering from endometritis/metritis while only 0.22% with pyometra were detected ([Modi et al., 2011](#)). Moreover, a much lower prevalence rate of endometritis was noted in Egyptian buffaloes i.e. 8.53% ([Hamouda et al., 2020](#)). A low prevalence (3.9% and 4.93%) of endometritis was found and reported in previous studies ([Abalti et al., 2006](#); [Mekibib et al., 2013](#)). [Azawi \(2008\)](#) found 12.3% and 0.5% incidence of endometritis and pyometra. Endometritis can be related to many predisposing factors including some extrinsic factors such as untrained artificial inseminators, unhygienic conditions during parturition, nutrition, and certain intrinsic factors: dystocia, negative energy balance, inflammation of the udder, retention of placenta, and decrease in body calcium, etc. ([Onyango et al., 2014](#); [Adnane et al., 2017](#)). In comparison to the current study, [Azawi and Ali \(2015\)](#) observed 2.2% and 4.9% of pyosalpinx and hydrosalpinx, respectively. A relatively low incidence of hydrosalpinx (1.25%) was observed and reported previously ([Mohammad, 2013](#)). Similarly, [Mekibib et al. \(2013\)](#) reported the incidence of Hydrosalpinx and pyosalpinx as 2.32% and 0.87%, respectively. The prevalence of hydrosalpinx and pyosalpinx in buffalo varies between 0.7%-14.2% and 0.6%-11.9%, respectively ([Purohit, 2014](#)).

In contrast to the present study, observed a low incidence of para-ovarian and luteal cysts (1.25% and 0.63%), and a high incidence of follicular cysts (7.5%), respectively was reported in a previous study ([Mohammad, 2013](#)). [Abalti et al. \(2006\)](#) reported incidence of para-ovarian and ovarian cysts as 1.5% and 3.5%, respectively.

In Iraqi buffaloes, the prevalence of para-ovarian cysts, luteal cysts, and follicular cysts was found at 4.4%, 0.2%, and 1.5%, respectively ([Azawi and Ali, 2015](#)). In another study, conducted on Pakistani non-descript cows, [Ali et al. \(2006\)](#) found the incidence of para-ovarian, luteal, and follicular cysts as 1.81%, 0.9%, and 2.72%, respectively. In comparison to the current study, a lower incidence of para-ovarian cysts (3.40% and 1.74%) luteal cyst incidence (1.5%, 0.87%), and a higher incidence of follicular cysts (8% and 4.35%) was also reported ([Saxena et al., 2006](#); [Mekibib et al., 2013](#)). A review of ovarian and oviduct pathologies by [Purohit, 2014](#) reported that 0.54% of buffalo have persistent corpus luteum while ovarian cysts range between 0.5-1.4%. A deficient luteinizing hormone is supposed to cause cystic ovaries ([Vanholder et al., 2002](#)). Moreover, this variation may be due to differences in sample size, breed, age and feeding practices. Moreover, several factors like breed, age, level of milk production, feeding, management and exercise are suggested as influencing the prevalence of cystic ovaries in cattle ([Nelson et al., 2010](#)).

In the current study, histological findings such as infiltration of the polymorphs and mononuclear cells in the endometrium, desquamation of epithelial cells, glandular degeneration, and peri-glandular fibrosis were in line with the work of other researchers ([Hatipoglu et al., 2002](#); [El-Sakkar et al., 2008](#); [Yilmaz et al., 2012](#)). Inflammation is a common process in the endometrium of postpartum dairy cattle. The presence of PMN cells in the endometrium was thought to be because of endometritis ([Healy et al., 2014](#)). Almost all cows have some inflammatory changes, especially in the epithelium and stratum compactum ([Adnane et al., 2017](#)). During the disease process of the uterus, inflammatory cells come into the endometrium and the superficial epithelium may become necrotic and the number of lymphocytes and neutrophils present in the endometrium increases ([Carneiro et al., 2016](#)). Thus, uterine disease causes a lower conception rate and increases calving intervals. These may result in the culling of the animals ([Sheldon et al., 2009](#)). In the current investigation, the prevalence of the bacterial isolates in normal cyclic animals was as follows: *E. coli* was most abundant followed by *Micrococcus* sp., *Staphylococcus* sp. and *Lactobacillus* sp. Previous studies told that normal microbial flora included *E. coli*, *Staphylococcus* sp., and *Bacillus* sp. ([Gani et al., 2008](#)). While others reported that normal microflora included *Y. enterocolitica*, *Citrobacter diversus*, *Enterococcus faecalis*, *Bacillus* sp., *Micrococcus* spp. and *Pasteurella multocida* ([El-Jakee et al., 2008](#)).

Different species of bacteria i.e. *Enterobacteriaceae* spp., *Escherichia coli*, *Corynebacterium* spp., *Streptococcus* spp., *Staphylococcus aureus*, *Pseudomonas*

spp., and *Bacillus* spp. have been isolated from the reproductive tract of cattle suffering from endometritis (Barman *et al.*, 2013). All the endometriotic or pyometric samples included in this study were positive for *E. coli*. Previous studies reported that endometrial biopsies in cows were positive for *E. coli* and *Streptococcus* sp. (Ata *et al.*, 2010). In this study, the most abundant organism was *E. coli*, it was present in all animals whether they were pregnant, non-pregnant, or aborted. Likewise, there was a 100% prevalence of *E. coli* in the uterus from normal prepubertal cattle and buffaloes (El-Jakee *et al.*, 2008). *E. coli* is known to be present in both normal and endometriotic animals (Hanafi *et al.*, 2008). The *E. coli* invading the endometrium is different from intestinal or extra-intestinal pathogenic *E. coli*, and its more adherent and invasive to the endometrium in comparison to other strains of *E. coli* (Sheldon *et al.*, 2010). *S. aureus* (51.6 %) and *S. epidermidis* (38.3 %) were also recovered in a higher percentage. *E. coli*, *S. aureus*, and *S. epidermidis* were isolated from healthy as well as inflamed uterus (Yilmaz *et al.*, 2012; Raza *et al.*, 2013). *S. aureus* is categorized as a potential uterine pathogen (Dolezel *et al.*, 2010). Liu *et al.* (2013) found *S. aureus* (21.8%) as the major pathogen in cows with endometritis. *Micrococcus*, *S. epidermidis*, and *Lactobacillus* observed in the current study, are more prevalent in healthy uterine tissues and considered opportunistic contaminants (Williams *et al.*, 2007; Raza *et al.*, 2013). *Aeromonas* spp. was reported along with other uterine bacteria and BHV-4 in cases of mild endometritis (Fabian *et al.*, 2008). The postpartum uterus is highly susceptible to the growth of a variety of aerobic and anaerobic microbes though many of them are removed by uterine defense. Some studies using conventional and molecular techniques have detected *E. coli* in very small numbers with a dominance of Fusobacteriaceae and Bacteroidetes from animals suffering from pyometra, endometritis, and repeat breeding (Knudsen *et al.*, 2015; Pothmann *et al.*, 2015). The current study has the limitation of covering only the aerobic microflora.

In the current study, all bacterial isolates were sensitive to chlortetracycline, ciprofloxacin, and norfloxacin and resistant to bacitracin. The results of this study partially agree with the findings of Malinowski *et al.* (2010) who found that *E. coli* was sensitive to norfloxacin and gentamicin while it was resistant to oxytetracycline. Low sensitivity of *T. pyogenes* to tetracycline and intermediate sensitivity to gentamicin has been reported. *E. coli* is highly resistant to tetracycline in a study conducted by Barman *et al.* (2013). Our study supports the findings of Gani *et al.* (2008) who found ciprofloxacin and oxytetracycline effective against uterine bacteria and a resistance of *E. coli* against penicillin. Udhayavel *et al.*

(2013) reported that ceftriaxone, gentamicin, Norfloxacin, and chlortetracycline were effective against uterine pathogens. Oxytetracycline was the best antimicrobial against endometritis and metritis in cows (Frieri *et al.*, 2017). *In vitro*, antibiotic sensitivity showed that the most effective antibiotic was ceftriaxone (64%). The sensitivity of gentamicin, norfloxacin, and chlortetracycline was found as 32% (Udhayavel *et al.*, 2013). In a study conducted in India, it was revealed that the sensitivity of bacteria isolated from the uterine lavage of clinically endometriotic buffaloes was 74.19% and 54.83% against ciprofloxacin and gentamycin, respectively (Bajaj *et al.*, 2018). In contrast to our results, bacteria isolated from the reproductive tract of endometriotic buffaloes have shown higher sensitivity against chloramphenicol (83.9%) and a lower sensitivity against gentamycin (80.06%) (Bhadaniya *et al.*, 2019). Similar to our findings, the sensitivity of *E. coli* isolated from endometriotic buffaloes was found as 90% against gentamycin (Ingale *et al.*, 2016). The variation in different antibiogram studies might be due to the improper large-scale administration and prescription of a variety of antimicrobial drugs and intrinsic factors of bacteria (Marston *et al.*, 2016).

CONCLUSION

The study has revealed that endometritis and pyometra are the most commonly occurring problems of female bovines in Pakistan. *E. coli* has been found as the most commonly occurring bacteria in the reproductive tract of female buffaloes and cattle. Though ciprofloxacin, norfloxacin, and chlortetracycline have been investigated as the most effective drug in the treatment of bacterial infections in female bovine reproductive tracts, further investigations on antimicrobial sensitivity and continuous monitoring of antimicrobial resistance are needed as future considerations.

DECLARATIONS

Ethics approval consent to participate

The ethical approval was taken from the University's Advanced Studies and Research Board of UVAS, Lahore to conduct the proposed research in Lahore.

Data analysis

Data was analyzed by MS Office 2016, GraphPad Prism 5, and SPSS 2016.

Availability of data and materials

Data on this article is available for any sort of publicity after publication.

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Abalti, A., Bekana, M., Woldemeskel, M. and Lobago, F.J., 2006. Female genital tract abnormalities of Zebu cattle slaughtered at Bahir-Dar Town, north-west Ethiopia. *Trop. Anim. Hlth. Prod.*, **38**: 505-510. <https://doi.org/10.1007/s11250-006-4319-2>
- Abbott, S.L., Cheung, W.K. and Janda, J.M., 2003. The genus aeromonas: Biochemical characteristics, atypical reactions, and phenotypic identification schemes. *J. clin. Microbiol.*, **41**: 2348-2357. <https://doi.org/10.1128/JCM.41.6.2348-2357.2003>
- Adnane, M., Kaidi, R., Hanzen, C. and England, G.C., 2017. Risk factors of clinical and subclinical endometritis in cattle: A review. *Turk. J. Vet. Anim. Sci.*, **41**: 1-11. <https://doi.org/10.3906/vet-1603-63>
- Ahuja, A.K., Cheema, R.S., Narang, D. and Dhindsa, S.S., 2017. Bacterial pathogens and antibiotic susceptibility patterns of cervico-vaginal discharges in cross bred repeat breeding heifer cows. *Int. J. Curr. Microbiol. appl. Sci.*, **6**: 1769-1775. <https://doi.org/10.20546/ijemas.2017.606.205>
- Al-Dori, O.A., Ismaeel, M.A. and Noumy, B.S., 2020. Bacterial infection causes of subclinical endometriositis associated with repeated conception failure in cattle. *Eur. J. Biosci.*, **14**: 6267-6274.
- Al-Fahad, T.A., Alwan, A.F. and Ibrahim, N., 2004. Histological and morphological study of abnormal cases of female reproductive system in Iraqi Buffaloes. *Iraqi J. Vet. Sci.*, **15**: 109-115.
- Ali, R., Raza, M., Jabbar, A. and Rasool, M.H., 2006. Pathological studies on reproductive organs of zebu cow. *J. Agric. Soc. Sci.*, **2**: 91-95.
- Ata, A., Turutoglu, H., Kale, M., Gulay, M.S. and Pehlivanoglu, F., 2010. Microbial flora of normal and abnormal cervical mucous discharge associated with reproductive performance of cows and heifers in Estrus. *Asian-Austral. J. Anim. Sci.*, **23**: 1007-1012. <https://doi.org/10.5713/ajas.2010.90645>
- Azawi, O., 2008. Postpartum uterine infection in cattle. Animal reproduction science. *Anim. Reprod. Sci.*, **105**: 187-208. <https://doi.org/10.1016/j.anireprosci.2008.01.010>
- Azawi, O., 2010. Uterine infection in buffalo cows: A review. *Buffalo Bull.*, **29**: 154-171.
- Azawi, O. and Ali, A., 2015. A study on the prevalence of pathological abnormalities of the ovaries and oviducts diagnosed at post mortem of buffaloes in Mosul. *Buffalo Bull.*, **34**: 51-62.
- Bajaj, N., Agrawal, S., Jain, S., Sharma, V., Mourya, A. and Shrivastava, O.J., 2018. Antibioqram and total viable bacteria count in uterine lavage of normal, sub-clinical and clinical endometritic postpartum buffaloes. *Buffalo Bull.*, **37**: 17-24.
- Bancroft, J.D. and Gamble, M., 2008. *Theory and practice of histological techniques*. 6th Edn. Elsevier Health Sciences, UK.
- Barman, P., Yadav, M., Bangthai, A. and Kumar, H., 2013. Antibioqram of bacteria isolated from bovine endometritis. *Vet. Res. Int.*, **1**: 20-24.
- Bergey, D.H., Breed, R.S., Murray, E.G. and Hitchens, A.P., 1994. *Manual of determinative bacteriology*. 9th edn. Lippincott Williams and Wilkins, USA.
- Beveridge, T.J., 2001. Use of the gram stain in microbiology. *Biotech. Histochem.*, **76**: 111-118. <https://doi.org/10.1080/714028139>
- Bhadaniya, A., Prasad, M., Savsani, H., Kalaria, V., Fefar, D., Mathapati, B., Javia, B., Ghodasara, S. and Odedra, M.J., 2019. Bacteriological and their antibioqram studies of endometritis in slaughtered buffaloes. *Buffalo Bull.*, **38**: 229-236.
- Carneiro, L.C., Cronin, J.G. and Sheldon, I.M., 2016. Mechanisms linking bacterial infections of the bovine endometrium to disease and infertility. *Reprod. Biol.*, **16**: 1-7. <https://doi.org/10.1016/j.repbio.2015.12.002>
- Dolezel, R., Palenik, T., Cech, S., Kohoutova, L. and Vyskocil, M.J., 2010. Bacterial contamination of the uterus in cows with various clinical types of metritis and endometritis and use of hydrogen peroxide for intrauterine treatment. *Vet. Med.*, **55**: 504-511. <https://doi.org/10.17221/2938-VETMED>
- Dubuc, J., Duffield, T., Leslie, K., Walton, J. and LeBlanc, S., 2011. Randomized clinical trial of antibiotic and prostaglandin treatments for uterine health and reproductive performance in dairy cows. *J. Dairy Sci.*, **94**: 1325-1338. <https://doi.org/10.3168/jds.2010-3757>
- El-Jakee, J., Ahmed, W., El-Seedy, F. and Abd El-Moez, S., 2008. Bacterial profile of the genital tract in female buffaloes during different reproductive stages. *Glob. Vet.*, **2**: 7-14.
- El-Sakkar, G.H., Ahmed, H.M. and Shahira, H.J., 2008. Histopathological, microbiological and biochemical studies on uteri and ovaries of infertile slaughtered buffaloes in Dakahlia Governorate. *Egypt. J. comp. Pathol. clin. Pathol.*, **21**: 59-76.
- Fabian, K., Makrai, L., Sachse, K., Szeredi, L. and Egyed, L.J., 2008. An investigation of the aetiological role of bovine herpesvirus 4 in bovine

- endometritis. *Vet. J.*, **177**: 289-292. <https://doi.org/10.1016/j.tvjl.2007.04.010>
- Fareed, S.K., Memon, K.H., Kachiwal, A.B., Azhar, S., Brula, M.I., Ali, M. and Khan, T.A., 2017. Prevalence and economic losses of reproductive disorders and mastitis in buffaloes at Karachi, Pakistan. *Indian J. Anim. Res.*, **51**: 1130-1133. <https://doi.org/10.18805/ijar.8602>
- Frieri, M., Kumar, K. and Boutin, A.J., 2017. Antibiotic resistance. *J. Infect. Publ. Hlth.*, **10**: 369-378. <https://doi.org/10.1016/j.jiph.2016.08.007>
- Gani, M., Amin, M., Alam, M., Kayesh, M., Karim, M., Samad, M. and Islam, M., 2008. Bacterial flora associated with repeat breeding and uterine infections in dairy cows. *Bangladesh J. Vet. Med.*, **6**: 79-86. <https://doi.org/10.3329/bjvm.v6i1.1342>
- Ghanem, M., Shalaby, A.H., Sharawy, S. and Saleh, N.J., 2002. Factors leading to endometritis in dairy cows in Egypt with special reference to reproductive performance. *J. Reprod. Dev.*, **48**: 371-375. <https://doi.org/10.1262/jrd.48.371>
- Gruner, B., Han, S.R., Meyer, H.G., Wulf, U., Bhakdi, S. and Siegel, E., 2007. Characterization of a catalase-negative methicillin-resistant *Staphylococcus aureus* strain. *J. clin. Microbiol.*, **45**: 2684-2685. <https://doi.org/10.1128/JCM.02457-06>
- Hamouda, M., Saber, A. and Al-Shabebi, A., 2020. Incidence of infertility in female buffaloes due to some reproductive disorders. *Adv. Anim. Vet. Sci.*, **8**: 1188-1193. <https://doi.org/10.17582/journal.aavs/2020/8.11.1188.1193>
- Hanafi, E.M., Ahmed, W., Abd El-Moez, S.I., El-Khadrawy, H. and Abd El-Hameed, A.R., 2008. Effect of clinical endometritis on ovarian activity and oxidative stress status in Egyptian buffalo-cows. *Am-Eur. J. Agric. environ. Sci.*, **4**: 530-536.
- Hatipoglu, F., Kiran, M., Ortatatl, M., Erer, H. and Ciftci, M., 2002. An abattoir study of genital pathology in cows: I. Ovary and oviduct. *Rev. Med. Vet.*, **153**: 29-34.
- Healy, L.L., Cronin, J.G., and Sheldon, I.M., 2014. Endometrial cells sense and react to tissue damage during infection of the bovine endometrium via interleukin 1. *Sci. Rep.*, **4**: 1-9. <https://doi.org/10.1038/srep07060>
- Hossein-Zadeh, N.G. and Ardalan, M., 2011. Cow-specific risk factors for retained placenta, metritis and clinical mastitis in Holstein cows. *Vet. Res. Commun.*, **35**: 345-354. <https://doi.org/10.1007/s11259-011-9479-5>
- Ingale, A.M., Rai, R., Saminathan, M., Vadhana, P., Hingade, S., Dhama, K., Milton, A. and Singh, R.J., 2016. Isolation, PCR detection, pathotyping and antibiogram profiling of *Escherichia coli* associated with endometritis in buffaloes. *J. Anim. Pl. Sci.*, **26**: 1247-1254.
- Iverson, W.G. and Millis, N.F., 1974. A method for the detection of starch hydrolysis by bacteria. *J. appl. Bact.*, **37**: 443-446. <https://doi.org/10.1111/j.1365-2672.1974.tb00460.x>
- Knudsen, L.R., Karstrup, C.C., Pedersen, H.G., Agerholm, J.S., Jensen, T.K. and Klitgaard, K.J., 2015. Revisiting bovine pyometra. New insights into the disease using a culture-independent deep sequencing approach. *Vet. Microbiol.*, **175**: 319-324. <https://doi.org/10.1016/j.vetmic.2014.12.006>
- Krishna, R., Mudiganti, R. and Ananda, N.S., 2019. β -Galactosidase optimization using response surface methodology. *Drug Invent. Today*, **11**: 1220.
- Lazim, H.E., Ali, A.J. and Azawi, O., 2008. Pathological and anatomical abnormalities affecting buffalo cows reproductive tracts in Mosul. *Iraqi J. Vet. Sci.*, **22**: 59-67. <https://doi.org/10.33899/ijvs.2008.5715>
- Liu, C., Wang, Y., Yang, Z., Cao, Y., Li, D., Liu, W. and Zhang, N.J., 2013. Prevalence and major pathogen causes of dairy cows clinical endometritis in Northeast China. *Asian J. Anim. Vet. Adv.*, **8**: 124-129. <https://doi.org/10.3923/ajava.2013.124.129>
- Ma, Z., Ginn, A., Kang, M., Galvão, K.N. and Jeong, K.C., 2018. Genomic and virulence characterization of intrauterine pathogenic *Escherichia coli* with multi-drug resistance isolated from cow uteri with metritis. *Front. Microbiol.*, **9**: 3137. <https://doi.org/10.3389/fmicb.2018.03137>
- Makwana, G.E., Gadhavi, H. and Sinha, M., 2012. Comparison of tube coagulase test with mannitol fermentation test for diagnosis of *Staphylococcus aureus*. *Natl. J. Integr. Res. Med.*, **3**: 73-75.
- Malinowski, E., Lassa, H., Markiewicz, H., Kaptur, M., Nadolny, M., Niewitecki, W. and Zietara, J.J., 2010. Antimicrobial resistance of aerobic bacteria isolated from the inflamed uterus of cows. *Med. Weter.*, **66**: 192-195.
- Marston, H.D., Dixon, D.M., Knisely, J.M., Palmore, T.N. and Fauci, A.S., 2016. Antimicrobial resistance. *J. Am. med. Assoc.*, **316**: 1193-1204. <https://doi.org/10.1001/jama.2016.11764>
- Mekibib, B., Desta, T. and Tesfaye, D.J., 2013. Gross pathological changes in the reproductive tracts of cows slaughtered at two abattoirs in Southern Ethiopia. *Vet. Med. Anim. Hlth.*, **5**: 46-50.
- Modi, L., Patel, P., Patel, S., Patel, G., Joshi, A. and Suthar, D.J., 2011. Prevalence of reproductive problems in buffalo in Mehsana milk-shed area of

- Gujarat. *Int. J. Agric. Vet. med. Sci.*, **5**: 424-428. <https://doi.org/10.5455/ijavms.20110801094631>
- Mohammad, A.G., 2013. Pathological abnormalities in genital tract of cows slaughtered at Dhamar Abattoirs, Yemen. *Yemeni J. Agric. Vet. Sci.*, **202**: 1-16. <https://doi.org/10.12816/0003907>
- Naz, S., Hanif, A., Maqbool, A., Ahmed, S. and Muhammad, K., 2012. Isolation, characterization and monitoring of antibiotic resistance in *Pasteurella multocida* isolates from buffalo (*Bubalus bubalis*) herds around Lahore. *J. Anim. Pl. Sci.*, **22**: 242-245.
- Nelson, S.T., Martin, A.D. and Østerås, O., 2010. Risk factors associated with cystic ovarian disease in Norwegian dairy cattle. *Acta Vet. Scand.*, **52**: 1-10. <https://doi.org/10.1186/1751-0147-52-60>
- Onyango, J., Deluna, C. and Blackie, N.J., 2014. Risk factors for postpartum uterine infections in dairy herds. *Int. J. Livest. Res.*, **4**: 29-40. <https://doi.org/10.5455/ijlr.20140512103231>
- Pothmann, H., Prunner, I., Wagener, K., Jaureguiberry, M., Sota, R.L., Erber, R., Aurich C., Ehling-Schulz, M. and Drillich, M.J., 2015. The prevalence of subclinical endometritis and intrauterine infections in repeat breeder cows. *Theriogenology*, **83**: 1249-1253. <https://doi.org/10.1016/j.theriogenology.2015.01.013>
- Price, J.C., Bromfield, J.J. and Sheldon, I.M., 2013. Pathogen-associated molecular patterns initiate inflammation and perturb the endocrine function of bovine granulosa cells from ovarian dominant follicles via TLR2 and TLR4 pathways. *Endocrinology*, **154**: 3377-3386. <https://doi.org/10.1210/en.2013-1102>
- Purohit, G.J., 2014. Ovarian and oviductal pathologies in the buffalo: Occurrence, diagnostic and therapeutic approaches. *Asian Pac. J. Reprod.*, **3**: 156-168. [https://doi.org/10.1016/S2305-0500\(14\)60020-8](https://doi.org/10.1016/S2305-0500(14)60020-8)
- Raza, S., Rabbani, M., Ahmad, N., Sheikh, A.A., Muhammad, K., Akhtar, F. and Rehman, H.U., 2013. Uterine microbial flora of Nili-Ravi buffalo during estrus and its relationship with pregnancy Rate in Pakistan. *Buffalo Bull.*, **32**: 564-567.
- Rehman, A., Jingdong, L., Chandio, A.A. and Hussain, I., 2017. Livestock production and population census in Pakistan: Determining their relationship with agricultural GDP using econometric analysis. *Inform. Process. Agric.* **4**: 168-177. <https://doi.org/10.1016/j.inpa.2017.03.002>
- Reynolds, J., Moyes, R.B. and Breakwell, D.P., 2009. Differential staining of bacteria: Acid fast stain. *Curr. Prot. Microbiol.*, **15**: 1-5. <https://doi.org/10.1002/9780471729259.mca03hs15>
- Saxena, G., Rani, S., Danodia, H. and Purohit, G.J., 2006. Pathological conditions in genital tract of female buffaloes (*Bubalus bubalis*). *Pak. Vet. J.*, **26**: 91-93.
- Servais, P. and Passerat, J., 2009. Antimicrobial resistance of fecal bacteria in waters of the Seine river watershed (France). *Sci. Tot. Environ.*, **408**: 365-372. <https://doi.org/10.1016/j.scitotenv.2009.09.042>
- Sharma, A., Singh, M., Kumar, P., Sharma, A., Neelam, A.M. and Sharma, P., 2017. Postpartum uterine infections in cows and factors affecting it. A review. *Int. J. Curr. Microbiol. appl. Sci.*, **6**: 1020-1028. <https://doi.org/10.20546/ijcmas.2017.609.123>
- Sheldon, I.M., Cronin, J., Goetze, L., Donofrio, G. and Schuberth, H.J., 2009. Defining postpartum uterine disease and the mechanisms of infection and immunity in the female reproductive tract in cattle. *Biol. Reprod.*, **81**: 1025-1032. <https://doi.org/10.1095/biolreprod.109.077370>
- Sheldon, I.M., Rycroft, A.N., Dogan, B., Craven, M., Bromfield, J.J., Chandler, A., Roberts, M.H., Price, S.B., Gilbert, R.O. and Simpson, K.W., 2010. Specific strains of *Escherichia coli* are pathogenic for the endometrium of cattle and cause pelvic inflammatory disease in cattle and mice. *PLoS One*, **5**: e9192. <https://doi.org/10.1371/journal.pone.0009192>
- Udhayavel, S., Malmarugan, S., Palanisamy, K. and Rajeswar, J., 2013. Antibioqram pattern of bacteria causing endometritis in cows. *Vet. World*, **6**: 100-102. <https://doi.org/10.5455/vetworld.2013.100-102>
- Vala, K., Solanki, G., Fefar, D., Borakhatariya, D. and Raval, R., 2019. Abattoir survey of genital abnormalities in Jaffrabadi Buffaloes. *Indian J. Vet. Sci. Biotechnol.*, **15**: 61-64. <https://doi.org/10.21887/ijvsbt.15.1.14>
- Vanholder, T., Opsomer, G., Govaere, J., Coryn, M. and de Kruif, A.J., 2002. Cystic ovarian disease in dairy cattle: aetiology, pathogenesis, and risk factors. *Tijdschr Diergeneesk.* **127**: 146-155.
- Verma, H., Rawat, S., Sharma, N., Jaiswal, V., Singh, R. and Harshit, V., 2018. Prevalence, bacterial etiology and antibiotic susceptibility pattern of bovine mastitis in Meerut. *J. Ent. Zool. Stud.*, **6**: 706-709.
- Williams, E.J., Fischer, D.P., Noakes, D.E., England, G.C., Rycroft, A., Dobson, H. and Sheldon, I.M., 2007. The relationship between uterine pathogen growth density and ovarian function in the postpartum dairy cow. *Theriogenology*, **68**: 549-559. <https://doi.org/10.1016/j.theriogenology.2007.05.013>

[theriogenology.2007.04.056](https://doi.org/10.1016/j.theriogenology.2007.04.056)

- Yilmaz, O., Kuyucuoglu, Y., Sevimli, A., Yazici, E. and Ucar, M.J., 2012. Uterine microbiology and histopathology in repeat breeder Anatolian Water buffaloes: An Abattoir study. *Kafkas. Univ. Vet. Fak. Derg.*, **18**: 791-798. <https://doi.org/10.9775/kvfd.2012.6446>