



The *In Vitro* Evaluation of Acaricide Paeonol against Human *Demodex* (Acari: Demodicidae)

Yueye Xu¹, Jingang Xu¹, Yujun Shuai¹, Qiao Teng¹, Huanxin Tu¹, Zhili Ren¹, Qingquan Chang¹, Junjie Guo², Yuanyuan Li^{1,3*}, Xiaoniu Tang^{1,3*} and Jinhong Zhao^{1, 3*}

¹Department of Medical Parasitology, Wannan Medical College, 22#, Wenchang West Road, Wuhu 241002, Anhui Province, China

²Department of Medical Parasitology, Qiqihaer Medical College, Qiqihaer 161000, Heilongjiang, China

³Anhui Provincial Key Laboratory of Biological Macromolecules, Wuhu 241002, Anhui, China

ABSTRACT

Demodex folliculorum and *Demodex brevis* that parasitize humans can cause multiple skin disorders, including pityriasis folliculorum, folliculitis, rosacea, blepharitis, seborrheic dermatitis, and perioral dermatosis. Paeonol is the main component isolated from the root bark of *Paeonia suffruticosa* which exhibits several beneficial effects such as anti-insect, anti-inflammatory, neuroprotective, anti-tumor, and anti-cardiovascular diseases. In this study, to evaluate the effectiveness of paeonol against human *Demodex in vitro*, the paeonol solution was directly used to contact and kill both *Demodex* species *in vitro*. The experiment showed that 40 mg/mL was the minimum effective concentration of paeonol for killing the two mite species; paeonol exhibited more remarkable killing effect on *D. brevis* than on *D. folliculorum*. This result suggests that paeonol has good acaricidal activity against human *Demodex* mite *in vitro*. Moreover, it is more effective against *D. brevis* than *D. folliculorum*.

Article Information

Received 19 September 2024

Revised 05 April 2025

Accepted 16 April 2025

Available online 22 October 2025

(early access)

Published 08 April 2026

Authors' Contribution

JZ, XT and YL designed the study. YX, JX, YS, QT, HT, ZR, QC, and JG conducted the experiment. YX wrote the manuscript.

Key words

Paeonol, *Demodex folliculorum*, *Demodex brevis*, *In vitro*, Acaricide

INTRODUCTION

Demodex, a genus of the family Demodicidae, sub-order Euacarophora, order Acariformes, and Arachnid, are a permanent ectoparasite of humans and mammals. Depending on the host and its morphological characteristics, more than 100 species or subspecies can live on humans, dogs, sheep, cattle, pigs, cats, mice, bats, pandas, and other mammals (Foley *et al.*, 2021; Smith *et al.*, 2022; Wei *et al.*, 2023). Human *Demodex*, including *Demodex brevis* and *Demodex folliculorum*, inhabit the sebaceous glands and follicles of eyelashes and other body hairs, such as the nose, ears, beard, and eyebrows. Under normal conditions, *D. brevis* is more broadly

dispersed throughout the body than *D. folliculorum*, despite the latter being more visible (Norm, 1982; Elston, 2010; Palopoli *et al.*, 2015). *D. folliculorum* and *D. brevis* survive by accessing food in the hair follicle and sebaceous gland cells through their needle-like mouths (Paichitrojjana, 2022). Human *Demodex* species are the most complex and commensal residents, but they can lead to some diseases and clinical symptoms termed demodicosis, such as decreased host immunity, sebaceous hyperplasia, and erythematotelangiectatic rosacea (Elston and Elston, 2014; Forton and De Maertelaer, 2021; Clanner-Engelshofen *et al.*, 2022). Hence, the prevention and control of demodicosis are beneficial for physical and mental health of patients with demodicosis. Currently, the treatment of demodicosis mainly depends on chemical agents, including metronidazole, ivermectin, permethrin, benzoyl benzoate, crotamiton, lindane, and sulfur (Jacob *et al.*, 2019; Paichitrojjana, 2022; Chudzicka-Strugala *et al.*, 2023). Nevertheless, several studies have confirmed that these medicines could induce additional adverse drug reactions (Jacob *et al.*, 2019).

Paeonol (2'-hydroxy-4'-methoxyacetophenone) is the principle bioactive compound isolated from the root bark of moutan cortex (*Paeonia suffruticosa* Andrew) and is known for its natural bioactive phenol with antioxidant

* Corresponding author: zhaojh@wnmc.edu.cn, txniu@163.com, hnlyy198717@163.com
0030-9923/2026/0003-1341 \$ 9.00/0



Copyright 2026 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

properties (Gyawali *et al.*, 2019; Zhang *et al.*, 2019; Adki and Kulkarni, 2020; Wang *et al.*, 2020; Ding *et al.*, 2023). Paeonol has prolonged clinical use history in modern China, and its dosage forms date back to as early as the 1970s (Zhang *et al.*, 2019). The pharmacological activity of paeonol was first recorded in traditional Chinese medicine 50 years ago (Adki and Kulkarni, 2020). Previous studies have shown that paeonol had been extensively used because of its great pharmacological potential as an anti-inflammatory, anti-fungal, anti-viral, and insecticidal compound (Xu *et al.*, 2003; Adki and Kulkarni, 2020; Lv *et al.*, 2020; Li *et al.*, 2021; Tang *et al.*, 2022). Some studies have reported that paeonol is active against multiple species of mites, such as *Dermatophagoides farina*, *Dermatophagoides pteronyssinus*, *Tyrophagus putrescentiae*, and *Aleuroglyphus ovatus* (Kim *et al.*, 2004; Tak *et al.*, 2006; Zou *et al.*, 2023a, 2023b). However, it remains unclear whether paeonol has an acaricidal effect on *Demodex*. The present study was performed to evaluate the acaricidal efficacy of paeonol against human *Demodex*.

MATERIALS AND METHODS

Mite collection

Live mites dwelling in human sebaceous glands and hair follicles were obtained anonymously through the cellophane tape method. Transparent glue was pasted on the nose, nasal groove, forehead, zygoma, and chin of *Demodex*-positive patients until the next day, after being instructed to clean their face with tap water before sleeping. Next, the tape was removed and placed on the center of a new clean slide. Paeonol and paraffin oil were purchased from Shanghai McLean Biochemical Technology Co., Ltd.

Acaricidal activity of paeonol against human *Demodex*

Paeonol was diluted to the following concentrations with paraffin oil at 50 °C: 60, 50, 40, 30, 20, and 10 mg/mL. The living adult mites on the slides with activity intensity were observed, classified, and screened by light microscopy (Nikon, Tokyo, Japan) at 25–27 °C. Subsequently, each selected mite was placed on an independent slide. And, the drug solution to be tested was added to uniformly cover the mite body surface, and the slides were placed in a wet box with 60–70% relative humidity at room temperature. The survival and activity of mites were observed under the microscope and recorded at fixed periods. The mites were considered dead when the body pincers and tarsi were motionless for 1 min and remained unmovable when stimulated with an anatomical needle. The survival time of the mites was recorded. Negative control individuals were also evaluated by treatment with paraffin oil. In the 60, 50, 40, 30, 20 and 10 mg/ml groups, the numbers of

D. folliculorum tested were 30, 31, 31, 33, 32 and 30, respectively; in the same concentration groups, the numbers of *D. brevis* were 27, 28, 27, 28, and 28, severally. In the negative control groups, the numbers of *D. folliculorum* and *D. brevis* were 10 and 13.

Statistical analysis

All statistical analyses were performed using IBM-SPSS software version 26 (IBM, Armonk, NY, USA). The primary statistical methods included the chi-square test and linear regression analysis.

RESULTS

Viability of paeonol against human *Demodex*

The mites were observed under a light microscope (Nikon, Tokyo, Japan) (Fig. 1). The initial state of the mites revealed that their holonomic bodies were active, with five or six tarsi; they also moved 6–8 times in a min (Fig. 2A, D). Following exposure to the paeonol solution, the bodies of both *D. folliculorum* and *D. brevis* significantly contracted and twisted; the body condition (more than six tarsi moved for more than 10 times in 1 min) was more strenuous than the initial activity state (Fig. 2B, E). After considerable struggle, their bodies gradually returned to their original appearance. The movement of the mites was delayed, podosoma and gnathosoma were expanded, opisthosoma was shortened, and the bodies of the mites became slightly more transparent than their primal form on death. There was no sign of movement in their pincers and tarsi for 1 min, even these parts were stimulated with a skinny anatomical needle, particularly in the 60, 50, and 40 mg/mL concentration groups (Fig. 2C, F). In the final stage of the 8-h observation period, the mites showed weak motor performance; only one leg moved 1–2 times in 1 min in response to the abovementioned stimulation, and the bodies of the mites did not change in the 30 mg/mL concentration group. The groups treated with paeonol at



Fig. 1. (A) Human *Demodex brevis* (Acari: Demodicidae). (B) Human *Demodex folliculorum* (Acari: Demodicidae). (C) Roots of moutan cortex. (D) Chemical structure of paeonol.




Fig. 2. Morphological changes in human *Demodex* mites following treatment with paeonol ($\times 400$). (A) *D. folliculorum* before treatment with paeonol. (B) *D. folliculorum* exposed to paeonol. (C) dead *D. folliculorum*. (D) *D. brevis* before treatment with paeonol. (E) *D. brevis* exposed to paeonol. (F) dead *D. brevis*. (G) *D. folliculorum* was not completely dead; however, all rotary movements of appendages of the mite had ceased. (H) *D. brevis* was not completely dead; however, all rotary movements of appendages of the mite had ceased. (I) Untreated *D. folliculorum*. The arrows indicate some coelom and/or internal hydroskeleton, which may help the body of the mite to move by decreasing the contact surface to reduce friction when walking.

10 or 20 mg/mL concentrations showed relatively vigorous movements without any needle, with the movement of two or three tarsi 3–5 times per min.

Acaricidal effect of paeonol on *D. folliculorum*

As shown in Table I no recorded deaths occurred in the pure paraffin oil negative control group. However, in the test groups, mite death was observed in the 60, 50, 40, 30, 20, and 10 mg/mL paeonol solutions containing paraffin oil for 8 h. Hence, the average dead time at 60, 50, and 40 mg/mL paeonol concentrations was 118, 142.2, and 240 min, respectively; furthermore, the average dead time was more than 480 min for 30, 20, and 10 mg/mL paeonol concentrations. The mortality rates in 8 h were 100%, 100%, 90.3%, 6.0%, 3.1%, and 3.3%, respectively. The results showed that with decreasing drug concentrations, the acaricidal effect faded over time, and the death time increased incrementally. No significant difference was observed between the 30 mg/mL group and the negative control group, the 20 mg/mL group and the negative control group, and the 10 mg/mL group and the negative control group ($\chi^2 = 0.001$, $P > 0.05$; $\chi^2 = 0.001$, $P > 0.05$; χ^2

$= 0.001$, $P > 0.05$) based on the statistical analysis of the 8-h mortality period. A significant difference in mortality was noted between the 60, 50, and 40 mg/mL groups and the negative control group ($\chi^2 = 34.844$, $P < 0.05$; $\chi^2 = 35.757$, $P < 0.05$; $\chi^2 = 24.468$, $P < 0.05$). Therefore, the 40 mg/mL dose was the lowest effective concentration of paeonol on *D. folliculorum* in this study in 8 h.

Table I. Acaricidal activities of paeonol against *D. folliculorum* and *D. brevis* at different concentrations for 8 h.

Concentration (mg/mL)	Total number	Dead	Mortality (%)	Average dead time (min)	Min. survival time (min)	Max. survival time (min)
<i>D. folliculorum</i>						
60	30	30	100.0	118.2	81	180
50	31	31	100.0	142.2	72	242
40	31	28	90.3	240	145	>480
30	33	2	6.0	>480	362	>480
20	32	1	3.1	>480	422	>480
10	30	1	3.3	>480	436	>480
Paraffin oil	10	0	0	>480	>480	>480
<i>D. brevis</i>						
60	27	27	100	98.6	59	180
50	28	28	100	123.4	63	195
40	27	25	92.6	181	96	>480
30	28	5	17.8	>480	322	>480
20	28	2	7.1	>480	391	>480
10	28	2	7.1	>480	391	>480
Paraffin oil	13	0	0	>480	>480	>480

Acaricidal effect of paeonol on *D. brevis*

Table I shows that the average dead time was 98.6, 123.4, and 181 min for the 60, 50, and 40 mg/mL paeonol concentrations, respectively. Additionally, more than 480 min was required to kill mites at 30, 20, and 10 mg/mL concentrations. In the negative controls, all mites remained active. The mortality rates in 8 h were 100%, 100%, 92.6%, 17.8%, 7.1%, and 7.1% for 60, 50, 40, 30, 20 and 10 mg/mL concentrations, respectively. No apparent difference in the mortality rate was noted between the 30, 20, and 10 mg/mL groups and the negative control group ($\chi^2 = 1.239$, $P > 0.05$; $\chi^2 = 0.440$, $P > 0.05$; $\chi^2 = 0.440$, $P > 0.05$, respectively). However, the 60, 50, and 40 mg/mL groups showed a significant difference when compared with the negative control group ($\chi^2 = 35.571$, $P < 0.05$; $\chi^2 = 36.512$, $P < 0.05$; $\chi^2 = 28.270$, $P < 0.05$, respectively). Thus, the

40 mg/mL dose was the lowest effective concentration of paeonol against *D. brevis* in 8 h.




Fig. 3. Effect of different concentrations of paeonol on the death times of human *Demodex* (A) and correlation analysis of the average death time of two species of *Demodex* (B).

The acaricidal effect of paeonol on the two species of human *Demodex*

Table I and Figure 3 suggest that the average death time for *D. brevis* (98.6, 123.4, and 181 mins) was less than that of *D. folliculorum* (118.2, 142.2, and 240 min) in 60, 50, and 40 mg/mL concentration groups, respectively. Thus, an apparent difference was noted between *D. brevis* and *D. folliculorum* in the average death time at the same concentration groups. *D. folliculorum* (81, 72, 145, 365, 422, and 436 mins) showed a longer survival time than *D. brevis* (59, 63, 96, 322, 391, and 391) at the same concentration groups of paeonol. The death rates of *D. folliculorum* in 8 h were 100%, 100%, 90.3%, 6.0%, 3.1%, and 3.3% for 60, 50, 40, 30, 20, and 10 mg/mL paeonol concentrations, respectively; the death rates of *D. brevis* were 100%, 100%, 92.6%, 17.8%, 7.1%, and 7.1% for 60, 50, 40, 30, 20, and 10 mg/mL paeonol concentrations, respectively. Additionally, paeonol could kill more *D. brevis*

than *D. folliculorum*, though higher drug concentrations resulted in faster death for both mite species. The average death time decreased with the increasing concentration of paeonol for both mite species. Thus, paeonol exhibited superior acaricidal activity against *D. brevis* than against *D. folliculorum* in terms of average death time, survival time, and mortality rate in 8 h.

DISCUSSION

Paeonol, with better therapeutic effects on pigmentation, psoriasis, eczema, and itchy skin, can bind to tyrosinase and inhibit the activity of enzymes to achieve whitening function (Wang *et al.*, 2020; Min *et al.*, 2023). Holmes (2013) found that bacteria in *Demodex* could be involved in the development of rosacea and other facial dermatoses (Lazaridou *et al.*, 2011; Jarmuda *et al.*, 2012). Other studies have shown that symptoms were relieved after antibacterial treatment (Hsu *et al.*, 2009; Martinez-Diaz *et al.*, 2012; Guerrero-González *et al.*, 2014). Zhao (2017) indicated that *Staphylococcus* and *Sphingomonas* might be associated with the development of facial dermatoses by the denaturing gradient gel electrophoresis (DGGE) technique (Zhao *et al.*, 2017). Zeng *et al.* (2022) demonstrated that paeonol exhibits an antibacterial effect on *Staphylococcus aureus* (Zeng *et al.*, 2022). Thus, we can reasonably expand the real-world applications of paeonol on skin care products. Heczko *et al.* (2023) reported that the mites were considered dead when all the rotary movements of *Demodex* appendages ceased (Fig. 2G, H). Clanner-Engelshofen (2022) proposed that the immotile mites with symmetrically spaced legs were potentially dead owing to muscular flaccidity (Clanner-Engelshofen *et al.*, 2020). In this study, we observed the body pincers and claw for 1 min and then stimulated the mite with an anatomical needle; if it remained motionless, we considered the mite to be dead (Clanner-Engelshofen *et al.*, 2020). We also discovered some coelom and/or internal hydroskeleton in live *D. folliculorum* (Fig. 2I). The solid hydroskeleton may help the body to move by decreasing the contact surface to reduce friction when walking (Clanner-Engelshofen *et al.*, 2020).

Although the 30, 20, and 10 mg/mL groups showed no significant difference with the negative control group in terms of the mortality rate in 8 h of treatment, the vitalities of mites were notably different at 8 h. The majority of mites exposed to the 30 mg/mL concentration of paeonol did not die but were carotic; only one leg moved with stimulation at the final stage of the 8-h in this research. However, the mites in the 20 or 10 mg/mL group actively moved even without needle stimulation at the end of 8 h. Therefore, a significant difference was noted between the 30 mg/

mL group and the negative control group when the drug application time was prolonged. A favorable correlation was noted between time and concentration and the total number of deceased human *Demodex* mites. The number of dead mites in 8 h increased as paeonol concentration increased, and the mortality rate reduced as paeonol concentration decreased. This finding was comparable to previous reports (Zhao *et al.*, 2006; Du *et al.*, 2021).

CONCLUSION

In this study, paeonol had a distinct killing effect on two types of human *Demodex* mites (*D. folliculorum* and *D. brevis*). The movements were delayed, podosoma and gnathosoma expansion occurred, even shortening arose in the opisthosoma, and the bodies became slightly more transparent than primal form when death. The killing time was lengthened with decreasing concentration, thus showing an evident dependence on concentration. According to the experimental findings, paeonol had a minimum effective dose of 40 mg/ml for killing both *D. folliculorum* and *D. brevis*, with paeonol having a higher killing effect on *D. brevis* than *D. folliculorum* during the 8-h period.

DECLARATIONS

Acknowledgment

The authors are grateful to the *Demodex*-positive patients for providing a large number of mites to carry out the work.

Funding

This work was supported by grants from the Academic Aid Program for Top-notch Talents in Provincial Universities (gxbjZD2020071), Wuhu Key Research and Development Program of China (2021yf391) and Young and Middle-aged Research Fund of Wannan Medical College (WK202216).

IRB approval

The study was approved by Wannan Medical college (Wuhu, China), and all participants signed a written informed consent forms.

Data availability statement

The authors confirm that the data supporting the findings of this study are available within the article.

Generative AI and AI-assisted technology statement

The authors have declared that no generative AI or AI-assisted technologies were used to create this manuscript.

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Adki, K.M. and Kulkarni, Y.A., 2020. Chemistry, pharmacokinetics, pharmacology and recent novel drug delivery systems of paeonol. *J. Life Sci.*, **250**: 18. <https://doi.org/10.1016/j.lfs.2020.117544>
- Chudzicka-Strugala, I., Golebiewska, I., Brudecki, G., Elamin, W. and Zwodziazak, B., 2023. Demodicosis in different age groups and alternative treatment options. A review. *J. clin. Med.*, **12**: 23. <https://doi.org/10.3390/jcm12041649>
- Clanner-Engelshofen, B.M., French, L.E. and Reinholz, M., 2020. Methods for extraction and *ex-vivo* experimentation with the most complex human commensal, *Demodex* spp. *J. exp. appl. Acarol.*, **80**: 59-70. <https://doi.org/10.1007/s10493-019-00450-9>
- Clanner-Engelshofen, B.M., Ständer, L.M., Steegmuller, T., Kämmerer, T., Frommherz, L.H., Stadler, P.C., Gurtler, A. and Reinholz, M., 2022. First *ex vivo* cultivation of human *Demodex* mites and evaluation of different drugs on mite proliferation. *J. Eur. Acad. Dermatol. Venereol.*, **36**: 2499-2503. <https://doi.org/10.1111/jdv.18468>
- Ding, M.G., Shi, R., Fu, F., Li, M., De, D.M., Du, Y.Y. and Li, Z.F., 2023. Paeonol protects against doxorubicin-induced cardiotoxicity by promoting Mfn2-mediated mitochondrial fusion through activating the PKC ϵ -Stat3 pathway. *J. Adv. Res.*, **47**: 151-162. <https://doi.org/10.1016/j.jare.2022.07.002>
- Du, J.J., Gao, R. and Zhao, J.H., 2021. The effect of volatile oil from chinese mugwort leaf on human demodecid mites *in vitro*. *J. Acta Parasitol.*, **66**: 615-622. <https://doi.org/10.1007/s11686-020-00314-y>
- Elston, C.A. and Elston, D.M., 2014. *Demodex* mites. *J. clin. Dermatol.*, **32**: 739-743. <https://doi.org/10.1016/j.clindermatol.2014.02.012>
- Elston, D.M., 2010. *Demodex* mites: Facts and controversies. *J. clin. Dermatol.*, **28**: 502-504. <https://doi.org/10.1016/j.clindermatol.2010.03.006>
- Foley, R., Kelly, P., Gatault, S. and Powell, F., 2021. *Demodex*: A skin resident in man and his best friend. *J. Eur. Acad. Dermatol. Venereol.*, **35**: 62-72. <https://doi.org/10.1111/jdv.16461>
- Forton, F.M. and De Maertelaer, V., 2021. Which factors influence *Demodex* proliferation? A retrospective pilot study highlighting a possible role of subtle

- immune variations and sebaceous gland status. *J. J. Dermatol.*, **48**: 1210-1220. <https://doi.org/10.1111/1346-8138.15910>
- Guerrero-González, G.A., Herz-Ruelas, M.E., Gómez-Flores, M. and Ocampo-Candiani, J., 2014. Crusted demodicosis in an immunocompetent pediatric patient. *J. Case Rep. Dermatol. Med.*, **2014**: 458046. <https://doi.org/10.1155/2014/458046>
- Gyawali, A., Krol, S. and Kang, Y.S., 2019. Involvement of a novel organic cation transporter in paeonol transport across the blood-brain barrier. *J. Biol. Ther.*, **27**: 290-301. <https://doi.org/10.4062/biomolther.2019.007>
- Heczko, J., Schell, C., Pansick, A., Stein, R. and Perry, H.D., 2023. Evaluation of a novel treatment, selenium disulfide, in killing *Demodex folliculorum* in vitro. *J. Can. J. Ophthalmol.*, **58**: 408-412. <https://doi.org/10.1016/j.jcjo.2022.04.012>
- Holmes, A.D., 2013. Potential role of microorganisms in the pathogenesis of rosacea. *J. Am. Acad. Dermatol.*, **69**: 1025-1032. <https://doi.org/10.1016/j.jaad.2013.08.006>
- Hsu, C.K., Hsu, M.M.L. and Lee, J.Y.Y., 2009. Demodicosis: A clinicopathological study. *J. Am. Acad. Dermatol.*, **60**: 453-462. <https://doi.org/10.1016/j.jaad.2008.10.058>
- Jacob, S., VanDaele, M.A. and Brown, J.N., 2019. Treatment of demodex-associated inflammatory skin conditions: A systematic review. *J. Dermatol. Ther.*, **32**: 7. <https://doi.org/10.1111/dth.13103>
- Jarmuda, S., O'Reilly, N., Żaba, R., Jakubowicz, O., Szkaradkiewicz, A. and Kavanagh, K., 2012. Potential role of *Demodex* mites and bacteria in the induction of rosacea. *J. med. Microbiol.*, **61**: 1504-1510. <https://doi.org/10.1099/jmm.0.048090-0>
- Kim, H.K., Tak, J.H. and Ahn, Y.J., 2004. Acaricidal activity of *Paeonia suffruticosa* root bark-derived compounds against *Dermatophagoides farinae* and *Dermatophagoides pteronyssinus* (Acari: Pyroglyphidae). *J. Agric. Fd. Chem.*, **52**: 7857-7861. <https://doi.org/10.1021/jf048708a>
- Lazaridou, E., Giannopoulou, C., Fotiadou, C., Vakirlis, E., Trigoni, A. and Ioannides, D., 2011. The potential role of microorganisms in the development of rosacea. *J. JD Tsch. Dermatol. Ges.*, **9**: 21-25. <https://doi.org/10.1111/j.1610-0387.2010.07513.x>
- Li, Q., Zhao, Y., Zhu, X.M. and Xie, Y.L., 2021. Antifungal efficacy of paeonol on *Aspergillus flavus* and its mode of action on cell walls and cell membranes. *J. LWT Fd. Sci. Technol.*, **149**: 9. <https://doi.org/10.1016/j.lwt.2021.111985>
- Lv, Q.H., Li, S.F., Wei, H.L., Wen, Z.M., Wang, Y.L., Tang, T.Z., Wang, J.F., Xia, L.N. and Deng, X.M., 2020. Identification of the natural product paeonol derived from peony bark as an inhibitor of the *Salmonella enterica* serovar *Typhimurium* type III secretion system. *J. appl. Microbiol. Biotechnol.*, **104**: 1673-1682. <https://doi.org/10.1007/s00253-019-10290-7>
- Martinez-Diaz, G.J., Clark, K.M., Vasquez, J.G. and English, J.C., 2012. Facial erythematous annular plaques: a case of annular *Demodex* facial dermatitis? *J. Am. Acad. Dermatol.*, **67**: e268-e269. <https://doi.org/10.1016/j.jaad.2012.05.015>
- Min, X.F., Lu, L., Xu, X.T., Wen, Y. and Zheng, X., 2023. Investigation on the inhibition mechanism and binding behavior of paeonol to tyrosinase and its anti-browning property by multi-spectroscopic and molecular docking methods. *J. Int. J. Biol. Macromol.*, **253**: 9. <https://doi.org/10.1016/j.ijbiomac.2023.126962>
- Norn, M.S.J., 1982. Incidence of *Demodex folliculorum* on skin of lids and nose. *J. Acta Ophthalmol.*, **60**: 575-583. <https://doi.org/10.1111/j.1755-3768.1982.tb00603.x>
- Paichitrojjana, A., 2022. Demodex: The worst enemies are the ones that used to be friends. *J. Dermatol. Rep.*, **14**: 9339. <https://doi.org/10.4081/dr.2022.9339>
- Palopoli, M.F., Fergus, D.J., Minot, S., Pei, D.T., Simison, W.B., Fernandez-Silva, I., Thoemmes, M.S., Dunn, R.R. and Trautwein, M., 2015. Global divergence of the human follicle mite *Demodex folliculorum*: Persistent associations between host ancestry and mite lineages. *J. Proc. natl. Acad. Sci. U.S.A.*, **112**: 15958-15963. <https://doi.org/10.1073/pnas.1512609112>
- Smith, G., Manzano-Marín, A., Reyes-Prieto, M., Antunes, C.S.R., Ashworth, V., Goselle, O.N., Jan, A.A.A., Moya, A., Latorre, A., Perotti, M.A. and Braig, H.R., 2022. Human follicular mites: Ectoparasites becoming symbionts. *J. mol. Biol. Evol.*, **39**: msac125. <https://doi.org/10.1093/molbev/msac125>
- Tak, J.H., Kim, H.K., Lee, S.H. and Ahn, Y.J., 2006. Acaricidal activities of paeonol and benzoic acid from *Paeonia suffruticosa* root bark and monoterpenoids against *Tyrophagus putrescentiae* (Acari: Acaridae). *J. Pest Manage. Sci.*, **62**: 551-557. <https://doi.org/10.1002/ps.1212>
- Tang, H.Q., Yang, D., Zhu, L., Shi, F., Ye, G., Guo, H.R., Deng, H.D., Zhao, L., Xu, Z.W. and Li, Y.L., 2022. Paeonol interferes with quorum-sensing in *Pseudomonas aeruginosa* and modulates

- inflammatory responses *in vitro* and *in vivo*. *J. Front. Immunol.*, **13**: 14. <https://doi.org/10.3389/fimmu.2022.896874>
- Wang, J.L., Wu, G.Y., Chu, H.P., Wu, Z.Y. and Sun, J.Y., 2020. Paeonol derivatives and pharmacological activities: A review of recent progress. *J. Mini Rev. Med. Chem.*, **20**: 466-482. <https://doi.org/10.2174/1389557519666191015204223>
- Wei, F., Li, L., Kong, Y., Yan, X.F., Varghese, K.J., Zhang, S., Jiang, J., Chai, B. and Chen, H.X., 2023. Evidence for the clinical association between *Demodex* and rosacea: A review. *J. Dermatology.*, **240**: 95-102. <https://doi.org/10.1159/000534245>
- Xu, H., Zhang, N. and Casida, J.E., 2003. Insecticides in Chinese medicinal plants: Survey leading to jacaranone, a neurotoxicant and glutathione-reactive quinol. *J. Agric. Fd. Chem.*, **51**: 2544-2547. <https://doi.org/10.1021/jf021164x>
- Zeng, Q., Fu, Y.T., Yang, M., Wang, T., Wang, Y., Lv, S.H. and Qian, W.D., 2022. Effect of paeonol against bacterial growth, biofilm formation and dispersal of *Staphylococcus aureus* and *Listeria monocytogenes* *in vitro*. *J. Biofouling.*, **38**: 173-185. <https://doi.org/10.1080/08927014.2022.2045014>
- Zhang, L., Li, D.C. and Lin, L.F., 2019. Paeonol: Pharmacological effects and mechanisms of action. *J. Int. Immunopharmacol.*, **72**: 413-421. <https://doi.org/10.1016/j.intimp.2019.04.033>
- Zhao, Y., Feng, L. and Shi, J., 2006. Studies on the experiment of killing *Demodex folliculorum* *in vitro* with crude extract of *Folium artemisiae argyi*. *J. Chin. J. Vector Biol. Contr.*, **17**: 209.
- Zhao, Y.E., Yang, F., Wang, R.L., Niu, D.L., Mu, X., Yang, R. and Hu, L., 2017. Association study of *Demodex* bacteria and facial dermatoses based on DGGE technique. *J. Parasitol. Res.*, **116**: 945-951. <https://doi.org/10.1007/s00436-016-5370-1>
- Zou, M.H., Xue, Q.Q., Guo, J.J., Teng, Q., Zhao, J.H. and Li, Y.Y., 2023a. Transcriptomics and proteomic-based study of the acaricidal mechanism of paeonol against *Aleuroglyphus ovatus*. *J. Int. J. Acarol.*, **11**. <https://doi.org/10.1080/01647954.2023.2268621>
- Zou, M.H., Xue, Q.Q., Teng, Q., Zhang, Q.Q., Liu, T., Li, Y.Y. and Zhao, J.H., 2023b. Acaricidal activities of paeonol from *Moutan cortex*, dried bark of *Paeonia x suffruticosa*, against the grain pest mite *Aleuroglyphus ovatus* (Acari: Acaridae). *J. exp. appl. Acarol.*, **91**: 615-628. <https://doi.org/10.1007/s10493-023-00861-9>