



Comparative Efficacy of Different Insecticides against *Bemisia tabaci* (Gennadius) under Controlled Conditions

Muddassir Ashiq, Shoaib Freed*, Javeria Abbas and Mudasar Raza

Department of Entomology, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan, Pakistan

ABSTRACT

Bemisia tabaci is the critical polyphagous pest of different crops and insecticides belonging to different groups are being used for its control. The present investigation was performed to estimate the efficacy of conventional and new synthetic insecticides along with their biochemical analysis on *B. tabaci*. The data affirmed that utmost nymphal mortality 89.80% was detected in the buprofezin and 81.46% mortality in thiamethoxam at highest concentration. Results of adult bioassay revealed maximum mortality i.e., 84.33% in the bifenthrin followed by 66.66%, 42.34%, 36.84% mortality in chlorfenapyr, profenofos and imidacloprid, respectively. Detoxification enzyme analysis showed maximum activities of acetylcholinesterase, esterases, glutathione S-transferase, acid and alkaline phosphatases in the highest concentration of insecticides and minimum activities were observed in the control groups. Findings of the current research showed that the rotational use of insecticides may be highly effective for *B. tabaci* and can also minimize risk of resistance development.

Article Information

Received 11 October 2024
Revised 10 October 2025
Accepted 28 October 2025
Available online 11 March 2026
(early access)
Published 18 June 2026

Authors' Contribution

MA and SF conceived and designed the research. MA and MR conducted experiments and collected data. SF supervised the experiments. MA, JA, MR and SF analyzed the data and wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

Key words

White fly, Insecticides, Biochemical characterization, Resistance, IGR, Enzymes

INTRODUCTION

Bemisia tabaci (Gennadius) is most significant pest of Ornamentals and vegetables (Perring *et al.*, 2018). Its host range varies from ornamentals to solanaceous crops including tobacco, brinjal, tomato, cotton, potato and chilies (Kanakala and Ghanim, 2019; Smith and Krey, 2019; Kunjwal and Srivastava, 2018). It damages the plants by sucking cell sap from leaves (Wang *et al.*, 2017; Eslamizadeh *et al.*, 2015; Mascarini *et al.*, 2013) and also excretes honeydew which causes the formation of mold and it inhibits the photosynthesis (Gao *et al.*, 2017; Xie *et al.*, 2014). In addition, nymphs inject enzymes that alter the plant physiology resulting in an unequal ripening of fruits and underdeveloped color (Legg *et al.*, 2014). It

also acts as vector of pathogens including torradoviruses, begomoviruses, ipomoviruses, carlaviruses and criniviruses (Cuthbertson, 2013; Stansly and Natwick, 2010; Kumarasinghe *et al.*, 2009). Out of these begomoviruses are well known for causing the losses in many crops (Gangwar and Charu, 2018).

In Pakistan, *B. tabaci* has developed resistance against carbamates, pyrethroids and organophosphates due to the excessive usage (Ahmad *et al.*, 2002, 2010; Ali, 2011). Neonicotinoids have novel mode as related to the central nervous system of insects (Stenersen, 2004). In the present cropping systems, neonicotinoids are being used against sucking (plant and leaf hoppers, aphid and whiteflies) and certain chewing (coleopterans and lepidopterans) pests (Jeschke *et al.*, 2011).

Alternatively, insect growth regulators (IGR's) have shown the potential for controlling *B. tabaci* under green house and field crops (Ishaaya *et al.*, 1988). IGR's act by interrupting the metamorphosis and development of insects and also disrupt their expression of activity, that's why these insecticides act in slow pace than other insecticides (Sial and Brunner, 2010; Horowitz *et al.*, 2005). Synthetic pyrethroids act on the voltage-gated calcium, sodium and chloride ducts and GABA-gated chloride channels

* Corresponding author: sfareed@bzu.edu.pk
0030-9923/2026/0004-1863 \$ 9.00/0



Copyright 2026 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

(Forshaw and Ray, 1990; Song and Narahashi, 1996). Bifenthrin being a 3rd generation pyrethroid is effective for managing the pests of cotton (Balakrishnan *et al.*, 2009), fruits (Reddy and Rao, 2002), rice (Veeravel and Ravivarman, 2010) and vegetables (Gupta *et al.*, 2009).

One of the causes of insecticide resistance is the physiological changes leading to an increased level of enzymes involved in the detoxification process (Heckel, 2012; Vontas *et al.*, 2002). Furthermore, the detoxification process can lead to cross-resistance to certain other insecticides (Zhang *et al.*, 2016; Mitchell *et al.*, 2012; Qian *et al.*, 2008). Considering the importance of *B. tabaci* as a potential economic pest, this study was planned to evaluate the effectiveness of insecticides on *B. tabaci* and the involvement of detoxification enzymes in the resistance development.

MATERIALS AND METHODS

Bemisia tabaci

Bemisia tabaci was collected with the help of manual aspirator from cotton fields of Bahauddin Zakariya University. Adults were reared on cotton plants in glasshouse without any exposure of insecticides under optimum conditions ($T = 25 \pm 2^\circ\text{C}$ and $RH = 65 \pm 5\%$) until 5 generations. Plants were monitored for egg laying after 48-72 h, and plants containing eggs were transferred into rearing cages to attain uniform generation.

Insecticides

Commercial formulations of imidacloprid 20 SL Confidor®, Syngenta, chlorfenapyr 360 SC Squadron® FMC, profenofos 50 EC Curacron®, Syngenta and bifenthrin 10EC Talstar®, FMC were used against adults, while thiamethoxam 25 WG Actara®, Syngenta, buprofezin 25 WP Buprofezin®, Syngenta were used against nymphs. All concentrations were made by serial dilution method.

Insecticides bioassay for adults

For bioassay, leaf dip method as reported by Luo *et al.* (2010) was used with slight variations to assess the effect of lethal and sublethal concentration (LC_{50} and LC_{30}). Briefly 2 mL out of 1.5% agar solution was taken into petri dishes of 1×3.5 cm and placed at a cool place. Petri dishes were modified by making a round hole in the cover lid with a cut of 2.8 cm in diameter for infusion of air into the petri dishes, mesh size 100 was used to make insect proof cover for the opening. To avoid large leaf veins, hole of 3.5 cm circumference was made. Leaves discs were carefully submerged in the dilutions while ddH₂O water was used to treat control insects. Leaves were left for drying with their

abaxial sides facing up and then put in petri dishes to get attached with agar having no space between them.

Bemisia tabaci adults were placed in the petri dishes by making the adults immobilized by placing them at -20°C for 10 s. 15 individuals per petri dishes were used for experiment. Bioassay was performed under completely randomized design (CRD) design having 5 treatments and 3 replications. Data was recorded by counting the dead insects and those not moving normally were also counted as dead.

Nymphal bioassay

Methodology of Zheng *et al.* (2017) was followed for nymphal bioassay. Cotton plants at two true leaves stage were grown in the glasshouse and then transferred into cages. Fifty *B. tabaci* adults were shifted in the cages for egg-laying. After 24 h, egg-laying was observed under microscope and adults were removed. Number of nymphs were counted using magnifying lens when they reached up to 2nd nymphal instar. Infested leaves were dipped into the dilutions of the insecticides for 20 s and mortality was assessed after seven days when nymphs developed into pupae. Those nymphs were considered as dead that could not transform into pupae. Experiment contained five treatments including one control and three replicates.

Determination enzymatic activity

Samples for biochemical analysis of *B. tabaci* were collected by following Malik *et al.* (2019). Five treated adults were taken into Eppendorf tubes (1.5 mL) and 100 μL of 0.15 M NaCl was added for crushing and volume was made 500 μL . Samples were spun for 10 min at 10,000 rpm and supernatant was used for determination of enzymatic activity glutathione (GST), esterases (EST), acetylcholinesterase (AChE), alkaline phosphatases (ALP) and acid phosphatase (ACP).

Habig *et al.* (1974) was followed for estimation of GST activity. Methodology of Damayanthi and Karunaratne (2005) was used for determining the EST activity. For determining the AChE activity, methodology of Ellman *et al.* (1961) was followed. The procedure of Otto *et al.* (1946) was followed to check the activities of ALP and ACP. Tris-HCl buffer 0.05M was used with 0.23mM disodium 4-nitrophenyl phosphate (PNPP), while for ACP 0.05M citrate buffer was used. Samples were incubated for 2 h at room temperature. After incubation, 425 μL 0.05M NaOH was mixed in the samples for coloration. Activities were observed at λ 410 nm.

Data analysis

The software POLO-PC (LeOra, 2003) was

used for establishing both lethal (LC_{50}) and sub-lethal concentrations (LC_{30}). Mortality data was then analyzed using the software Statistix 8.1, and the Tukey's test was used to differentiate between the means of different treatment. Graphs describing enzymatic activities were created using Microsoft Excel 2010 and Prism Graph Pad (version 9.0.0). Mean values with standard errors were determined using the bootstrap methods (Meyer *et al.*, 1986). All experiment were repeated twice.

RESULTS

Mortality of B. tabaci after exposure to insecticides

Toxicity of insecticides was evaluated using leaf dip bioassay. Mortality data for insecticides efficacy on adults of *B. tabaci* was observed 2nd day post treatment which depicted highest mortality in bifenthrin i.e., 84.33 % followed by chlorfenapyr, profenofos and imidacloprid 66.66, 42.34 and 36.84%, respectively compared to the untreated groups (Table I). The data depicted varying LC_{50} , LC_{30} and slopes values for different insecticides used against adult *B. tabaci* (Table II).

Mortality data for insecticides efficacy on *B. tabaci* nymphs was observed 7th day post treatment until nymphs changed to pupae. Results showed that highest mortality of nymphs was caused by buprofezin (89.80 %). Thiamethoxam also demonstrated notable mortality of 81.46% as compared to the untreated groups (Table 1). Similarly, buprofezin and thiamethoxam treatments on *B. tabaci* nymphs showed LC_{50} and slope 1.039, LC_{50} , LC_{30} and slope 0.796 values, respectively (Table II).

Enzymatic activity of B. tabaci after insecticides exposure

In bifenthrin treatment, highest AChE activity 23.53 $\mu\text{mol/min mg protein}$ was observed in the maximum concentration treatment followed by GST, EST, ALP activities 18.53, 15.53, 7.17 $\mu\text{mol/min mg protein}$, while least activity of ACP i.e., 5.39 was recorded in the similar treatment (Fig. 1). In chlorfenapyr treatment highest activity of GST i.e., 29.20 was found in the maximum concentration followed by EST 28.53, AChE 22.54, ALP 6.05 $\mu\text{mol/min mg protein}$ and minimum activity of ACP 5.19 was found in highest concentration (Fig. 2).

Table I. Percent mortality of different stages of *B. tabaci* after exposure to insecticides.

Concentration (ppm)	Adult				Concentration (ppm)	Nymph	
	Bifenthrin (n=270)	Chlorfenapyr (n=270)	Profenofos (n=270)	Imidacloprid (n=270)		Buprofezin (n=270)	Thiamethoxam (n=270)
100	84.33 $\pm$ 0.64 ^a	66.66 $\pm$ 1.70 ^a	42.34 $\pm$ 1.11 ^a	36.84 $\pm$ 0.69 ^a	20	89.80 $\pm$ 1.82 ^a	81.46 $\pm$ 1.11 ^a
75	46.66 $\pm$ 0.74 ^{ab}	56.66 $\pm$ 0.74 ^{ab}	35.66 $\pm$ 1.29 ^{ab}	27.33 $\pm$ 0.74 ^a	10	74.87 $\pm$ 1.02 ^a	70.00 $\pm$ 1.29 ^{ab}
50	33.33 $\pm$ 0.74 ^b	40.00 $\pm$ 1.82 ^{bc}	25.32 $\pm$ 1.29 ^{bc}	20.00 $\pm$ 1.58 ^{ab}	5	69.33 $\pm$ 0.74 ^{ab}	60.00 $\pm$ 1.29 ^{ab}
25	20.00 $\pm$ 1.82 ^c	30.00 $\pm$ 1.82 ^c	20.00 $\pm$ 1.58 ^{ca}	16.66 $\pm$ 1.05 ^b	1	53.33 $\pm$ 0.91 ^{bc}	50.00 $\pm$ 1.58 ^{bc}
12.50	13.33 $\pm$ 1.29 ^{cd}	23.33 $\pm$ 1.29 ^{cd}	13.33 $\pm$ 1.05 ^a	10.00 $\pm$ 1.58 ^b	0.5	36.66 $\pm$ 1.05 ^{cd}	33.33 $\pm$ 1.05 ^c
Control	6.66 $\pm$ 1.82 ^d	3.33 $\pm$ 1.82 ^d	6.66 $\pm$ 1.82 ^c	3.33 $\pm$ 1.82 ^c	Control	7.66 $\pm$ 1.82 ^d	6.66 $\pm$ 0.82 ^d

Table II. LC_{50} and LC_{30} of different insecticides treated against laboratory population of *B. tabaci*.

Stage	Insecticides	LC_{50}	LC_{30}	Slope $\pm$ SE	χ^2	df	P
Adult	Profenofos	0.742-3.186	0.039-1.004	0.681 $\pm$ 0.179	2.344	3	0.781
	Bifenthrin	1.636-9.009	0.843-1.554	0.546 $\pm$ 0.175	0.171	3	0.057
	Imidacloprid	1.559-3.893	0.302-1.258	0.973 $\pm$ 0.186	0.655	3	0.218
	Chlorfenapyr	3.113-14.159	3.622-4.202	0.636 $\pm$ 0.178	1.418	3	0.473
Nymph	Buprofezin	1.076-2.688	0.234-1.214	1.039 $\pm$ 0.190	5.385	3	0.185
	Thiamethoxam	1.047-3.387	0.112-1.203	0.796 $\pm$ 0.181	0.253	3	0.084

* S.E, Standard Error; χ^2 , Chi-square value; df, Degree of freedom; P, p-value.




Fig. 1. Activities of AChE, EST, GST, ALP and ACP in *B. tabaci* adults treated with bifenthrin. Bars with same letters represent non-significant difference ($P > 0.05$).




Fig. 2. Activities of AChE, EST, GST, ALP and ACP in *B. tabaci* adults treated with chlorfenapyr. Bars with same letters represent non-significant difference ($P > 0.05$).

In the imidacloprid treatment maximum activity of EST 31.20 was recorded in highest concentration treatment followed by 25.53 in GST, 15.53 in AChE, 4.41 in ACP while minimum activity of ALP 3.97 was observed in highest concentration treatment (Fig. 3). In the profenofos treatment, maximum activity of GST 29.25 was found in the highest concentration treatment followed by 26.75, 20.15, 5.17 $\mu\text{mol}/\text{min mg}$ protein of EST, AChE, ALP while minimum activity was observed in the ACP (Fig. 4).




Fig. 3. Activities of AChE, EST, GST, ALP and ACP in *B. tabaci* adults treated with imidacloprid. Bars with same letters represent non-significant difference ($P > 0.05$).

Enzymatic activity in nymphs of *B. tabaci*

In buprofezin treatment maximum activity of GST i.e., 25.53 $\mu\text{mol}/\text{min mg}$ protein followed by AChE 23.20, EST 18.87, ALP 9.72 $\mu\text{mol}/\text{min mg}$ protein, and minimum level of ACP 9.05 was recorded in highest concentration treatment (Fig. 5). Similarly, in thiamethoxam treatment, maximum activity of AChE i.e., 25.47 was found in highest concentration treatment followed by 20.20, 18.20, 5.73 in GST, EST and ACP while minimum activity of ALP 4.17 was at similar concentration, respectively (Fig. 6).




Fig. 4. Activities of AChE, EST, GST, ALP and ACP in *B. tabaci* adults treated with profenofos. Bars with same letters represent non-significant difference ($P > 0.05$).




Fig. 6. Activities of AChE, EST, GST, ALP and ACP in *B. tabaci* nymphs treated with thiamethoxam. Bars with same letters represent non-significant difference ($P > 0.05$).




Fig. 5. Activities of AChE, EST, GST, ALP and ACP in *B. tabaci* nymphs treated with buprofezin. Bars with same letters represent non-significant difference ($P > 0.05$).

DISCUSSION

Bemisia tabaci is a versatile pest found in plains of Pakistan all over the year. During summer, cotton, okra, brinjal, and tomatoes flourish in a hotter and drier climate. Small farmers commonly sow these crops adjacently, and whiteflies continue to migrate between them. The occurrence of *B. tabaci* resistance relies on the regularity of pesticide uses instead of whitefly populations on specific crops. Cotton, brinjal and okra are treated with insecticide four times, while tomato is sprayed only two times. In Pakistan, it is common to control the infestation of sucking pests on vegetables and ornamental plants by using new chemicals (Basit *et al.*, 2013). In the 1990s, Pakistan permitted the use of novel insecticides in response to a notable rise in resistance to insecticides in the country (Ahmad *et al.*, 2002, 2010). For *B. tabaci*, pesticides with novel modes of action proved to be highly effective and were extensively used. In the year 2000, the vulnerability of *B. tabaci* to common insecticides was regained due to increased use of novel chemical insecticides and the decreased reliance on conventional insecticides (Ahmad and Khan, 2017). This study presents the insecticidal effectiveness of seven different insecticides classified as

organophosphates, pyrethroids, neonicotinoids, and IGRs. Maximum mortality was observed by buprofezin 89.80%, which is in agreement with Gogi *et al.* (2021) who deliberated the efficacy of insecticides against whitefly and *Encarsia formosa* and stated buprofezin to cause 84.4% nymphal mortality on Bt-cotton. Neonicotinoids are the major insecticides used for the management of whiteflies (Yang *et al.*, 2016). Findings of this study indicated that thiamethoxam showed 81.46% mortality of nymphs which is in line with Chen *et al.* (2018) who reported toxicity of thiamethoxm to *B. tabaci* nymphs.

Imidacloprid provides effective control of adults of *B. tabaci* and in the current research 84.33% mortality was recorded which is partially consistent with the outcomes of Babar *et al.* (2013) and Ghosh (2020) who reported 76.0 and 80.1% mortalities of adult *B. tabaci*. Whiteflies in cotton have been difficult to eradicate using standard pesticides instead of pyrethroids (Dennehy and Williams III, 1997). The tolerance of whitefly may be due to the decreased insecticide permeation, the high insecticide dissolution, and lack of sensitivity of the insecticides' target location (Guedes, 1999; Yu, 1988). Low mortalities 36.84 and 42.34%, were caused by bifenthrin and profenofos treatment which agree with the findings of Bacci *et al.* (2007) and Kady and Devine (2003) who reported inefficiency of these insecticides for controlling *B. tabaci*.

AChE, ALP, ACP, EST, and GST enzymes are responsible for neutralizing toxic metabolites in the insect body. These enzymes are essential for maintaining the normal functioning of insect physiology (Chen *et al.*, 2017; Zibace *et al.*, 2009). The metabolism of insects depends on enzymes that neutralize the impact of insecticides. The resistance in insect pests to these pesticides depends on the amplified activities of such enzymes (Gong *et al.*, 2021).

The study tested the impact of pesticide application on the activities of detoxification enzymes in *B. tabaci*. Increased GST and AChE activities were noted following pesticide treatment on *B. tabaci*, which aligns with the outcomes of Yang *et al.* (2016) and Bilal *et al.* (2017, 2018a, b) who detected a similar rise in AChE activity following exposure to pesticides. Similarly, elevated levels of EST, ALP, and ACP were detected in insects treated with insecticides, which correlates with Wang *et al.* (2002) and Jia *et al.* (2016) who described amplified enzymatic activities in *Aphis gossypii* and *Locusta migratoria*.

CONCLUSION

This study exposes the difference in vulnerability of *B. tabaci* to different insecticides under laboratory conditions and its biochemical characterization. Results propose that appropriate use of insecticides with rotation

can help to reduce *B. tabaci* population and reduce the risk of insecticides resistance.

DECLARATIONS

Funding

The study received no external funding.

Generative AI and AI-assisted technology statement

The authors declare that no generative AI and AI assisted technology was used in the creation of this manuscript.

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Ahmad, M., Arif, M.I., Ahmad, Z. and Denholm, I., 2002. Cotton whitefly (*Bemisia tabaci*) resistance to organophosphate and pyrethroid insecticides in Pakistan. *Pest Manage. Sci.*, **58**: 203-208. <https://doi.org/10.1002/ps.440>
- Ahmad, M., Arif, M.I. and Naveed, M., 2010. Dynamics of resistance to organophosphate and carbamate insecticides in the cotton whitefly *Bemisia tabaci* (Hemiptera: Aleyrodidae) from Pakistan. *J. Pest. Sci.*, **83**: 409-420. <https://doi.org/10.1007/s10340-010-0311-8>
- Ahmad, M. and Khan, R.A., 2017. Field-evolved resistance of *Bemisia tabaci* (Hemiptera: Aleyrodidae) to carbodiimide and neonicotinoids in Pakistan. *J. econ. Ent.*, **110**: 1235-1242. <https://doi.org/10.1093/jee/tox058>
- Ali, M., 2011. *Handbook for agriculture extension agents on pesticides standardized in the Punjab*. Agriculture Extension Wing, Agriculture Department, Government of the Punjab.
- Babar, T.K., Karar, H., Saleem, M., Ali, A., Ahmad, S. and Hameed, A., 2013. Comparative efficacy of various insecticides against whitefly, *Bemisia tabaci* (Genn.) adult (Homoptera: Aleyrodidae) on transgenic cotton variety Bt-886. *Pak. Entomol.*, **35**: 99-104.
- Bacci, L., Crespo, A.L., Galvan, T.L., Pereira, E.J., Picanço, M.C., Silva, G.A. and Chediak, M., 2007. Toxicity of insecticides to the sweetpotato whitefly (Hemiptera: Aleyrodidae) and its natural enemies. *Pest Manage. Sci.*, **63**: 699-706. <https://doi.org/10.1002/ps.1393>
- Balakrishnan, N., Kumar, B.V. and Sivasubramanian, P., 2009. Bioefficacy of bifenthrin 10 EC against

- sucking insects, bollworms and natural enemies in cotton. *Madras Agric. J.*, **96**: 225-229.
- Basit, M., Saeed, S., Saleem, M.A., Denholm, I. and Shah, M., 2013. Detection of resistance, cross-resistance, and stability of resistance to new chemistry insecticides in *Bemisia tabaci* (Homoptera: Aleyrodidae). *J. econ. Ent.*, **106**: 1414-1422. <https://doi.org/10.1603/EC12414>
- Bilal, M., Freed, S., Ashraf, M.Z. and Muhammad, S., 2017. Enhanced activities of acetylcholinesterase, acid and alkaline phosphatases in *Helicoverpa armigera* after exposure to entomopathogenic fungi. *Invertebr. Surviv. J.*, **14**: 464-476.
- Bilal, M., Freed, S., Ashraf, M.Z., Zaka, S.M. and Khan, M.B., 2018a. Activity of acetylcholinesterase and acid and alkaline phosphatases in different insecticide-treated *Helicoverpa armigera* (Hübner). *Environ. Sci. Pollut. Res.*, **25**: 22903-22910. <https://doi.org/10.1007/s11356-018-2394-3>
- Bilal, M., Freed, S., Muhammad, S., Ashraf, M.Z. and Khan, M.B., 2018b. Activity of glutathione S transferase and esterase enzymes in *Helicoverpa armigera* (Hübner) after exposure to entomopathogenic fungi. *Entomol. Res.*, **48**: 279-287. <https://doi.org/10.1111/1748-5967.12284>
- Chen, J., Wang, Z., Cao, L., Gong, Y., Hoffmann, A.A. and Wei, S., 2018. Toxicity of seven insecticides to different developmental stages of the whitefly *Bemisia tabaci* MED (Hemiptera: Aleyrodidae) in multiple field populations of China. *Ecotoxicology*, **27**: 742-751. <https://doi.org/10.1007/s10646-018-1956-y>
- Chen, X.D., Seo, M. and Stelinski, L.L., 2017. Behavioral and hormetic effects of the butenolide insecticide, flupyradifurone, on Asian citrus psyllid, *Diaphorina citri*. *Crop Prot.*, **98**: 102-107. <https://doi.org/10.1016/j.cropro.2017.03.017>
- Cuthbertson, A.G., 2013. Update on the status of *Bemisia tabaci* in the UK and the use of entomopathogenic fungi within eradication programmes. *Insects*, **4**: 198-205. <https://doi.org/10.3390/insects4020198>
- Damayanthi, B. and Karunaratne, S., 2005. Biochemical characterization of insecticide resistance in insect pests of vegetables and predatory ladybird beetles. *J. natl. Sci. Found. Sri Lanka*, **33**: 115-122. <https://doi.org/10.4038/jnsfsv.33i2.2341>
- Dennehy, T.J. and Williams III, L., 1997. Management of resistance in *Bemisia* in Arizona cotton. *Pestic. Sci.*, **51**: 398-406. [https://doi.org/10.1002/\(SICI\)1096-9063\(199711\)51:3<398::AID-PS655>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1096-9063(199711)51:3<398::AID-PS655>3.0.CO;2-C)
- Ellman, G.L., Courtney, K.D., Andres Jr, V. and Featherstone, R.M., 1961. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.*, **7**: 88-95. [https://doi.org/10.1016/0006-2952\(61\)90145-9](https://doi.org/10.1016/0006-2952(61)90145-9)
- Eslamizadeh, R., Sajap, A.S.B., Omar, D.B., Azura, N. and Adam, B., 2015. Evaluation of different isolates of entomopathogenic fungus, *Paecilomyces fumosoroseus* (Deuteromycotina: Hyphomycetes) against *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Biol. Control Pl. Prot.*, **2**: 82-90.
- Forshaw, P. and Ray, D., 1990. A novel action of deltamethrin on membrane resistance in mammalian skeletal muscle and non-myelinated nerve fibres. *Neuropharmacology*, **29**: 75-81. [https://doi.org/10.1016/0028-3908\(90\)90086-7](https://doi.org/10.1016/0028-3908(90)90086-7)
- Gangwar, R. and Charu, G., 2018. Lifecycle, distribution, nature of damage and economic importance of whitefly, *Bemisia tabaci* (Gennadius). *Acta Sci. Agric.*, **2**: 36-39.
- Gao, T., Wang, Z., Huang, Y., Keyhani, N.O. and Huang, Z., 2017. Lack of resistance development in *Bemisia tabaci* to *Isaria fumosorosea* after multiple generations of selection. *Sci. Rep.*, **7**: 1-11. <https://doi.org/10.1038/srep42727>
- Ghosh, S.K., 2020. Environmentally sound approach for management of tomato whitefly (*Bemisia tabaci* Genn.). *J. Ent. Zool. Stud.*, **8**: 814-818. <https://doi.org/10.22271/j.ento.2020.v8.i6j.7926>
- Gogi, M.D., Syed, A.H., Atta, B., Sufyan, M., Arif, M.J., Arshad, M., Nawaz, A., Khan, M.A., Mukhtar, A. and Liburd, O.E., 2021. Efficacy of biorational insecticides against *Bemisia tabaci* (Genn.) and their selectivity for its parasitoid *Encarsia formosa* Gahan on Bt cotton. *Sci. Reports.*, **11**: 2101. <https://doi.org/10.1038/s41598-021-81585-x>
- Gong, P., Chen, D., Wang, C., Li, M., Li, X., Zhang, Y., Li, X. and Zhu, X., 2021. Susceptibility of four species of aphids in wheat to seven insecticides and its relationship to detoxifying enzymes. *Front. Physiol.*, **11**: 1852-1860. <https://doi.org/10.3389/fphys.2020.623612>
- Guedes, R., 1999. *Resistência de insetos a inseticidas*. Manejo integrado de doenças e pragas, pp. 101-107.
- Gupta, S., Sharma, R., Gupta, R., Sinha, S., Singh, R. and Gajbhiye, V., 2009. Persistence of new insecticides and their efficacy against insect pests of okra. *Bull. environ. Contam. Toxicol.*, **82**: 243-247. <https://doi.org/10.1007/s00128-008-9581-8>
- Habig, W.H., Pabst, M.J. and Jakoby, W.B., 1974. Glutathione S-transferases: The first enzymatic step in mercapturic acid formation. *J. biol. Chem.*, **249**: 7130-7139. <https://doi.org/10.1016/S0021->

- 9258(19)42083-8
- Heckel, D.G., 2012. Insecticide resistance after silent spring. *Science*, **337**: 1612-1614. <https://doi.org/10.1126/science.1226994>
- Horowitz, A.R., Kontsedalov, S., Khasdan, V. and Ishaaya, I., 2005. Biotypes B and Q of *Bemisia tabaci* and their relevance to neonicotinoid and pyriproxyfen resistance. *Arch. Insect Biochem. Physiol.*, **58**: 216-225. <https://doi.org/10.1002/arch.20044>
- Ishaaya, I., Mendelson, Z. and Melamed-Madjar, V., 1988. Effect of buprofezin on embryo genesis and progeny formation of sweet potato whitefly (Homoptera: Aleyrodidae). *J. econ. Ent.*, **81**: 781-784. <https://doi.org/10.1093/jee/81.3.781>
- Jeschke, P., Nauen, R., Schindler, M. and Elbert, A., 2011. Overview of the status and global strategy for neonicotinoids. *J. Agric. Fd. Chem.*, **59**: 2897-2908. <https://doi.org/10.1021/jf101303g>
- Jia, M., Cao, G., Li, Y., Tu, X., Wang, G., Nong, X., Whitman, D.W. and Zhang, Z., 2016. Biochemical basis of synergism between pathogenic fungus *Metarhizium anisopliae* and insecticide chlorantraniliprole in *Locusta migratoria* (Meyen). *Sci. Reports*, **6**: 1-15. <https://doi.org/10.1038/srep28424>
- Kady, H.E. and Devine, G.J., 2003. Insecticide resistance in Egyptian populations of the cotton whitefly, *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Pest Manage. Sci.*, **59**: 865-871. <https://doi.org/10.1002/ps.687>
- Kanakala, S. and Ghanim, M., 2019. Global genetic diversity and geographical distribution of *Bemisia tabaci* and its bacterial endosymbionts. *PLoS One*, **14**: e0213946. <https://doi.org/10.1371/journal.pone.0213946>
- Kumarasinghe, N.C., Salim, N. and Wijayarathne, W., 2009. Identification and biology of two whitefly species on cassava in Sri Lanka. *J. Pl. Prot. Res.*, **49**: 373-377. <https://doi.org/10.2478/v10045-009-0058-5>
- Kunjwal, N. and Srivastava, R.M., 2018. *Insect pests of vegetables*. Springer, Singapore. https://doi.org/10.1007/978-981-10-8687-8_7
- Legg, J.P., Shirima, R., Tajebe, L.S., Guastella, D., Boniface, S., Jeremiah, S., Nsami, E., Chikoti, P. and Rapisarda, C., 2014. Biology and management of *Bemisia* whitefly vectors of cassava virus pandemics in Africa. *Pest Manage. Sci.*, **70**: 1446-1453. <https://doi.org/10.1002/ps.3793>
- LeOra, S., 2003. *Poloplus, a user's guide to probit or logit analysis*. LeOra Software, Berkeley, CA.
- Luo, C., Jones, C., Devine, G., Zhang, F., Denholm, I. and Gorman, K., 2010. Insecticide resistance in *Bemisia tabaci* biotype Q (Hemiptera: Aleyrodidae) from China. *Crop Prot.*, **29**: 429-434. <https://doi.org/10.1016/j.cropro.2009.10.001>
- Malik, S.J., Freed, S., Ali, N., Ismail, H.M. and Naeem, A., 2019. Nitenpyram selection, resistance and biochemical characterization in dusky cotton bug, *Oxycarenus hyalinipennis* Costa (Hemiptera: Lygaeidae). *Crop Prot.*, **125**: 104-114. <https://doi.org/10.1016/j.cropro.2019.104904>
- Mascarin, G.M., Kobori, N.N., Quintela, E.D. and Delalibera Jr, I., 2013. The virulence of entomopathogenic fungi against *Bemisia tabaci* biotype B (Hemiptera: Aleyrodidae) and their conidial production using solid substrate fermentation. *Biol. Contr.*, **66**: 209-218. <https://doi.org/10.1016/j.biocontrol.2013.05.001>
- Meyer, J.S., Ingersoll, C.G., McDonald, L.L. and Boyce, M.S., 1986. Estimating uncertainty in population growth rates: jackknife vs. bootstrap techniques. *Ecology*, **67**: 1156-1166. <https://doi.org/10.2307/1938671>
- Mitchell, S.N., Stevenson, B.J., Müller, P., Wilding, C.S., Egyir-Yawson, A., Field, S.G., Hemingway, J., Paine, M.J., Ranson, H. and Donnelly, M.J., 2012. Identification and validation of a gene causing cross-resistance between insecticide classes in *Anopheles gambiae* from Ghana. *Proc. natl. Acad. Sci.*, **109**: 6147-6152. <https://doi.org/10.1073/pnas.1203452109>
- Otto, A., Oliver, H. and Jane, M., 1946. A method for the rapid determination of alkaline phosphatase with five cubic millimeters of serum. *J. biol. Chem.*, **164**: 321-329. [https://doi.org/10.1016/S0021-9258\(18\)43072-4](https://doi.org/10.1016/S0021-9258(18)43072-4)
- Perring, T.M., Stansly, P.A., Liu, T., Smith, H.A. and Andreason, S.A., 2018. *Whiteflies: Biology, ecology, and management, Sustainable management of arthropod pests of tomato*. Elsevier, pp. 73-110. <https://doi.org/10.1016/B978-0-12-802441-6.00004-8>
- Qian, L., Cao, G., Song, J., Yin, Q. and Han, Z., 2008. Biochemical mechanisms conferring cross-resistance between tebufenozide and abamectin in *Plutella xylostella*. *Pestic. Biochem. Physiol.*, **91**: 175-179. <https://doi.org/10.1016/j.pestbp.2008.03.011>
- Reddy, D.J. and Rao, B.N., 2002. Efficacy of selected insecticides against pests of grapevine. *Pestic. Res. J.*, **14**: 92-99.
- Sial, A.A. and Brunner, J.F., 2010. Lethal and sublethal

- effects of an insect growth regulator, pyriproxyfen, on obliquebanded leafroller (Lepidoptera: Tortricidae). *J. econ. Ent.*, **103**: 340-347. <https://doi.org/10.1603/EC09295>
- Smith, H.A. and Krey, K.L., 2019. Three release rates of *Dicyphus hesperus* (Hemiptera: Miridae) for management of *Bemisia tabaci* (Hemiptera: Aleyrodidae) on greenhouse tomato. *Insects*, **10**: 213. <https://doi.org/10.3390/insects10070213>
- Song, J. and Narahashi, T., 1996a. Differential effects of the pyrethroid tetramethrin on tetrodotoxin-sensitive and tetrodotoxin-resistant single sodium channels. *Brain Res.*, **712**: 258-264. [https://doi.org/10.1016/0006-8993\(95\)01449-7](https://doi.org/10.1016/0006-8993(95)01449-7)
- Stansly, P.A. and Natwick, E.T., 2010. Integrated systems for managing *Bemisia tabaci* in protected and open field agriculture. In: *Bemisia: Bionomics management of a global pest* (eds. P.A. Stansly and S.E. Naranjo). Springer, Dordrecht, pp. 467-497. https://doi.org/10.1007/978-90-481-2460-2_17
- Stenersen, J., 2004. *Chemical pesticides mode of action and toxicology*. CRC Press. <https://doi.org/10.1201/9780203646830>
- Veeravel, R. and Ravivarman, B., 2010. Bioefficacy evaluation of bifenthrin 10EC against major pests of rice. *Madras Agric. J.*, **97**: 164-167. <https://doi.org/10.29321/MAJ.10.100372>
- Vontas, J.G., Small, G.J., Nikou, D.C., Ranson, H. and Hemingway, J., 2002. Purification, molecular cloning and heterologous expression of a glutathione S-transferase involved in insecticide resistance from the rice brown planthopper, *Nilaparvata lugens*. *Biochem. J.*, **362**: 329-337. <https://doi.org/10.1042/bj3620329>
- Wang, K., Liu, T., Yu, C., Jiang, X. and Yi, M., 2002. Resistance of *Aphis gossypii* (Homoptera: Aphididae) to fenvalerate and imidacloprid and activities of detoxification enzymes on cotton and cucumber. *J. econ. Ent.*, **95**: 407-413. <https://doi.org/10.1603/0022-0493-95.2.407>
- Wang, S., Zhang, Y., Yang, X., Xie, W. and Wu, Q., 2017. Resistance monitoring for eight insecticides on the sweetpotato whitefly (Hemiptera: Aleyrodidae) in China. *J. econ. Ent.*, **110**: 660-666. <https://doi.org/10.1093/jee/tox040>
- Xie, W., Liu, Y., Wang, S., Wu, Q., Pan, H., Yang, X., Guo, L. and Zhang, Y., 2014. Sensitivity of *Bemisia tabaci* (Hemiptera: Aleyrodidae) to several new insecticides in China: effects of insecticide type and whitefly species, strain, and stage. *J. Insect Sci.*, **14**: 261-273. <https://doi.org/10.1093/jisesa/ieu123>
- Yang, X., He, C., Xie, W., Liu, Y., Xia, J., Yang, Z., Guo, L., Wen, Y., Wang, S. and Wu, Q., 2016. Glutathione S-transferases are involved in thiamethoxam resistance in the field whitefly *Bemisia tabaci* Q (Hemiptera: Aleyrodidae). *Pestic. Biochem. Physiol.*, **134**: 73-78. <https://doi.org/10.1016/j.pestbp.2016.04.003>
- Yu, S., 1988. Selectivity of insecticides to the spined soldier bug (Heteroptera: Pentatomidae) and its Lepidopterous prey. *J. econ. Ent.*, **81**: 119-122. <https://doi.org/10.1093/jee/81.1.119>
- Zhang, X., Liao, X., Mao, K., Zhang, K., Wan, H. and Li, J., 2016. Insecticide resistance monitoring and correlation analysis of insecticides in field populations of the brown planthopper *Nilaparvata lugens* (Stål) in China 2012–2014. *Pestic. Biochem. Physiol.*, **132**: 13-20. <https://doi.org/10.1016/j.pestbp.2015.10.003>
- Zheng, H., Xie, W., Wang, S., Wu, Q., Zhou, X. and Zhang, Y., 2017. Dynamic monitoring (B versus Q) and further resistance status of Q-type *Bemisia tabaci* in China. *Crop Prot.*, **94**: 115-122. <https://doi.org/10.1016/j.cropro.2016.11.035>
- Zibae, A., Bandani, A.R. and Tork, M., 2009. Effect of the entomopathogenic fungus, *Beauveria bassiana*, and its secondary metabolite on detoxifying enzyme activities and acetylcholinesterase (AChE) of the Sunn pest, *Eurygaster integriceps* (Heteroptera: Scutellaridae). *Biocont. Sci. Technol.*, **19**: 485-498. <https://doi.org/10.1080/09583150902847127>